

Cardiac Rehabilitation in the Outpatient Setting

Policy # 00570

Original Effective Date: 09/20/2017

Current Effective Date: 07/01/2026

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Investigational or experimental services are not covered. This includes any drug, device, procedure, or service provided under the investigational arm of a clinical trial or clinical study. These services are excluded from coverage under benefits.

When Services Are Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of available data, the Company may consider outpatient cardiac rehabilitation (CR) programs to be **eligible for coverage**** when **ALL** of the following criteria are met:

- History of **ANY** of the following conditions or procedures:
 - Acute myocardial infarction (heart attack) within the preceding 12 months; **OR**
 - Coronary artery bypass graft surgery; **OR**
 - Percutaneous transluminal coronary angioplasty or coronary stenting; **OR**
 - Heart valve surgery; **OR**
 - Heart or heart-lung transplantation; **OR**
 - Current stable angina pectoris unresponsive to optimal guideline-directed medical therapy; **OR**
 - Compensated heart failure with New York Heart Association (NYHA) class II to IV symptoms treated with optimal guideline-directed medical therapy; **AND**
- CR program starts within 90 days of the cardiac event and will be completed within 6 months of the cardiac event (with exception of acute myocardial infarction as noted above); **AND**
- The cardiac rehabilitation program includes **ALL** of the following components:
 - Physician-prescribed exercise each day cardiac rehabilitation services are provided; **AND**
 - Cardiac risk factor modification; **AND**
 - Psychosocial assessment; **AND**
 - Outcomes assessment; **AND**
 - Individualized treatment plan detailing how each of the above components are utilized; **AND**
- A cardiac rehabilitation exercise program is eligible for coverage for 3 sessions per week up to a 12-week period (36 sessions).

Cardiac Rehabilitation in the Outpatient Setting

Policy # 00570

Original Effective Date: 09/20/2017

Current Effective Date: 07/01/2026

Note:

A comprehensive evaluation may be performed before initiation of cardiac rehabilitation to evaluate the individual and determine an appropriate exercise program. In addition to a medical examination, an electrocardiogram stress test may be performed. An additional stress test may be performed at the completion of the program.

Physical and/or occupational therapy are not medically necessary in conjunction with cardiac rehabilitation unless performed for an unrelated diagnosis.

When Services Are Considered Not Medically Necessary

The use of cardiac rehabilitation in the outpatient setting when patient selection criteria are not met is considered to be **not medically necessary.****

When Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers repeat participation in an outpatient cardiac rehabilitation program in the absence of another qualifying cardiac event to be **investigational.***

Based on review of available data, the Company considers intensive cardiac rehabilitation with the Ornish Program for Reversing Heart Disease, Pritikin Program, or Benson-Henry Institute Program to be **investigational.***

Based on review of available data, the Company considers virtual cardiac rehabilitation to be **investigational.***

Based on review of available data, the Company considers outpatient cardiac rehabilitation for all other indications (e.g., SARS-CoV-2) to be **investigational.***

Background/Overview

Heart disease is the leading cause of mortality in the United States, accounting for more than half of all deaths. Coronary artery disease is the most common cause of heart disease. In a 2024 update on heart disease and stroke statistics from the American Heart Association, it was estimated that 720,000 Americans have a new coronary attack (first hospitalized myocardial infarction or coronary heart disease death) and 335,000 have a recurrent attack annually. Both coronary artery disease and various other disorders—structural heart disease and other genetic, metabolic, endocrine, toxic, inflammatory, and infectious causes—can lead to the clinical syndrome of heart failure, of which there are about 650,000 new cases in the United States annually. Given the burden of heart disease, preventing secondary cardiac events and treating the symptoms of heart disease and heart failure have received much attention from national organizations.

Policy # 00570

Original Effective Date: 09/20/2017

Current Effective Date: 07/01/2026

Stable Angina

Stable chest pain is a symptom of myocardial ischemia characterized by chest pain that is provoked with stress (physical or emotional). 2021 Report of the American College of Cardiology and American Heart Association Joint Committee on Clinical Practice Guidelines notes that anginal symptoms are perceived as retrosternal chest discomfort (e.g., pain, discomfort, heaviness, tightness, pressure, constriction, squeezing) that gradually build in intensity over a few minutes. Common symptoms associated with myocardial ischemia include, but are not limited to, dyspnea, palpitations, diaphoresis, lightheadedness, presyncope or syncope, upper abdominal pain, or heartburn unrelated to meals and nausea or vomiting.

For intermediate-high risk patients with stable chest pain and no known coronary artery disease (CAD), Computed Tomography Angiography (CCTA) is effective for diagnosis of CAD, risk stratification, and guiding treatment decisions (1A). Stress imaging (stress echocardiography, PET/SPECT myocardial perfusion imaging or cardiovascular magnetic resonance) is effective for diagnosis of myocardial ischemia and for estimating risk of major adverse cardiac event (1). For intermediate-high risk patients with stable chest pain and known coronary stenosis of 40-90% in a proximal or middle coronary segment on CCTA, fractional flow reserve (FFR)-CT can be useful for diagnosis of vessel-specific ischemia and to guide decision-making regarding the use of coronary revascularization (2a).

The term “obstructive” is used to indicate coronary artery disease (CAD) with $\geq 50\%$ stenosis and nonobstructive is used to indicate CAD with $< 50\%$ stenosis. “High risk CAD” is used to denote patients with obstructive stenosis who have left main stenosis $\geq 50\%$ or anatomically significant 3-vessel disease ($\geq 70\%$ stenosis).

For patients with obstructive CAD and stable chest pain, it is recommended to optimize guideline directed medical therapy (GDMT) including optimization of anti-ischemic and preventive therapies with the goal to reduce the patient’s angina burden and improve clinical outcomes (1A). For patients with known nonobstructive CAD and stable chest pain, it is recommended to optimize preventive therapies according to ACC/AHA clinical practice guidelines (1). Although GDMT exists for obstructive CAD, there are no current guidelines that are specific to nonobstructive CAD. Thus, adhering to atherosclerotic CV prevention guidelines is recommended.

Guideline-directed medical therapy (GDMT) consists of risk factor management and, in symptomatic patients, antianginal medications which improve quality of life.

- Risk factor management: All patients with stable CAD should be encouraged to adopt healthy lifestyles including tobacco cessation/avoidance, regular physical activity, maintenance of a healthy weight and adherence to a healthy diet. In addition, absent a contraindication, all stable CAD patients should be taking the following evidence-supported medications:
 - Antiplatelet agents – Aspirin and/or P2Y₁₂ receptor antagonist
 - Statin – Maximum tolerated dose of high-intensity statin (atorvastatin 40-80 mg or rosuvastatin 20-40 mg). Patients intolerant of statins and/or not reaching LDL

Cardiac Rehabilitation in the Outpatient Setting

Policy # 00570

Original Effective Date: 09/20/2017

Current Effective Date: 07/01/2026

- cholesterol goal on maximum tolerated statin dose should be treated with ezetimibe, a PCSK9 inhibitor, or bempedoic acid.
 - Beta blockers – In patients with a history of myocardial infarction, who have left ventricular systolic dysfunction (ejection fraction $\leq 40\%$), or as an option for management of hypertension.
 - ACE Inhibitor or Angiotensin Receptor Blocker – In patients with left ventricular systolic dysfunction (ejection fraction $\leq 40\%$), diabetes, chronic kidney disease, or as an option for management of hypertension
 - Antidiabetic agents – For patients who are diabetic (Hemoglobin A1c goal should be $< 8\%$ in all patients although more aggressive management may be appropriate for some)
- Symptom control: Most patients with stable CAD who have symptoms should be offered antianginal medications as an initial approach with revascularization reserved for those who have persistent unacceptable symptoms despite maximally tolerated doses.
 - Beta blockers – Unless contraindicated beta blockers are first-line therapy with dose escalation until symptoms resolve or side effects develop.
 - Calcium channel blockers and/or long acting-nitrates should be used as alternative initial therapy in symptomatic patients who have contraindication to, or intolerance of, beta blockers. They should also be prescribed when symptoms persist despite maximum tolerated doses of beta blockers.
 - Ranolazine may be prescribed either as initial therapy in symptomatic patients who have contraindication to, or intolerance of, other antianginal medication, or for those with persistent symptoms despite treatment with other medications as described above.

Heart failure

The NYHA classification is used to characterize symptoms and functional capacity of patients with symptomatic (stage C, NYHA class II and III) heart failure or advanced heart failure (stage D). A 2022 Report of the American College of Cardiology and American Heart Association Joint Committee on Clinical Practice Guidelines for the management of heart failure noted for stage C heart failure that exercise training in patients with HF is safe and has numerous benefits. In a major trial of exercise and HF, exercise training was associated with a reduction in CVD mortality or hospitalizations in the exercise training group after adjustment for risk factors. Meta-analyses show that cardiac rehabilitation improves functional capacity, exercise duration, and health-related QOL. A cardiac rehabilitation program for patients with HF usually includes a medical evaluation, education regarding the importance of medical adherence, dietary recommendations, psychosocial support, and an exercise training and physical activity counseling program. Patients with HF on optimal GDMT, who are in stable medical condition and are able to participate in an exercise program, are candidates for an exercise rehabilitation program. In a diverse population of older patients who were hospitalized for acute decompensated HF, an early, transitional, tailored, progressive rehabilitation intervention that included multiple physical-function domains (strength, balance, mobility, and endurance) initiated during, or early after hospitalization for HF, and continued after discharge, resulted in greater improvement in physical function than usual care.

Cardiac Rehabilitation in the Outpatient Setting

Policy # 00570

Original Effective Date: 09/20/2017

Current Effective Date: 07/01/2026

Cardiac Rehabilitation

In 1995, the U.S. Public Health Service defined cardiac rehabilitation services as, in part, “comprehensive, long-term programs involving medical evaluation, prescribed exercise, cardiac risk factor modification, education, and counseling.... [These programs] are designed to limit the physiologic and psychological effects of cardiac illness, reduce the risk for sudden death or reinfarction, control cardiac symptoms, stabilize or reverse the atherosclerotic process, and enhance the psychosocial and vocational status of selected patients.” The U.S. Public Health Service recommended cardiac rehabilitation services for patients with coronary heart disease and heart failure, including those awaiting or following cardiac transplantation. A 2010 definition of cardiac rehabilitation from the European Association of Cardiovascular Prevention and Rehabilitation stated: “Cardiac rehabilitation can be viewed as the clinical application of preventive care by means of a professional multi-disciplinary integrated approach for comprehensive risk reduction and global long-term care of cardiac patients.” Since the 1995 release of the U.S. Public Health Service guidelines, other societies, including in 2005 the American Heart Association⁴ and in 2010 the Heart Failure Society of America have developed guidelines on the role of cardiac rehabilitation in patient care.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

Not applicable.

Rationale/Source

This medical policy was developed through consideration of peer-reviewed medical literature generally recognized by the relevant medical community, U.S. Food and Drug Administration approval status, nationally accepted standards of medical practice and accepted standards of medical practice in this community, technology evaluation centers, reference to regulations, other plan medical policies, and accredited national guidelines.

Description

Cardiac rehabilitation refers to comprehensive medically supervised programs in the outpatient setting that aim to improve the function of patients with heart disease and prevent future cardiac events. National organizations have specified core components to be included in cardiac rehabilitation programs.

Summary of Evidence

For individuals who have been diagnosed with heart disease and receive outpatient cardiac rehabilitation, the evidence includes multiple randomized controlled trials (RCTs) and systematic reviews of these trials. Relevant outcomes are overall survival (OS), disease-specific survival, symptoms, and morbid events. Meta-analyses of the available trials have found that cardiac rehabilitation improves health outcomes for select patients, particularly those with coronary heart disease, heart failure, and who have had cardiac surgical interventions. The available evidence has limitations, including lack of blinded outcome assessment, but for the survival-related outcomes of

Cardiac Rehabilitation in the Outpatient Setting

Policy # 00570

Original Effective Date: 09/20/2017

Current Effective Date: 07/01/2026

interest, this limitation is less critical. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have been diagnosed with heart disease without a second event and receive repeat outpatient cardiac rehabilitation, the evidence includes no trials. Relevant outcomes are OS, disease-specific survival, symptoms, and morbid events. No studies were identified evaluating the effectiveness of repeat participation in a cardiac rehabilitation program. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have been diagnosed with heart disease and receive intensive cardiac rehabilitation with the Ornish Program for Reversing Heart Disease, the evidence includes an RCT. Relevant outcomes are OS, disease-specific survival, symptoms, and morbid events. No RCTs have compared the Ornish Program with a “standard” cardiac rehabilitation program; an RCT compared it with usual care. The trial included patients with coronary artery disease and no recent cardiac events and had mixed findings at 1 and 5 years. The trial had a small sample size for a cardiac trial (N=48), and only 35 patients were available for the 5-year follow-up. The Ornish Program is considered by the Centers for Medicare & Medicaid Services as an intensive cardiac rehabilitation program, but the program described in the RCT could meet criteria for standard cardiac rehabilitation. No studies were identified comparing the Ornish Program with any other cardiac rehabilitation program. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have been diagnosed with heart disease and receive intensive cardiac rehabilitation with the Pritikin Program, the evidence includes 2 nonrandomized studies. Relevant outcomes are OS, disease-specific survival, symptoms, and morbid events. Studies are needed that compare the impact of intensive cardiac rehabilitation using the Pritikin Program with standard outpatient cardiac rehabilitation programs for these outcomes. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have been diagnosed with heart disease and receive intensive cardiac rehabilitation with the Benson-Henry Institute Program, the evidence includes a case-control study and case series. Relevant outcomes are OS, disease-specific survival, symptoms, and morbid events. Studies are needed that compare the impact of intensive cardiac rehabilitation using the Benson-Henry Institute Program with standard outpatient cardiac rehabilitation programs. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have heart disease due to post-acute sequelae of SARS-CoV-2 infection who receive cardiac rehabilitation in the outpatient setting, no relevant evidence was identified. Relevant outcomes are OS, disease-specific survival, symptoms, and morbid events. Reports examining the outcomes of rehabilitation in patients with post-acute COVID-19 have not primarily focused on cardiac rehabilitation. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Cardiac Rehabilitation in the Outpatient Setting

Policy # 00570

Original Effective Date: 09/20/2017

Current Effective Date: 07/01/2026

For individuals who have been diagnosed with heart disease and receive virtual cardiac rehabilitation, the evidence includes systematic reviews/meta-analyses, RCTs, and observational studies. Relevant outcomes are OS, disease-specific survival, symptoms, and morbid events. Meta-analyses have found beneficial effects of virtual cardiac rehabilitation on physical activity and quality of life, but not on cardiovascular hospitalization or mortality. The few available prospective randomized studies have conflicting findings on the effect of virtual cardiac rehabilitation compared to traditional outpatient cardiac rehabilitation on hospital readmission. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Supplemental Information

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American College of Cardiology Foundation/American Heart Association

The 2022 American College of Cardiology (ACC) and the American Heart Association (AHA) heart failure guidelines recommend rehabilitation for Stage C heart failure stating, "In patients with HF, a cardiac rehabilitation program can be useful to improve functional capacity, exercise tolerance, and health-related QOL." In 2023, the ACC/AHA published a statement on supervised exercise training specific to patients with chronic heart failure with preserved ejection fraction (HFpEF) and concluded, "data reviewed herein demonstrate a comparable or larger magnitude of improvement in exercise capacity from supervised exercise training in patients with chronic HFpEF compared with those with heart failure with reduced ejection fraction."

American Heart Association

In 2024, the AHA and the American Association of Cardiovascular and Pulmonary Rehabilitation issued an updated consensus statement on the core components of cardiac rehabilitation programs. The core components included patient assessment before beginning the program, nutritional counseling, weight management and body composition, blood pressure management, lipid management, diabetes management, tobacco cessation, psychosocial management, aerobic exercise training, strength training, physical activity counseling, and program quality. Programs that only offered supervised exercise training were not considered cardiac rehabilitation. The guidelines specified the assessment, interventions, and expected outcomes for each of the core components. For example, symptom-limited exercise testing before exercise training was strongly recommended. The guidelines did not specify the optimal overall length of programs or the number or duration of sessions.

In 2019, the AHA, with the American Association of Cardiovascular and Pulmonary Rehabilitation and the ACC, released a scientific statement on home-based cardiac rehabilitation (HBCR). They make the following suggestions for healthcare providers:

Cardiac Rehabilitation in the Outpatient Setting

Policy # 00570

Original Effective Date: 09/20/2017

Current Effective Date: 07/01/2026

- Recommend center-based cardiac rehabilitation (CBCR) to all eligible patients.
- As an alternative, recommend HBCR to clinically stable low- and moderate-risk patients who cannot attend CBCR.
- Design and test HBCR “using effective processes of care for CVD [cardiovascular disease] secondary prevention.”
- For healthcare organizations, develop and support the following:
 - Maximization of cardiac rehabilitation (CR) referrals
 - High-quality CBCR and HBCR programs “using evidence-based standards and guidelines, strategies to maximize patient adherence both in the shorter and longer-term, and outcome tracking methods to help promote continuous quality improvement.”
 - “Testing and implementation of an evidence-based hybrid approach to CR” that are optimized for each patient and that “promote long-term adherence and favorable behavior change.”
- For CR professionals, “work with other healthcare professionals and policymakers to implement additional research and...expand the evidence base for HBCR.”

The guideline does not use the terminology “virtual” cardiac rehabilitation, but it states that electronic tools such as text messaging, smartphone applications, and wearable sensors may allow patients to follow personalized recommendations for exercise, dietary, and behavioral interventions, and thus expand the number of patients who can participate in cardiac rehabilitation. Other benefits of technology-assisted HBCR include greater patient engagement and patient-provider communication. The panel stated that studies were needed regarding the effect of technology-assisted HBCR on outcomes.

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

Cardiac Rehabilitation

Since 1989, Medicare has had a national coverage determination (NCD) for cardiac rehabilitation. The NCD was retired in April 2023. CMS periodically retires NCDs that no longer contain clinically pertinent and/or current information or no longer reflect current medical practice. In the absence of NCDs, coverage determinations are made by the Medicare Administrative Contractors (MACs) under section 1862(a)(1)(A) of the Social Security Act.⁵⁷

In October 2020, virtual cardiac rehabilitation and intensive cardiac rehabilitation were added to the list of telehealth services that Medicare would cover during the COVID-19 public health emergency. Virtual cardiac rehabilitation was only covered through the end of 2024.

Intensive Cardiac Rehabilitation

In January 2010, Medicare added intensive cardiac rehabilitation as a benefit. Intensive cardiac rehabilitation programs must be approved by Medicare on an individual basis.

Cardiac Rehabilitation in the Outpatient Setting

Policy # 00570

Original Effective Date: 09/20/2017

Current Effective Date: 07/01/2026

The national coverage determination described intensive cardiac rehabilitation in the following manner:

“Intensive cardiac rehabilitation (ICR) refers to a physician-supervised program that furnishes cardiac rehabilitation services more frequently and often in a more rigorous manner. As required by §1861(eee)(4)(A) of the Social Security Act (the Act), an ICR program must show, in peer-reviewed published research, that it accomplished 1 or more of the following for its patients: (1) positively affected the progression of coronary heart disease; (2) reduced the need for coronary bypass surgery; and, (3) reduced the need for percutaneous coronary interventions. The ICR program must also demonstrate through peer-reviewed published research that it accomplished a statistically significant reduction in 5 or more of the following measures for patients from their levels before cardiac rehabilitation services to after cardiac rehabilitation services: (1) low density lipoprotein; (2) triglycerides; (3) body mass index; (4) systolic blood pressure; (5) diastolic blood pressure; and, (6) the need for cholesterol, blood pressure, and diabetes medications. Individual ICR programs must be approved through the national coverage determination process to ensure that they demonstrate these accomplishments.”

In 2010, the Centers for Medicare & Medicaid Services also issued 2 decision memos on specific programs. One stated that the Ornish Program for Reversing Heart Disease met the intensive cardiac rehabilitation program requirements and was included on the list of approved intensive cardiac rehabilitation programs. It provided the following description of the Ornish Program:

“The Ornish Program for Reversing Heart Disease (also known as the Multisite Cardiac Lifestyle Intervention Program, Multicenter Cardiac Lifestyle Intervention Program and the Lifestyle Heart Trial program) ... incorporates comprehensive lifestyle modifications including exercise, a low-fat diet, smoking cessation, stress management training, and group support sessions. Over the years, the Ornish program has been refined but continues to focus on these specific risk factors.”

The other stated that the Pritikin Program met program requirements and was included on the list of approved intensive cardiac rehabilitation programs. As described in the decision memo:

“The Pritikin program (also known as the Pritikin Longevity Program) evolved into a comprehensive program that is provided in a physician’s office and incorporates a specific diet (10% to 15% of calories from fat, 15% to 20% from protein, 65% to 75% from complex carbohydrates), exercise and counseling lasting 21 to 26 days. An optional residential component is also available for participants.”

In 2014, Centers of Medicare & Medicaid Services issued another decision memo on the Benson-Henry Institute Cardiac Wellness Program. The memo stated that "the evidence is sufficient to expand the intensive care rehabilitation benefit to include the Benson-Henry Institute Cardiac Wellness Program. The Cardiac Wellness Program is a multicomponent intervention program that includes supervised exercise, behavioral interventions, and counseling, and is designed to reduce cardiovascular risk and improve health outcomes."

Cardiac Rehabilitation in the Outpatient Setting

Policy # 00570

Original Effective Date: 09/20/2017

Current Effective Date: 07/01/2026

Ongoing and Unpublished Clinical Trials

Some currently ongoing and unpublished trials that might influence this review are listed in Table 1.

Table 1. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT06077201	Home-Based Cardiac Rehabilitation Using a Novel Mobile Health Exercise Regimen Following Transcatheter Heart Valve Interventions	375	Oct 2026
NCT05933083	MCNAIR Study: comparative effectiveness of in-person and telehealth Cardiac Rehabilitation	516	Oct 2027
NCT05972070 ^a	Integration of Telemedicine and Home-Based Cardiac Rehabilitation: Feasibility, Efficacy, and Adherence	500	Dec 2025
NCT04245813	Effectiveness of a Cardiac Rehabilitation Program in Patients With Heart Failure	144	May 2023 (unknown status)
NCT02984449	Preventive Heart Rehabilitation in Patients Undergoing Elective Open Heart Surgery to Prevent Complications and to Improve Quality of Life (Heart-ROCQ) - A Prospective Randomized Open Controlled Trial, Blinded End-point (PROBE)	350	Aug 2025
NCT05689385	The Effectiveness of eHealth-based Cardiac Rehabilitation in Post-myocardial Infarction Patients: a Randomized Controlled Trial	150	Dec 2028

NCT: national clinical trial.

^a Indicates industry sponsored or cosponsored trial.

References

1. Martin SS, Aday AW, Almarzooq ZI, et al. 2024 Heart Disease and Stroke Statistics: A Report of US and Global Data From the American Heart Association. *Circulation*. Feb 20 2024; 149(8): e347-e913. PMID 38264914
2. Balady GJ, Williams MA, Ades PA, et al. Core components of cardiac rehabilitation/secondary prevention programs: 2007 update: a scientific statement from the American Heart Association Exercise, Cardiac Rehabilitation, and Prevention Committee, the Council on Clinical Cardiology; the Councils on Cardiovascular Nursing, Epidemiology and Prevention, and

Cardiac Rehabilitation in the Outpatient Setting

Policy # 00570

Original Effective Date: 09/20/2017

Current Effective Date: 07/01/2026

- Nutrition, Physical Activity, and Metabolism; and the American Association of Cardiovascular and Pulmonary Rehabilitation. *Circulation*. May 22 2007; 115(20): 2675-82. PMID 17513578
3. Corrà U, Piepoli MF, Carré F, et al. Secondary prevention through cardiac rehabilitation: physical activity counselling and exercise training: key components of the position paper from the Cardiac Rehabilitation Section of the European Association of Cardiovascular Prevention and Rehabilitation. *Eur Heart J*. Aug 2010; 31(16): 1967-74. PMID 20643803
 4. Leon AS, Franklin BA, Costa F, et al. Cardiac rehabilitation and secondary prevention of coronary heart disease: an American Heart Association scientific statement from the Council on Clinical Cardiology (Subcommittee on Exercise, Cardiac Rehabilitation, and Prevention) and the Council on Nutrition, Physical Activity, and Metabolism (Subcommittee on Physical Activity), in collaboration with the American association of Cardiovascular and Pulmonary Rehabilitation. *Circulation*. Jan 25 2005; 111(3): 369-76. PMID 15668354
 5. Lindenfeld J, Albert NM, Boehmer JP, et al. HFSA 2010 Comprehensive Heart Failure Practice Guideline. *J Card Fail*. Jun 2010; 16(6): e1-194. PMID 20610207
 6. Oldridge N. Exercise-based cardiac rehabilitation in patients with coronary heart disease: meta-analysis outcomes revisited. *Future Cardiol*. Sep 2012; 8(5): 729-51. PMID 23013125
 7. Anderson L, Thompson DR, Oldridge N, et al. Exercise-based cardiac rehabilitation for coronary heart disease. *Cochrane Database Syst Rev*. Jan 05 2016; 2016(1): CD001800. PMID 26730878
 8. Davies EJ, Moxham T, Rees K, et al. Exercise based rehabilitation for heart failure. *Cochrane Database Syst Rev*. Apr 14 2010; (4): CD003331. PMID 20393935
 9. Heran BS, Chen JM, Ebrahim S, et al. Exercise-based cardiac rehabilitation for coronary heart disease. *Cochrane Database Syst Rev*. Jul 06 2011; (7): CD001800. PMID 21735386
 10. Dibben G, Faulkner J, Oldridge N, et al. Exercise-based cardiac rehabilitation for coronary heart disease. *Cochrane Database Syst Rev*. Nov 06 2021; 11(11): CD001800. PMID 34741536
 11. Long L, Mordi IR, Bridges C, et al. Exercise-based cardiac rehabilitation for adults with heart failure. *Cochrane Database Syst Rev*. Jan 29 2019; 1(1): CD003331. PMID 30695817
 12. Molloy CD, Long L, Mordi IR, et al. Exercise-based cardiac rehabilitation for adults with heart failure - 2023 Cochrane systematic review and meta-analysis. *Eur J Heart Fail*. Dec 2023; 25(12): 2263-2273. PMID 37850321
 13. Molloy C, Long L, Mordi IR, et al. Exercise-based cardiac rehabilitation for adults with heart failure. *Cochrane Database Syst Rev*. Mar 07 2024; 3(3): CD003331. PMID 38451843
 14. West RR, Jones DA, Henderson AH. Rehabilitation after myocardial infarction trial (RAMIT): multi-centre randomised controlled trial of comprehensive cardiac rehabilitation in patients following acute myocardial infarction. *Heart*. Apr 2012; 98(8): 637-44. PMID 22194152
 15. Doherty P, Lewin R. The RAMIT trial, a pragmatic RCT of cardiac rehabilitation versus usual care: what does it tell us?. *Heart*. Apr 2012; 98(8): 605-6. PMID 22505460
 16. Pandey A, Kitzman DW, Brubaker P, et al. Response to Endurance Exercise Training in Older Adults with Heart Failure with Preserved or Reduced Ejection Fraction. *J Am Geriatr Soc*. Aug 2017; 65(8): 1698-1704. PMID 28338229
 17. Kitzman DW, Brubaker PH, Morgan TM, et al. Exercise training in older patients with heart failure and preserved ejection fraction: a randomized, controlled, single-blind trial. *Circ Heart Fail*. Nov 2010; 3(6): 659-67. PMID 20852060

Cardiac Rehabilitation in the Outpatient Setting

Policy # 00570

Original Effective Date: 09/20/2017

Current Effective Date: 07/01/2026

18. Opotowsky AR, Rhodes J, Landzberg MJ, et al. A Randomized Trial Comparing Cardiac Rehabilitation to Standard of Care for Adults With Congenital Heart Disease. *World J Pediatr Congenit Heart Surg.* Mar 2018; 9(2): 185-193. PMID 29544423
19. Sumner J, Harrison A, Doherty P. The effectiveness of modern cardiac rehabilitation: A systematic review of recent observational studies in non-attenders versus attenders. *PLoS One.* 2017; 12(5): e0177658. PMID 28498869
20. Nilsson BB, Lunde P, Grøgaard HK, et al. Long-Term Results of High-Intensity Exercise-Based Cardiac Rehabilitation in Revascularized Patients for Symptomatic Coronary Artery Disease. *Am J Cardiol.* Jan 01 2018; 121(1): 21-26. PMID 29096886
21. Jafri SH, Imran TF, Medbury E, et al. Cardiovascular Outcomes of Patients Referred to Home Based Cardiac Rehabilitation. *Heart Lung.* 2022; 52: 1-7. PMID 34801771
22. Ornish D, Brown SE, Scherwitz LW, et al. Can lifestyle changes reverse coronary heart disease? The Lifestyle Heart Trial. *Lancet.* Jul 21 1990; 336(8708): 129-33. PMID 1973470
23. Ornish D, Scherwitz LW, Billings JH, et al. Intensive lifestyle changes for reversal of coronary heart disease. *JAMA.* Dec 16 1998; 280(23): 2001-7. PMID 9863851
24. Lakhani F, Racette SB, Park LK, et al. Prospective Study of the Impact of Outpatient Intensive Cardiac Rehabilitation on Diet Quality, Health-related Quality of Life, and Cardiovascular Health Indices. *Am J Cardiol.* Apr 01 2023; 192: 60-66. PMID 36736014
25. Racette SB, Park LK, Rashdi ST, et al. Benefits of the First Pritikin Outpatient Intensive Cardiac Rehabilitation Program. *J Cardiopulm Rehabil Prev.* Nov 01 2022; 42(6): 449-455. PMID 35861951
26. Zeng W, Stason WB, Fournier S, et al. Benefits and costs of intensive lifestyle modification programs for symptomatic coronary disease in Medicare beneficiaries. *Am Heart J.* May 2013; 165(5): 785-92. PMID 23622916
27. Casey A, Chang BH, Huddleston J, et al. A model for integrating a mind/body approach to cardiac rehabilitation: outcomes and correlators. *J Cardiopulm Rehabil Prev.* 2009; 29(4): 230-8; quiz 239-40. PMID 19451830
28. Centers for Disease Control and Prevention (CDC). Clinical Overview of Long COVID. July 24, 2025; <https://www.cdc.gov/covid/hcp/clinical-overview/index.html>.
29. Soriano JB, Murthy S, Marshall JC, et al. A clinical case definition of post-COVID-19 condition by a Delphi consensus. *Lancet Infect Dis.* Apr 2022; 22(4): e102-e107. PMID 34951953
30. Jin Choo Y, Chang MC. Effects of telecardiac rehabilitation on coronary heart disease: A PRISMA-compliant systematic review and meta-analysis. *Medicine (Baltimore).* Jul 15 2022; 101(28): e29459. PMID 35839029
31. Nacarato D, Sardeli AV, Mariano LO, et al. Cardiovascular telerehabilitation improves functional capacity, cardiorespiratory fitness and quality of life in older adults: A systematic review and meta-analysis. *J Telemed Telecare.* Sep 2024; 30(8): 1238-1248. PMID 36469017
32. Cavalheiro AH, Silva Cardoso J, Rocha A, et al. Effectiveness of Tele-rehabilitation Programs in Heart Failure: A Systematic Review and Meta-analysis. *Health Serv Insights.* 2021; 14: 11786329211021668. PMID 34188484
33. Cruz-Cobo C, Bernal-Jiménez MÁ, Vázquez-García R, et al. Effectiveness of mHealth Interventions in the Control of Lifestyle and Cardiovascular Risk Factors in Patients After a Coronary Event: Systematic Review and Meta-analysis. *JMIR Mhealth Uhealth.* Dec 02 2022; 10(12): e39593. PMID 36459396

Cardiac Rehabilitation in the Outpatient Setting

Policy # 00570

Original Effective Date: 09/20/2017

Current Effective Date: 07/01/2026

34. Maulana S, Trisyani Y, Mirwanti R, et al. The Potential of Cardiac Telerehabilitation as Delivery Rehabilitation Care Model in Heart Failure during COVID-19 and Transmissible Disease Outbreak: A Systematic Scoping Review of the Latest RCTs. *Medicina (Kaunas)*. Sep 21 2022; 58(10). PMID 36295482
35. Ramachandran HJ, Jiang Y, Tam WWS, et al. Effectiveness of home-based cardiac telerehabilitation as an alternative to Phase 2 cardiac rehabilitation of coronary heart disease: a systematic review and meta-analysis. *Eur J Prev Cardiol*. May 25 2022; 29(7): 1017-1043. PMID 34254118
36. Pagliari C, Isernia S, Rapisarda L, et al. Different Models of Cardiac Telerehabilitation for People with Coronary Artery Disease: Features and Effectiveness: A Systematic Review and Meta-Analysis. *J Clin Med*. Jun 10 2024; 13(12). PMID 38929925
37. McDonagh ST, Dalal H, Moore S, et al. Home-based versus centre-based cardiac rehabilitation. *Cochrane Database Syst Rev*. Oct 27 2023; 10(10): CD007130. PMID 37888805
38. Zhou M, Xu Y, Zhang L, et al. Effectiveness of smartphone-assisted cardiac rehabilitation: a systematic review and meta-analysis. *Disabil Rehabil*. Jul 2024; 46(15): 3256-3265. PMID 37559408
39. Li L, Ringeval M, Wagner G, et al. Effectiveness of home-based cardiac rehabilitation interventions delivered via mHealth technologies: a systematic review and meta-analysis. *Lancet Digit Health*. Apr 2025; 7(4): e238-e254. PMID 40023729
40. Nagatomi Y, Ide T, Higuchi T, et al. Home-based cardiac rehabilitation using information and communication technology for heart failure patients with frailty. *ESC Heart Fail*. Aug 2022; 9(4): 2407-2418. PMID 35534907
41. Brouwers RWM, Kemps HMC, Herkert C, et al. A 12-week cardiac telerehabilitation programme does not prevent relapse of physical activity levels: long-term results of the FIT@Home trial. *Eur J Prev Cardiol*. May 25 2022; 29(7): e255-e257. PMID 35040993
42. Brouwers RWM, Kraal JJ, Regis M, et al. Effectiveness of Cardiac Telerehabilitation With Relapse Prevention: SmartCare-CAD Randomized Controlled Trial. *J Am Coll Cardiol*. Jun 01 2021; 77(21): 2754-2756. PMID 34045031
43. Indraratna P, Biswas U, McVeigh J, et al. A Smartphone-Based Model of Care to Support Patients With Cardiac Disease Transitioning From Hospital to the Community (TeleClinical Care): Pilot Randomized Controlled Trial. *JMIR Mhealth Uhealth*. Feb 28 2022; 10(2): e32554. PMID 35225819
44. Snoek JA, Prescott EI, van der Velde AE, et al. Effectiveness of Home-Based Mobile Guided Cardiac Rehabilitation as Alternative Strategy for Nonparticipation in Clinic-Based Cardiac Rehabilitation Among Elderly Patients in Europe: A Randomized Clinical Trial. *JAMA Cardiol*. Apr 01 2021; 6(4): 463-468. PMID 33112363
45. Piotrowicz E, Pencina MJ, Opolski G, et al. Effects of a 9-Week Hybrid Comprehensive Telerehabilitation Program on Long-term Outcomes in Patients With Heart Failure: The Telerehabilitation in Heart Failure Patients (TELEREH-HF) Randomized Clinical Trial. *JAMA Cardiol*. Mar 01 2020; 5(3): 300-308. PMID 31734701
46. Yudi MB, Clark DJ, Tsang D, et al. SMARTphone-based, early cardiac REHABilitation in patients with acute coronary syndromes: a randomized controlled trial. *Coron Artery Dis*. Aug 01 2021; 32(5): 432-440. PMID 32868661

Cardiac Rehabilitation in the Outpatient Setting

Policy # 00570

Original Effective Date: 09/20/2017

Current Effective Date: 07/01/2026

47. Hakala S, Kivistö H, Paajanen T, et al. Effectiveness of Distance Technology in Promoting Physical Activity in Cardiovascular Disease Rehabilitation: Cluster Randomized Controlled Trial, A Pilot Study. *JMIR Rehabil Assist Technol*. Jun 18 2021; 8(2): e20299. PMID 34142970
48. Dalli Peydró E, Sanz Sevilla N, Tuzón Segarra MT, et al. A randomized controlled clinical trial of cardiac telerehabilitation with a prolonged mobile care monitoring strategy after an acute coronary syndrome. *Clin Cardiol*. Jan 2022; 45(1): 31-41. PMID 34952989
49. Maddison R, Rawstorn JC, Stewart RAH, et al. Effects and costs of real-time cardiac telerehabilitation: randomised controlled non-inferiority trial. *Heart*. Jan 2019; 105(2): 122-129. PMID 30150328
50. Lear SA, Singer J, Banner-Lukaris D, et al. Randomized trial of a virtual cardiac rehabilitation program delivered at a distance via the Internet. *Circ Cardiovasc Qual Outcomes*. Nov 2014; 7(6): 952-9. PMID 25271050
51. Dodson JA, Adhikari S, Schoenthaler A, et al. Rehabilitation at Home Using Mobile Health for Older Adults Hospitalized for Ischemic Heart Disease: The RESILIENT Randomized Clinical Trial. *JAMA Netw Open*. Jan 02 2025; 8(1): e2453499. PMID 39775808
52. Schmidt C, Magalhães S, Gois Basilio P, et al. Center- vs Home-Based Cardiac Rehabilitation in Patients With Heart Failure: EXIT-HF Randomized Controlled Trial. *JACC Heart Fail*. May 2025; 13(5): 695-706. PMID 39895436
53. Nkonde-Price C, Reynolds K, Najem M, et al. Comparison of Home-Based vs Center-Based Cardiac Rehabilitation in Hospitalization, Medication Adherence, and Risk Factor Control Among Patients With Cardiovascular Disease. *JAMA Netw Open*. Aug 01 2022; 5(8): e2228720. PMID 36006642
54. Shah ND, Banta CW, Berger AL, et al. Retrospective Comparison of Outcomes and Cost of Virtual Versus Center-Based Cardiac Rehabilitation Programs. *J Am Heart Assoc*. Dec 17 2024; 13(24): e036861. PMID 39673292
55. Wang Z, Elkoustaf R, Batiste C, et al. Home based cardiac rehabilitation: A retrospective cohort analysis on all-cause mortality and hospital readmission rates across sexes and races. *Am J Prev Cardiol*. Sep 2024; 19: 100708. PMID 39157645
56. Heidenreich PA, Bozkurt B, Aguilar D, et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol*. May 03 2022; 79(17): e263-e421. PMID 35379503
57. Sachdev V, Sharma K, Keteyian SJ, et al. Supervised Exercise Training for Chronic Heart Failure With Preserved Ejection Fraction: A Scientific Statement From the American Heart Association and American College of Cardiology. *J Am Coll Cardiol*. Apr 18 2023; 81(15): 1524-1542. PMID 36958952
58. Brown TM, Pack QR, Aberegg E, et al. Core Components of Cardiac Rehabilitation Programs: 2024 Update: A Scientific Statement From the American Heart Association and the American Association of Cardiovascular and Pulmonary Rehabilitation. *Circulation*. Oct 29 2024; 150(18): e328-e347. PMID 39315436
59. Thomas RJ, Beatty AL, Beckie TM, et al. Home-Based Cardiac Rehabilitation: A Scientific Statement From the American Association of Cardiovascular and Pulmonary Rehabilitation, the American Heart Association, and the American College of Cardiology. *J Am Coll Cardiol*. Jul 09 2019; 74(1): 133-153. PMID 31097258

Cardiac Rehabilitation in the Outpatient Setting

Policy # 00570

Original Effective Date: 09/20/2017

Current Effective Date: 07/01/2026

60. Centers for Medicare & Medicaid Services (CMS). National Coverage Determination (NCD) for Intensive Cardiac Rehabilitation Programs (20.31). 2010; <https://www.cms.gov/medicare-coverage-database/view/ncd.aspx?NCID=339&ncdver=1&fromdb=true>.
61. Centers for Medicare and Medicaid Services. Trump Administration Drives Telehealth Services in Medicaid and Medicare. October 14, 2020. <https://www.cms.gov/newsroom/press-releases/trump-administration-drives-telehealth-services-medicare-and-medicare>.
62. Centers for Medicare and Medicaid Services. List of Telehealth Services. Updated December 4, 2025. <https://www.cms.gov/Medicare/Medicare-General-Information/Telehealth/Telehealth-Codes>.
63. Centers for Medicare and Medicaid Services. Billing and Coding: Outpatient Cardiac Rehabilitation. Retired November 1, 2023. https://localcoverage.cms.gov/mcd_archive/view/article.aspx?articleInfo=54070:43&.
64. Centers for Medicare & Medicaid Services (CMS). Decision Memo for Intensive Cardiac Rehabilitation (ICR) Program - Dr. Ornish's Program for Reversing Heart Disease (CAG-00419N). 2010; <https://www.cms.gov/medicare-coverage-database/view/ncacal-decision-memo.aspx?proposed=N&NCAId=240&NcaName=Intensive+Cardiac+Rehabilitation+%28ICR%29+Program+-+Dr.+Ornish>.
65. Centers for Medicare & Medicaid Services (CMS). Decision Memo for Intensive Cardiac Rehabilitation (ICR) Program - Pritikin Program (CAG-00418N). 2010; <https://www.cms.gov/medicare-coverage-database/view/ncacal-decision-memo.aspx?proposed=N&NCAId=239&bc=AiAAAAAAAgAAAA%>.
66. Centers for Medicare and Medicaid Services. Decision Memo: Intensive Cardiac Rehabilitation (ICR) Program - Benson-Henry Institute Cardiac Wellness Program. May 6, 2014. <https://www.cms.gov/medicare-coverage-database/view/ncacal-decision-memo.aspx?proposed=N&ncid=271>.
67. National Institute for Health and Clinical Excellence (NICE). Recent-onset chest pain of suspected cardiac origin: assessment and diagnosis. NICE Clinical Guideline 95. London, UK: NICE; May 2010.

Policy History

Original Effective Date: 09/20/2017

Current Effective Date: 07/01/2026

09/07/2017	Medical Policy Committee review
09/20/2017	Medical Policy Implementation Committee approval. New policy.
09/06/2018	Medical Policy Committee review
09/19/2018	Medical Policy Implementation Committee approval. No change to coverage.
09/05/2019	Medical Policy Committee review
09/11/2019	Medical Policy Implementation Committee approval. No change to coverage.
09/03/2020	Medical Policy Committee review
09/09/2020	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
09/02/2021	Medical Policy Committee review

Cardiac Rehabilitation in the Outpatient Setting

Policy # 00570

Original Effective Date: 09/20/2017

Current Effective Date: 07/01/2026

09/08/2021 Medical Policy Implementation Committee approval. Added PICO and investigational policy statement for intensive cardiac rehabilitation with the Benson-Henry Institute Program.

09/01/2022 Medical Policy Committee review

09/14/2022 Medical Policy Implementation Committee approval. No change to coverage.

05/04/2023 Medical Policy Committee review

05/10/2023 Medical Policy Implementation Committee approval. Added “Based on review of available data, the Company considers virtual cardiac rehabilitation to be investigational.”

05/02/2024 Medical Policy Committee review

05/08/2024 Medical Policy Implementation Committee approval. Investigational policy statement added for all other indications, including post-acute sequelae of SARS-CoV-2 infection.

06/05/2025 Medical Policy Committee review

06/11/2025 Medical Policy Implementation Committee approval. Coverage eligibility unchanged. Two new MN statements added for Chronic and Stable Angina. Section on stable angina added to background policy for support.

06/04/2026 Medical Policy Committee review

06/10/2026 Medical Policy Implementation Committee review

08/20/2026 Medical Quality Management Committee approval. The following statement was changed from investigational to not medically necessary: “The use of cardiac rehabilitation in the outpatient setting when patient selection criteria are not met is considered to be not medically necessary.”

Next Scheduled Review Date: 06/2027

Coding

The five character codes included in the Louisiana Blue Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)†, copyright 2025 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Louisiana Blue Medical Policy Coverage Guidelines is with Louisiana Blue and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Louisiana Blue Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Louisiana Blue Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

Cardiac Rehabilitation in the Outpatient Setting

Policy # 00570

Original Effective Date: 09/20/2017

Current Effective Date: 07/01/2026

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	93797, 93798
HCPCS	G0422, G0423, S9472
ICD-10 Diagnosis	All related diagnoses

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
 1. Consultation with technology evaluation center(s);
 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
 3. Reference to federal regulations.

**Medically Necessary (or “Medical Necessity”) - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

For these purposes, “nationally accepted standards of medical practice” means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

Cardiac Rehabilitation in the Outpatient Setting

Policy # 00570

Original Effective Date: 09/20/2017

Current Effective Date: 07/01/2026

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: If the Patient's health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

NOTICE: Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage.

NOTICE: If an authorization for an ongoing course of treatment has been provided to a member and the member changes from one health plan to another health plan (e.g., a member moves from carrier A to Louisiana Blue), Louisiana Blue may honor the previous health plan's authorization for the same service under the same type of in-network benefit for a 90-day transition period. Documentation of the authorization for the ongoing course of treatment from the previous health plan must be provided to us by the member or their provider and the services provided for the course of treatment must otherwise be a covered service under the Louisiana Blue health plan. This provision does not apply to medications covered under the plan's pharmacy benefit.