

Hematopoietic Cell Transplantation for Chronic Lymphocytic Leukemia and Small Lymphocytic Lymphoma

Policy # 00052

Original Effective Date: 01/28/2002

Current Effective Date: 10/01/2025

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Note: Hematopoietic Cell Transplantation for Non-Hodgkin Lymphomas is addressed separately in medical policy 00062.

When Services Are Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of available data, the Company may consider allogeneic hematopoietic cell transplantation (HCT) to treat chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma in individuals with markers of poor-risk disease to be **eligible for coverage.**** (see Policy Guidelines and Rationale). Use of a myeloablative or reduced-intensity pretransplant conditioning regimen should be individualized based on factors that include age, the presence of comorbidities, and disease burden.

When Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers allogeneic hematopoietic cell transplantation (HCT) to treat chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL) except as noted above to be **investigational.***

Based on review of available data, the Company considers autologous hematopoietic cell transplantation (HCT) to treat chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL) to be **investigational.***

Hematopoietic Cell Transplantation for Chronic Lymphocytic Leukemia and Small Lymphocytic Lymphoma

Policy # 00052

Original Effective Date: 01/28/2002

Current Effective Date: 10/01/2025

Policy Guidelines

Staging and Prognosis of Chronic Lymphocytic Leukemia or Small Lymphocytic Lymphoma

Two scoring systems are used to determine stage and prognosis of individuals with chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL). As outlined in Table 1, the Rai and Binet staging systems classify patients into 3 risk groups with different prognoses and are used to make therapeutic decisions.

Table 1. Rai and Binet Classification for Chronic Lymphocytic Leukemia or Small Lymphocytic Lymphoma

Rai Stage	Risk	Description	Median Survival, y	Binet Stage	Description	Median Survival, y
0	Low	Lymphocytosis	>10	A	< 3 lymphoid areas, normal hemoglobin and platelets	>10
I	Intermediate	Lymphocytosis + lymphadenopathy	7 to 9	B	≥3 lymphoid areas, normal hemoglobin and platelets	7
II	Intermediate	Lymphocytosis + splenomegaly/hepatomegaly ± lymphadenopathy	7 to 9			
III	High	Lymphocytosis + anemia ± lymphadenopathy, hepatomegaly, or splenomegaly	1.5 to 5	C	Any number of lymphoid areas, anemia, thrombocytopenia	5
IV	High	Lymphocytosis + thrombocytopenia ± anemia, splenomegaly, or lymphadenopathy	1.5 to 5			

Because prognoses of individuals vary within the different Rai and Binet classifications, other prognostic markers are used in conjunction with staging to determine clinical management.

The National Comprehensive Cancer Network guideline on CLL/SLL stated the following as unfavorable prognostic factors: DNA sequencing with mutated TP53 or ≤2% immunoglobulin heavy-chain variable (IGHV) mutation; interphase cytogenetics with del17p or deletion of 11q (del11q); or complex karyotype (≥3 unrelated chromosome abnormalities in more than 1 cell on karyotype).

Hematopoietic Cell Transplantation for Chronic Lymphocytic Leukemia and Small Lymphocytic Lymphoma

Policy # 00052

Original Effective Date: 01/28/2002

Current Effective Date: 10/01/2025

Reduced-Intensity Conditioning for Allogeneic Hematopoietic Cell Transplantation

Use of a myeloablative or reduced-intensity pretransplant conditioning regimen should be individualized based on factors that include age, the presence of comorbidities, and disease burden.

Some individuals for whom a conventional myeloablative allotransplant could be curative may be considered as candidates for reduced-intensity conditioning (RIC) allogeneic hematopoietic cell transplantation (allo-HCT). These include those individuals whose age (typically >60 years) or comorbidities (eg, liver or kidney dysfunction, generalized debilitation, prior intensive chemotherapy, low Karnofsky Performance Status score) preclude use of a standard myeloablative conditioning regimen. An individual who relapses following a conventional myeloablative allo-HCT could undergo a second myeloablative procedure if a suitable donor is available and his or her medical status would permit it. However, this type of individual would likely undergo RIC before a second allo-HCT if complete remission could be reinduced with chemotherapy.

The ideal allogeneic donors are human leukocyte antigen (HLA)- identical siblings, matched at the HLA-A, -B, and -DR loci on each arm of chromosome 6. Related donors mismatched at a single locus are also considered suitable donors. A matched, unrelated donor identified through the National Marrow Donor Registry is typically the next option considered. Recently, haploidentical donors-typically a parent or a child of the individual -with whom usually there is sharing of only 3 of the 6 major histocompatibility antigens, have been under investigation as a stem cell source. Most individuals will have such a donor; however, the risk of graft-versus-host disease and overall morbidity of the procedure may be severe, and experience with these donors is not as extensive as that with matched donors.

Background/Overview

Chronic Lymphocytic Leukemia and Small Lymphocytic Lymphoma

Chronic lymphocytic leukemia (CLL) and small lymphocytic lymphoma (SLL) are neoplasms of hematopoietic origin characterized by the accumulation of lymphocytes with a mature, generally well-differentiated morphology. In CLL, these cells accumulate in the blood, bone marrow, lymph nodes, and spleen; in SLL they are generally confined to lymph nodes. The Revised European-American/World Health Organization Classification of Lymphoid Neoplasms considers B-cell CLL and SLL a single disease entity.

Chronic lymphocytic leukemia and SLL share many common features and are often referred to as blood and tissue counterparts of each other, respectively. Both tend to present as asymptomatic enlargement of the lymph nodes, tend to be indolent, but can undergo transformation to a more aggressive form of the disease (eg, Richter transformation). The median age at diagnosis of CLL is approximately 72 years, but it may present in younger individuals, often as a poor-risk disease with significantly reduced life expectancy.

Hematopoietic Cell Transplantation for Chronic Lymphocytic Leukemia and Small Lymphocytic Lymphoma

Policy # 00052

Original Effective Date: 01/28/2002

Current Effective Date: 10/01/2025

Treatment regimens used for CLL are generally the same as those used for SLL, and treatment outcomes are comparable for both diseases. Both low- and intermediate-risk CLL and SLL demonstrate relatively good prognoses, with median survivals of 6 to 10 years; however, the median survival of high-risk CLL or SLL may only be 2 years. Although typically responsive to initial therapy, CLL and SLL are rarely cured by conventional therapy, and nearly all patients ultimately die of their disease. This natural disease history prompted an investigation of HCT as a possible curative regimen.

Hematopoietic Cell Transplantation

Hematopoietic cell transplantation (HCT) is a procedure in which hematopoietic stem cells are infused to restore bone marrow function in cancer patients who receive bone marrow-toxic doses of drugs with or without whole-body radiotherapy. Hematopoietic stem cells may be obtained from the transplant recipient (autologous HCT) or a donor (allogeneic HCT [allo-HCT]). These cells can be harvested from bone marrow, peripheral blood, or umbilical cord blood shortly after delivery of neonates. Although cord blood is an allogeneic source, the stem cells in it are antigenically "naive" and thus are associated with a lower incidence of rejection or graft-versus-host disease (GVHD).

Immunologic compatibility between infused hematopoietic stem cells and the recipient is not an issue in autologous HCT. However, immunologic compatibility between donor and patient is critical for achieving a good outcome of allo-HCT. Compatibility is established by typing of human leukocyte antigens (HLA) using cellular, serologic, or molecular techniques. HLA refers to the tissue type expressed at the HLA-A, -B, and -DR loci on each arm of chromosome 6. Depending on the disease being treated, an acceptable donor will match the patient at all or most of the HLA loci.

Conditioning for Hematopoietic Cell Transplantation

Conventional Conditioning for Hematopoietic Cell Transplantation

The conventional practice of allo-HCT involves administration of cytotoxic agents (eg, cyclophosphamide, busulfan) with or without total body irradiation at doses sufficient to destroy endogenous hematopoietic capability in the recipient. The beneficial treatment effect in this procedure is due to a combination of initial eradication of malignant cells and subsequent graft-versus-malignancy effect that develops after engraftment of allogeneic stem cells within the patient's bone marrow space. The slower graft-versus-malignancy effect is considered the potentially curative component, but it may be overwhelmed by extant disease without the use of pretransplant conditioning. However, intense conditioning regimens are limited to patients who are sufficiently fit medically to tolerate substantial adverse events that include pre-engraftment opportunistic infections secondary to loss of endogenous bone marrow function and organ damage and failure caused by the cytotoxic drugs. Furthermore, in any allo-HCT, immunosuppressant drugs are required to minimize graft rejection and GVHD, which also increases the susceptibility of the patient to opportunistic infections.

Hematopoietic Cell Transplantation for Chronic Lymphocytic Leukemia and Small Lymphocytic Lymphoma

Policy # 00052

Original Effective Date: 01/28/2002

Current Effective Date: 10/01/2025

The success of autologous HCT is predicated on the ability of cytotoxic chemotherapy with or without radiation to eradicate cancerous cells from the blood and bone marrow. This permits subsequent engraftment and repopulation of bone marrow space with presumably normal hematopoietic stem cells obtained from the patient before undergoing bone marrow ablation. As a consequence, autologous HCT is typically performed as consolidation therapy when the patient's disease is in complete remission. Patients who undergo autologous HCT are susceptible to chemotherapy-related toxicities and opportunistic infections before engraftment, but not GVHD.

Reduced-Intensity Conditioning for Allogeneic Hematopoietic Cell Transplantation

Reduced-intensity conditioning (RIC) refers to the pretransplant use of lower doses or less intense regimens of cytotoxic drugs or radiation than are used in conventional full-dose myeloablative conditioning treatments. The goal of RIC is to reduce disease burden but also to minimize as much as possible associated treatment-related morbidity and nonrelapse mortality in the period during which the beneficial graft-versus-malignancy effect of allogeneic transplantation develops. Although the definition of RIC remains arbitrary, with numerous versions employed, all seek to balance the competing effects of nonrelapse mortality and relapse due to residual disease. Reduced-intensity conditioning regimens can be viewed as a continuum in effects, from near totally myeloablative to minimally myeloablative with lymphoablation, with intensity tailored to specific diseases and patient condition. Patients who undergo RIC with allo-HCT initially demonstrate donor cell engraftment and bone marrow-mixed chimerism. Most will subsequently convert to full-donor chimerism, which may be supplemented with donor lymphocyte infusions to eradicate residual malignant cells. For this evidence review, the term reduced-intensity conditioning will refer to all conditioning regimens intended to be nonmyeloablative, as opposed to fully myeloablative (conventional) regimens.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

The U.S. Food and Drug Administration regulates human cells and tissues intended for implantation, transplantation, or infusion through the Center for Biologics Evaluation and Research, under the Code of Federal Regulation title 21, parts 1270 and 1271. Hematopoietic cells are included in these regulations.

Rationale/Source

This medical policy was developed through consideration of peer-reviewed medical literature generally recognized by the relevant medical community, U.S. Food and Drug Administration approval status, nationally accepted standards of medical practice and accepted standards of medical practice in this community, technology evaluation centers, reference to regulations, other plan medical policies, and accredited national guidelines.

Hematopoietic Cell Transplantation for Chronic Lymphocytic Leukemia and Small Lymphocytic Lymphoma

Policy # 00052

Original Effective Date: 01/28/2002

Current Effective Date: 10/01/2025

Risk stratification of patients with chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL) guides therapy decisions, which may include hematopoietic cell transplantation (HCT) for those with poor risk features.

Summary of Evidence

For individuals who have chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL) and markers of poor-risk disease who receive allogeneic hematopoietic cell transplantation (allo-HCT), the evidence includes single-arm prospective and registry-based studies. Relevant outcomes are overall survival, disease-specific survival, change in disease status, and treatment-related mortality and morbidity. Data have suggested that allo-HCT can provide long-term disease control and overall survival in patients with poor-risk CLL/SLL. High rates of treatment-related morbidity discourage this approach in lower-risk disease, particularly among older patients whose health status typically precludes the use of myeloablative conditioning. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have CLL/SLL who receive autologous HCT, the evidence includes randomized controlled trials (RCTs) and a systematic review. Relevant outcomes are overall survival, disease-specific survival, change in disease status, and treatment-related mortality and morbidity. Autologous HCT is feasible in younger patients but is not curative, particularly in those with poor-risk CLL. Studies of autologous HCT published to date have not shown improvement in overall survival in patients with CLL/SLL, and results must be considered in the context of improved outcomes with the use of newer chemoimmunotherapy agents. Furthermore, evidence from the European Intergroup RCT has suggested quality of life issues are important in selecting patients for autologous HCT and may dictate the management course for patients who are otherwise candidates for this approach. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Supplemental Information

Clinical Input From Physician Specialty Societies and Academic Medical Centers

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American Society for Blood and Marrow Transplantation

In 2015, the American Society for Blood and Marrow Transplantation published guidelines on indications for allo-HCT and autologous HCT for CLL. Recommendations described the current

Hematopoietic Cell Transplantation for Chronic Lymphocytic Leukemia and Small Lymphocytic Lymphoma

Policy # 00052

Original Effective Date: 01/28/2002

Current Effective Date: 10/01/2025

consensus on the use of HCT in and out of the clinical trial setting. Treatment recommendations are shown in Table 2.

Table 2. 2015 Recommendations for Allogeneic and Autologous Hematopoietic Cell Transplantation for Chronic Lymphocytic Leukemia

Adult Indications	Allogeneic HCT	Autologous HCT
High-risk, first or greater remission	C	N
T cell, prolymphocytic leukemia	R	R
B cell, prolymphocytic leukemia	R	R
Transformation to high-grade lymphoma	C	C

C: standard of care, clinical evidence available; HCT: hematopoietic cell transplantation; N: not generally recommended; R: standard of care, rare indication.

In 2016, the Society published clinical practice recommendations with additional detail on allo-HCT for CLL. Recommendations are shown in Table 3 .

Table 3. 2016 Recommendations for Allogeneic Hematopoietic Cell Transplantation for Chronic Lymphocytic Leukemia

Indications	Allogeneic HCT
High-risk CLL	Not recommended in the first-line consolidation setting
	Not recommended for patients who relapse after first-line therapy and demonstrate sensitive disease after second-line therapy (not BCR inhibitors)
	Recommended for patients who relapse after first-line therapy, have refractory disease after second-line therapy (not BCR inhibitors), and show an objective response to BCR inhibitors or to a clinical trial
	Recommended for patients who relapse after first-line therapy, have refractory disease after second-line therapy (including BCR inhibitors but not BCL-2 inhibitors), and show an objective response to BCL-2 inhibitors or to a clinical trial

Hematopoietic Cell Transplantation for Chronic Lymphocytic Leukemia and Small Lymphocytic Lymphoma

Policy # 00052

Original Effective Date: 01/28/2002

Current Effective Date: 10/01/2025

	Recommended when there is a lack of response or there is progression after BCL-2 inhibitors
Richter transformation	Recommended after achieving an objective response to anthracycline-based chemotherapy
Purine analogue relapsed and/or refractory disease	Not recommended

BCL-2: B-cell lymphoma 2; BCR: B-cell receptor; CLL: chronic lymphocytic leukemia; HCT: hematopoietic cell transplantation.

American Society for Transplantation and Cellular Therapy

In 2020, the American Society for Transplantation and Cellular Therapy (ASTCT) published guidelines on indications for HCT and immune effector cell therapy. Recommendations for CLL are shown in Table 4.

Table 4. 2020 Recommendations for Allogeneic Hematopoietic Cell Transplantation for Chronic Lymphocytic Leukemia

Adult Indications	Allogeneic HCT	Autologous HCT
High-risk, first or greater remission	S	N
T cell, prolymphocytic leukemia	S	R
B cell, prolymphocytic leukemia	R	R
Transformation to high-grade lymphoma	C	S

C: standard of care, clinical evidence available; HCT: hematopoietic cell transplantation; N: not generally recommended; R: standard of care, rare indication; S: standard of care

National Comprehensive Cancer Network Guidelines

Current National Comprehensive Cancer Network guidelines (v.1.2025) for CLL and small lymphocytic lymphoma (SLL) state the following regarding HCT:

- "Given the favorable outcome of patients with del(17p) or TP53 mutation treated with covalent BTKi as first-line therapy and the availability of venetoclax as an effective treatment option for relapsed or refractory CLL, allogeneic HCT is not considered as a reasonable treatment option for relapsed/refractory CLL after initial purine analogue-based therapy."
- "Allogeneic HCT can be considered for relapsed/refractory disease after prior therapy with BTKi- and venetoclax-based regimens in patients without significant comorbidities."

Hematopoietic Cell Transplantation for Chronic Lymphocytic Leukemia and Small Lymphocytic Lymphoma

Policy # 00052

Original Effective Date: 01/28/2002

Current Effective Date: 10/01/2025

- In patients with histologic transformation (Richter's) and progression, allogeneic HCT can be considered for certain patients with disease responding to initial chemotherapy. In addition, "autologous HCT may also be appropriate for patients with disease responding to initial therapy but who are not candidates for allogeneic HCT due to age, comorbidities, or lack of a suitable donor."

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

Ongoing and Unpublished Clinical Trials

A search of [ClinicalTrials.gov](https://clinicaltrials.gov) in November 2024 did not identify any ongoing or unpublished trials that would likely influence this review.

References

1. Giné E, Moreno C, Esteve J, et al. The role of stem-cell transplantation in chronic lymphocytic leukemia risk-adapted therapy. *Best Pract Res Clin Haematol*. Sep 2007; 20(3): 529-43. PMID 17707838
2. Gribben JG. Role of allogeneic hematopoietic stem-cell transplantation in chronic lymphocytic leukemia. *J Clin Oncol*. Oct 20 2008; 26(30): 4864-5. PMID 18794537
3. Kharfan-Dabaja MA, Anasetti C, Santos ES. Hematopoietic cell transplantation for chronic lymphocytic leukemia: an evolving concept. *Biol Blood Marrow Transplant*. Apr 2007; 13(4): 373-85. PMID 17382245
4. Gladstone DE, Fuchs E. Hematopoietic stem cell transplantation for chronic lymphocytic leukemia. *Curr Opin Oncol*. Mar 2012; 24(2): 176-81. PMID 22234253
5. Delgado J, Milligan DW, Dreger P. Allogeneic hematopoietic cell transplantation for chronic lymphocytic leukemia: ready for prime time?. *Blood*. Sep 24 2009; 114(13): 2581-8. PMID 19641189
6. Dreger P. Allogeneic transplantation for chronic lymphocytic leukemia. *Hematology Am Soc Hematol Educ Program*. 2009: 602-9. PMID 20008245
7. Brown JR, Kim HT, Li S, et al. Predictors of improved progression-free survival after nonmyeloablative allogeneic stem cell transplantation for advanced chronic lymphocytic leukemia. *Biol Blood Marrow Transplant*. Oct 2006; 12(10): 1056-64. PMID 17084369
8. Delgado J, Thomson K, Russell N, et al. Results of alemtuzumab-based reduced-intensity allogeneic transplantation for chronic lymphocytic leukemia: a British Society of Blood and Marrow Transplantation Study. *Blood*. Feb 15 2006; 107(4): 1724-30. PMID 16239425

Hematopoietic Cell Transplantation for Chronic Lymphocytic Leukemia and Small Lymphocytic Lymphoma

Policy # 00052

Original Effective Date: 01/28/2002

Current Effective Date: 10/01/2025

9. Dreger P, Brand R, Hansz J, et al. Treatment-related mortality and graft-versus-leukemia activity after allogeneic stem cell transplantation for chronic lymphocytic leukemia using intensity-reduced conditioning. *Leukemia*. May 2003; 17(5): 841-8. PMID 12750695
10. Khouri IF, Saliba RM, Admirand J, et al. Graft-versus-leukaemia effect after non-myeloablative haematopoietic transplantation can overcome the unfavourable expression of ZAP-70 in refractory chronic lymphocytic leukaemia. *Br J Haematol*. May 2007; 137(4): 355-63. PMID 17456058
11. Schetelig J, Thiede C, Bornhauser M, et al. Evidence of a graft-versus-leukemia effect in chronic lymphocytic leukemia after reduced-intensity conditioning and allogeneic stem-cell transplantation: the Cooperative German Transplant Study Group. *J Clin Oncol*. Jul 15 2003; 21(14): 2747-53. PMID 12860954
12. Sorrow ML, Storer BE, Sandmaier BM, et al. Five-year follow-up of patients with advanced chronic lymphocytic leukemia treated with allogeneic hematopoietic cell transplantation after nonmyeloablative conditioning. *J Clin Oncol*. Oct 20 2008; 26(30): 4912-20. PMID 18794548
13. Dreger P, Döhner H, Ritgen M, et al. Allogeneic stem cell transplantation provides durable disease control in poor-risk chronic lymphocytic leukemia: long-term clinical and MRD results of the German CLL Study Group CLL3X trial. *Blood*. Oct 07 2010; 116(14): 2438-47. PMID 20595516
14. Reljic T, Kumar A, Djulbegovic B, et al. High-dose therapy and autologous hematopoietic cell transplantation as front-line consolidation in chronic lymphocytic leukemia: a systematic review. *Bone Marrow Transplant*. Aug 2015; 50(8): 1144. PMID 26242579
15. Michallet M, Dreger P, Sutton L, et al. Autologous hematopoietic stem cell transplantation in chronic lymphocytic leukemia: results of European intergroup randomized trial comparing autografting versus observation. *Blood*. Feb 03 2011; 117(5): 1516-21. PMID 21106985
16. de Wreede LC, Watson M, van Os M, et al. Improved relapse-free survival after autologous stem cell transplantation does not translate into better quality of life in chronic lymphocytic leukemia: lessons from the randomized European Society for Blood and Marrow Transplantation-Intergroup study. *Am J Hematol*. Feb 2014; 89(2): 174-80. PMID 24123244
17. Majhail NS, Farnia SH, Carpenter PA, et al. Indications for Autologous and Allogeneic Hematopoietic Cell Transplantation: Guidelines from the American Society for Blood and Marrow Transplantation. *Biol Blood Marrow Transplant*. Nov 2015; 21(11): 1863-1869. PMID 26256941
18. Kharfan-Dabaja MA, Kumar A, Hamadani M, et al. Clinical Practice Recommendations for Use of Allogeneic Hematopoietic Cell Transplantation in Chronic Lymphocytic Leukemia on Behalf of the Guidelines Committee of the American Society for Blood and Marrow Transplantation. *Biol Blood Marrow Transplant*. Dec 2016; 22(12): 2117-2125. PMID 27660167
19. Kanate AS, Majhail NS, Savani BN, et al. Indications for Hematopoietic Cell Transplantation and Immune Effector Cell Therapy: Guidelines from the American Society for Transplantation and Cellular Therapy. *Biol Blood Marrow Transplant*. Jul 2020; 26(7): 1247-1256. PMID 32165328

Hematopoietic Cell Transplantation for Chronic Lymphocytic Leukemia and Small Lymphocytic Lymphoma

Policy # 00052

Original Effective Date: 01/28/2002

Current Effective Date: 10/01/2025

20. National Comprehensive Cancer Network (NCCN). NCCN clinical practice guidelines in oncology: chronic lymphocytic leukemia/small lymphocytic lymphoma. Version 1.2025. https://www.nccn.org/professionals/physician_gls/pdf/cll.pdf.

Policy History

Original Effective Date: 01/28/2002

Current Effective Date: 10/01/2025

12/06/2001	Medical Policy Committee review
01/28/2002	Managed Care Advisory Council approval
06/24/2002	Format revision. Coverage eligibility policy unchanged.
03/31/2004	Medical Director review
04/20/2004	Medical Policy Committee review. Format revision.
04/26/2004	Managed Care Advisory Council approval
06/03/2005	Medical Director review
06/21/2005	Medical Policy Committee review. Format revision. Rationale and Source added.
07/15/2005	Managed Care Advisory Council approval
07/07/2006	Format revision, including addition of FDA and or other governmental regulatory approval and rationale/source. Coverage eligibility unchanged.
08/01/2007	Medical Director review
08/15/2007	Medical Policy Committee approval. Rationale/Source updated. Coverage eligibility unchanged.
08/06/2009	Medical Policy Committee approval.
08/26/2009	Medical Policy Implementation Committee approval. Rationale/Source updated. Coverage eligibility unchanged. Title changed.
07/01/2010	Medical Policy Committee approval.
07/21/2010	Medical Policy Implementation Committee approval. Policy statement regarding allogeneic transplant in patients with markers of poor-risk disease changed; now may be considered medically necessary.
07/07/2011	Medical Policy Committee approval.
07/20/2011	Medical Policy Implementation Committee approval. No change to coverage.
06/28/2012	Medical Policy Committee review
07/27/2012	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
03/04/2013	Coding updated
08/01/2013	Medical Policy Committee review
08/21/2013	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
09/04/2014	Medical Policy Committee review

Hematopoietic Cell Transplantation for Chronic Lymphocytic Leukemia and Small Lymphocytic Lymphoma

Policy # 00052

Original Effective Date: 01/28/2002

Current Effective Date: 10/01/2025

09/17/2014	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
08/03/2015	Coding update: ICD10 Diagnosis code section added; ICD9 Procedure code section removed.
09/03/2015	Medical Policy Committee review
09/23/2015	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
09/08/2016	Medical Policy Committee review
09/21/2016	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
01/01/2017	Coding update: Removing ICD-9 Diagnosis Codes
09/07/2017	Medical Policy Committee review
09/20/2017	Medical Policy Implementation Committee approval. The word stem was removed from title and body of the policy.
09/06/2018	Medical Policy Committee review
09/19/2018	Medical Policy Implementation Committee approval. Added policy guidelines.
09/05/2019	Medical Policy Committee review
09/11/2019	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
09/03/2020	Medical Policy Committee review
09/09/2020	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
09/02/2021	Medical Policy Committee review
09/08/2021	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
09/01/2022	Medical Policy Committee review
09/14/2022	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
09/07/2023	Medical Policy Committee review
09/13/2023	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
09/05/2024	Medical Policy Committee review
09/11/2024	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
09/04/2025	Medical Policy Committee review
09/10/2025	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
05/01/2026	Coding update.

Next Scheduled Review Date: 09/2026

Hematopoietic Cell Transplantation for Chronic Lymphocytic Leukemia and Small Lymphocytic Lymphoma

Policy # 00052

Original Effective Date: 01/28/2002

Current Effective Date: 10/01/2025

Coding

The five character codes included in the Louisiana Blue Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)[‡], copyright 2024 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Louisiana Blue Medical Policy Coverage Guidelines is with Louisiana Blue and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Louisiana Blue Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Louisiana Blue Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	38204, 38205, 38206, 38207, 38208, 38209, 38210, 38211, 38212, 38213, 38214, 38215, 38230, 38232, 38240, 38241, 38243 Delete code effective 05/01/2026: 38242
HCPCS	S2140, S2142, S2150
ICD-10 Diagnosis	All related diagnoses

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or

Hematopoietic Cell Transplantation for Chronic Lymphocytic Leukemia and Small Lymphocytic Lymphoma

Policy # 00052

Original Effective Date: 01/28/2002

Current Effective Date: 10/01/2025

- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
1. Consultation with technology evaluation center(s);
 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
 3. Reference to federal regulations.

****Medically Necessary (or “Medical Necessity”)** - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

For these purposes, “nationally accepted standards of medical practice” means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: If the Patient’s health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

NOTICE: Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage.