

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Investigational or experimental services are not covered. This includes any drug, device, procedure, or service provided under the investigational arm of a clinical trial or clinical study. These services are excluded from coverage under benefits.

When Services May Be Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of available data, the Company may consider systemic hyperbaric oxygen therapy (HBO) to be **eligible for coverage**.**

Patient Selection Criteria

Coverage eligibility may be considered for systemic hyperbaric oxygen therapy (HBO) in the treatment of **ANY** of the following conditions:

- Non healing diabetic wounds of the lower extremities in patients who meet **ALL** of the following criteria:
 - Patient has type 1 or type 2 diabetes and has a lower-extremity wound due to diabetes; **AND**
 - Patient has a wound classified as Wagner grade 3 or higher:
 - Grade 3- lesion has penetrated deeper than grade 2, and there is abscess, osteomyelitis, pyarthrosis, plantar space abscess, or infection of the tendon and tendon sheaths (Grade 2 ulcer penetrates to tendon, bone, or joint)
 - Grade 4- wet or dry gangrene in the toes or forefoot
 - Grade 5- gangrene involves the whole foot or such a percentage that no local procedures are possible and amputation is indicated; **AND**
 - Patient has no measurable signs of healing after 30 days of an adequate course of standard wound therapy, which includes all of the following:
 - Assessment and correction of any vascular problems in the affected limb (as feasible)
 - Optimization of nutritional status and glucose control
 - Debridement to remove devitalized tissue (if present)
 - Maintenance of a clean, moist bed of granulation tissue
 - Appropriate off-loading

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

- Treatment of infection (if present); **AND**
 - Standard wound therapy will be continued; **AND**
 - Wounds will be evaluated at least every 30 days and HBO continued only if measurable signs of healing have been documented within 30-day period of treatment; **OR**
- Acute traumatic peripheral ischemia (e.g., crush injuries, suturing of severed limbs, reperfusion injury, compartment syndrome); **OR**
- Peripheral arterial insufficiency related to non-healing arterial insufficiency ulcers with no measurable signs of healing after 30 days of an adequate course of standard wound therapy; **OR**
- Decompression sickness; **OR**
- Gas embolism, acute; **OR**
- Gas gangrene (i.e., clostridial myonecrosis); **OR**
- Progressive necrotizing soft tissue infections (necrotizing fasciitis), while receiving broad spectrum antibiotics with either performed or planned surgical debridement; **OR**
- Compromised skin grafts or flaps (not for primary management of wounds) with any of the following:
 - Visible ischemic changes such as pallor, mottling, or frank necrosis of the overlying skin; **OR**
 - Documented hypoxia or decreased perfusion and transcutaneous oxygen tension TcPO₂ less than 40 mm Hg on room air; **OR**
- Cyanide poisoning, acute; **OR**
- Acute carbon monoxide poisoning; **OR**
- Soft-tissue radiation necrosis (e.g., radiation enteritis, cystitis, proctitis) and osteoradionecrosis; **OR**
- Pre- and post-treatment for patients undergoing dental surgery (non-implant-related) of an irradiated jaw; **OR**
- Profound anemia with exceptional blood loss: only when blood transfusion is impossible or must be delayed; **OR**
- Treatment-refractory osteomyelitis; **OR**
- Idiopathic sudden sensorineural hearing loss (ISSHL) in individuals with one of the following:
 - Sudden hearing loss of 30 decibels or more; **OR**
 - Hearing loss at least 61 decibels of pure tone thresholds when HBOT is combined with steroid therapy (systemic or intra-tympanic) and initiated within 1 month of onset of SSNHL; **OR**
- Central retinal artery occlusion (CRAO).

When Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

Based on review of available data, the Company considers topical hyperbaric oxygen therapy (HBOT) to be **investigational**.*

Based on review of available data, the Company considers limb specific hyperbaric oxygen therapy (HBOT) to be **investigational**.*

Based on review of available data, the Company considers systemic hyperbaric oxygen therapy (HBO) in all other situations, including but not limited to, the treatment of the following conditions to be **investigational**.*

- Bisphosphonate-related osteonecrosis of the jaw;
- Acute peripheral artery insufficiency (outside of other listed medically necessary indications involving arterial insufficiency) (see Policy Guidelines);
- Acute thermal burns;
- Acute surgical and traumatic wounds not meeting criteria specified in the eligible for coverage statement;
- Chronic wounds, other than those in patients with diabetes who meet the criteria specified in the eligible for coverage statement;
- Spinal cord injury;
- Traumatic brain injury;
- Inflammatory bowel disease (Crohn disease or ulcerative colitis);
- Brown recluse spider bites;
- Bone grafts;
- Carbon tetrachloride poisoning, acute;
- Cerebrovascular disease, acute (thrombotic or embolic) or chronic;
- Fracture healing;
- Hydrogen sulfide poisoning;
- Intra-abdominal and intracranial abscesses;
- Lepromatous leprosy;
- Meningitis;
- Pseudomembranous colitis (antimicrobial agent-induced colitis);
- Radiation myelitis;
- Sickle cell crisis and/or hematuria;
- Demyelinating diseases (e.g., multiple sclerosis, amyotrophic lateral sclerosis);
- Retinopathy, adjunct to scleral buckling procedures in patients with sickle cell peripheral retinopathy and retinal detachment;
- Pyoderma gangrenosum;
- Acute coronary syndromes and as an adjunct to coronary interventions, including but not limited to, percutaneous coronary interventions and cardiopulmonary bypass;
- Refractory mycoses: mucormycosis, actinomycosis, conidiobolus coronato;
- Cerebral edema, acute;
- Migraine;
- In vitro fertilization;

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

- Cerebral palsy;
- Tumor sensitization for cancer treatments, including but not limited to, radiotherapy or chemotherapy;
- Delayed-onset muscle soreness;
- Idiopathic femoral neck necrosis;
- Chronic arm lymphedema following radiotherapy for cancer;
- Early treatment (beginning at completion of radiotherapy) to reduce adverse events of radiotherapy;
- Autism spectrum disorder;
- Bell palsy;
- Acute ischemic stroke;
- Motor dysfunction associated with stroke;
- Herpes zoster;
- Vascular dementia;
- Fibromyalgia; and
- Mental illness (i.e., posttraumatic stress disorder, generalized anxiety disorder or depression).

Policy Guidelines

Evidence

There is limited comparative evidence for HBOT. The policy is based on the best available evidence and is largely informed by clinical input and guidelines.

Topical Hyperbaric Oxygen

HCPCS code A4575 is used to describe a disposable topical hyperbaric oxygen appliance that creates a “chamber” around the wound area which is pressurized with “hyperbaric oxygen.” Conventional oxygen tanks, typically gas, are used to supply the oxygen. An example of such a device is the AOTI Hyper-Box^{TM†}.

This policy addresses topical hyperbaric oxygen therapy (HBOT) but not topical oxygen wound care.

Topical HBOT may be performed in the office, clinic, or may be self-administered by the patient in the home. Typically, the therapy is offered for 90 minutes per day for 4 consecutive days. After a 3-day break, the cycle is repeated. The regimen may last for 8 to 10 weeks.

Systemic Hyperbaric Oxygen

The Wagner classification system categorizes wounds as follows: grade 0, no open lesion; grade 1, superficial ulcer without penetration to deeper layers; grade 2, ulcer penetrates to tendon, bone, or joint; grade 3, lesion has penetrated deeper than grade 2, and there is abscess, osteomyelitis, pyarthrosis, plantar space abscess, or infection of the tendon and tendon sheaths; grade 4, wet or dry

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

gangrene in the toes or forefoot; grade 5, gangrene involves the whole foot or such a percentage that no local procedures are possible and amputation (at least at the below the knee level) is indicated.

Progressive Necrotizing Soft Tissue Infections

Necrotizing soft tissue infection (NSTI) is a set of disorders characterized by a rapidly progressive infection with necrosis or gangrene. No definition of "progressive" was identified. However, definition of NSTI includes progression of infection despite antibiotic therapy. UHMC clinical input speaks to progressive NSTI in terms of NSTI while receiving broad spectrum antibiotics with either performed or planned therapeutic and diagnostic surgical debridement. The UHMC input also notes that frozen section soft-tissue biopsy is the gold standard of diagnosis, but is not feasible in practice. There are no unique clinical considerations based on the wound characteristics, site and/or depth of infection or time to treatment. By their very nature, NSTI are life and limb threatening.

Idiopathic Sudden Sensorineural Hearing Loss (ISSHL)

Idiopathic sudden sensorineural hearing loss (ISSHL) is an abrupt loss of hearing, typically unilaterally, without a definitive or identifiable cause upon investigation, as is the case for 90% of sudden sensorineural hearing loss patients. The degree of hearing loss is typically defined as a loss of 30 decibels or more across 3 contiguous frequencies on audiogram. The hearing loss initially occurring on one side can occur subsequently on the contralateral side in the future. The exact etiology of ISSHL has not been elucidated but of the major proposed mechanisms may be mitigated by HBOT. ISSHL is included in the FDA approved uses of HBOT.

Central Retinal Artery Occlusion (CRAO) and Other Retinal Conditions

CRAO is relatively rare yet devastating diagnosis with poor prognosis for spontaneous recovery. Factors which influence outcome include the length of time of occlusion, the anatomical site of the occlusion, and the presence of a patent cilioretinal artery. The diagnosis of CRAO is typically and reliably made with a fundoscopic exam. Advanced diagnostic studies can confirm CRAO but are not required for the diagnosis. Treatments for CRAO include ocular massage, anterior chamber paracentesis, fibrinolysis, and ocular pressure lowering agents. However, none of these demonstrate improved outcomes compared to control. The FDA has added CRAO to the list of cleared indications for HBOT.

CRAO is a rare complication associated with CaHA (calcium hydroxylapatite) cosmetic filler injection, likely due to embolism.

In addition to CRAO, there are related clinical syndromes for which HBOT could be considered. This includes individuals with branch retinal artery occlusion, particularly those with complete or near complete blindness in the contralateral eye. Also, Susac's Syndrome which is a rare disorder thought to be an autoimmune endotheliopathy causing vascular injury and deposition of thrombotic material in the lumen of small vessels. Treatments for this syndrome include steroids, anticoagulation, and IVIG; HBOT might improve visual acuity for these individuals.

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

Acute Peripheral Artery Insufficiency

For this policy review, the indication of acute peripheral artery insufficiency is too broad to include as a stand alone indication for HBOT.

Acute peripheral artery insufficiency is not included in the FDA list of approved conditions for HBOT. The Undersea and Hyperbaric Medical Society guidelines (15th edition) include peripheral artery insufficiency as an indication for HBOT related to diabetic foot ulcers and non-healing arterial insufficiency ulcers but does not have a stand-alone indication for acute peripheral artery insufficiency.

Acute arterial Insufficiencies (AAI) are interruptions, complete or partial, of perfusion that put the tissues distal to the interruption at risk for loss of function or dying. AAIs thereby span all arteries including a variety of conditions already included in this review (e.g., central retinal artery occlusion, ischemic stroke, compartment syndrome). Acute peripheral artery insufficiency (also called peripheral arterial insufficiency) would be a subtype of AAI. Peripheral artery insufficiency is also referred to as peripheral artery disease (PAD). PAD is defined by the American Heart Association as a narrowing of the peripheral arteries that carry blood away from the heart to other parts of the body, and is typically further defined to narrowed arteries reducing blood flow to the arms or legs. The AHA states the most common type of PAD is lower-extremity PAD. In 2024, the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines published a Guideline for the Management of Lower Extremity PAD. The Guideline suggests that HBOT may have a limited role, and states HBOT: “*may be considered as an adjunctive therapy to revascularization for wound healing in the context of CLTI (chronic limb threatening ischemia) and diabetic foot ulcers*”. No other mention of HBOT is made, including no mention of HBOT for acute limb ischemia.

Timing and Duration of HBOT Treatment

While broad indications are given above, the decision to treat with HBOT and timing of HBOT should be made on a case-by-case basis. For example, acute arterial ischemias have a spectrum of times that vary by tissue type: minutes for neurological tissues, hours for muscle, days for skin and bone, and even longer for relatively avascular connective tissues, cartilage, and adnexal structures. Even for indications with guideline-based time periods there are case studies showing improvement outside of such windows. For example, the Undersea and Hyperbaric Medical Society Committee recommends HBOT treatment for central retinal artery occlusion (CRAO) within 24 hours of onset, as studies demonstrate the outcome of HBOT is improved with early treatment. However, successful cases have been reported in which treatment began later, sometimes up to weeks later. Given the safety of HBOT, the lack of successful alternative medical treatments, the debilitating impact of vision loss, and the challenges faced in getting a patient to a hyperbaric facility, it is difficult to provide a specific time cutoff after which HBOT should not be tried for CRAO.

The policy supplemental information includes considerations for recommended treatment dose and number of treatment sessions for various conditions as found in the UHMS Hyperbaric Oxygen Therapy Guidelines (15th edition).

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

Background/Overview

Hyperbaric Oxygen Therapy

Hyperbaric oxygen therapy (HBOT) is a technique for delivering higher pressures of oxygen to tissue. Two methods of administration are available: topical and systemic.

Topical Hyperbaric Oxygen Therapy

Topical hyperbaric therapy is a technique of delivering 100% oxygen directly to an open, moist wound at a pressure slightly higher than atmospheric pressure. It is hypothesized that the high concentrations of oxygen diffuse directly into the wound to increase the local cellular oxygen tension, which in turn promotes wound healing. Devices consist of an appliance to enclose the wound area (frequently an extremity) and a source of oxygen; conventional oxygen tanks may be used. The appliances may be disposable and may be used without supervision in the home by well-trained patients. Topical hyperbaric therapy has been investigated as a treatment of skin ulcerations resulting from diabetes, venous stasis, postsurgical infection, gangrenous lesion, decubitus ulcers, amputations, skin graft, burns, or frostbite.

Systemic Hyperbaric Oxygen Therapy

In systemic or large hyperbaric oxygen chambers, the patient is entirely enclosed in a pressure chamber and breathes oxygen at a pressure greater than 1 atmosphere (the pressure of oxygen at sea level). Thus, this technique relies on systemic circulation to deliver highly oxygenated blood to the target site, typically a wound. Systemic HBOT can be used to treat systemic illness, such as air or gas embolism, carbon monoxide poisoning, or clostridial gas gangrene. Treatment may be carried out either in a monoplace chamber pressurized with pure oxygen or in a larger, multiplace chamber pressurized with compressed air, in which case the patient receives pure oxygen by mask, head tent, or endotracheal tube.

Adverse Events

HBOT is a generally safe therapy, with an estimated adverse side effect rate of 0.4%. Adverse events may occur either from pressure effects or the oxygen. The pressure effect (barotrauma) may affect any closed air-filled cavity such as ears, sinus, teeth, and lungs. Pain and/or swelling may occur at these sites as pressure increases during the procedure and decreases as the procedure is ending. Oxygen toxicity may affect the pulmonary, neurologic, or ophthalmologic systems. Pulmonary symptoms include a mild cough, substernal burning, and dyspnea. Neurologic effects include tunnel vision, tinnitus, nausea, and dizziness. Ophthalmologic effects include retinopathy in neonates, cataract formation, and transient myopic vision changes.

Note that this evidence review does not address topical oxygen therapy in the absence of pressurization.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

Since 1979, the U.S. Food and Drug Administration (FDA) has cleared multiple topical and systemic hyperbaric oxygen administration devices through the 510(k) pathway. In 2013, the FDA published

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

a statement warning that non-FDA approved uses of HBOT may endanger the health of patients.³ If patients mistakenly believe that HBOT devices have been proven safe for uses not cleared by the FDA, they may delay or forgo proven medical therapies.

As of July 2021, the FDA has cleared hyperbaric chambers for the following disorders:

- Air and gas bubbles in blood vessels
- Anemia (severe anemia when blood transfusions cannot be used)
- Burns (severe and large burns treated at a specialized burn center)
- Carbon monoxide poisoning
- Crush injury
- Decompression sickness (diving risk)
- Gas gangrene
- Hearing loss (complete hearing loss that occurs suddenly and without any known cause)
- Infection of the skin and bone (severe)
- Radiation injury
- Skin graft flap at risk of tissue death
- Vision loss (when sudden and painless in one eye due to blockage of blood flow)
- Wounds (non-healing, diabetic foot ulcers).

HBOT is being studied for other conditions, including COVID-19. However, at this time, the FDA has not cleared or authorized the use of any HBOT device to treat COVID-19 or any conditions beyond those listed above. The website, clinicaltrials.gov, has more information on HBOT clinical trials for COVID-19 and other conditions.

Risks of hyperbaric oxygen therapy

When HBOT chambers are used for indications cleared by the FDA, HBOT is generally safe, and serious complications are rare.

Because of the increased pressure and increased concentration of the oxygen during HBOT, potential risks include:

- Ear and sinus pain
- Middle ear injuries, including tympanic membrane rupture
- Temporary vision changes
- Lung collapse (rare)

High concentrations of oxygen also pose the risk of fire, which is one reason why the FDA recommends treatment at an accredited facility. Explosions and fires have occurred in HBOT chambers that have not been reviewed by the FDA and are located at unaccredited facilities.

Other hyperbaric devices

The FDA has also cleared a large, zippered bag that is intended to treat altitude sickness only.

These zippered chambers for treating altitude sickness provide pressure but do not attach to oxygen tanks. The FDA has not cleared these bags for use with oxygen tanks or oxygen concentrators.

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

However, the FDA is aware of instances in which people used these bags to create homemade HBOT devices, which can pose the risk of fire and suffocation.

Rationale/Source

This medical policy was developed through consideration of peer-reviewed medical literature generally recognized by the relevant medical community, U.S. Food and Drug Administration approval status, nationally accepted standards of medical practice and accepted standards of medical practice in this community, technology evaluation centers, reference to regulations, other plan medical policies, and accredited national guidelines.

Hyperbaric oxygen therapy (HBOT) involves breathing 100% oxygen at pressures between 1.5 and 3.0 atmospheres. It is generally applied systemically with the patient inside a hyperbaric chamber. HBOT can also be applied topically; ie, the body part to be treated is isolated (eg, in an inflatable bag and exposed to pure oxygen). HBOT has been investigated for various conditions that have potential to respond to increased oxygen delivery to tissue.

Summary of Evidence

For individuals with wounds, burns or infections who receive topical hyperbaric oxygen therapy (HBOT), the evidence includes a systematic review, case series, and a randomized controlled trial (RCT). Relevant outcomes are overall survival (OS), symptoms, change in disease status, and functional outcomes. The systematic review identified 3 RCTs including patients with sacral pressure ulcers, ischial pressure ulcers, and refractory venous ulcers. All trials reported that healing improved significantly after HBOT than after standard of care. Pooling of results was not possible due to heterogeneity in patient populations and treatment regimens. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with chronic diabetic ulcers who receive systemic HBOT, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms and change in disease status. Meta-analyses of RCTs found significantly higher diabetic ulcer healing rates with HBOT than with control conditions. Two of the 3 meta-analyses found that HBOT was associated with a significantly lower rate of major amputation. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with carbon monoxide poisoning who receive systemic HBOT, the evidence includes RCTs and a systematic review. Relevant outcomes are OS and symptoms. A meta-analysis in a Cochrane review of low-quality RCT data did not find HBOT to be associated with a significantly lower risk of neurologic deficits after carbon monoxide poisoning. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with radionecrosis, osteoradionecrosis, or treatment of irradiated jaw who receive systemic HBOT, the evidence includes RCTs and a systematic review. Relevant outcomes are symptoms and change in disease status. A meta-analysis in a Cochrane review of RCTs found evidence that HBOT improved radionecrosis and osteoradionecrosis outcomes and resulted in better

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

outcomes before tooth extraction in an irradiated jaw. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with chronic refractory osteomyelitis who receive systemic HBOT, the evidence includes case series. Relevant outcomes are symptoms and change in disease status. The case series reported high rates of successful outcomes (no drainage, pain, tenderness, or cellulitis) in patients with chronic refractory osteomyelitis treated with HBOT. However, controlled studies are needed to determine conclusively the impact of HBOT on health outcomes compared with other interventions. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with acute thermal burns who receive systemic HBOT, the evidence includes a systematic review of 2 RCTs. Relevant outcomes are OS, symptoms, and change in disease status. Both RCTs were judged to have poor methodologic quality. Evidence from well-conducted controlled trials is needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with acute surgical and traumatic wounds who receive systemic HBOT, the evidence includes RCTs, controlled nonrandomized studies, and systematic reviews. Relevant outcomes are OS, symptoms, change in disease status, and functional outcomes. There was considerable heterogeneity across the 4 RCTs identified (eg, patient population, comparison group, treatment regimen, outcomes). This heterogeneity prevented pooling of trial findings and limits the ability to definitively conclude the impact of HBOT on health outcomes for patients with acute surgical and traumatic wounds. Additional evidence from high-quality RCTs is needed. A systematic review of controlled Chinese studies suggests HBOT may increase the survival rate of compromised skin grafts and flaps when initiated within 72 hours; however, risk of bias in the original Chinese publications cannot be evaluated. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with bisphosphonate-related osteonecrosis of the jaw who receive systemic HBOT, the evidence includes an RCT. Relevant outcomes are symptoms and change in disease status. The RCT was unblinded and reported initial benefits at 3-month follow-up; however, there were no significant benefits of HBOT for most health outcomes compared with standard care in the long-term (6 months to 2 years). The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with necrotizing soft tissue infections who receive systemic HBOT, the evidence includes systematic reviews. Relevant outcomes are OS, symptoms, and change in disease status. A Cochrane review did not identify any RCTs. Another systematic review of retrospective cohort studies with methodological limitations did not find consistent benefit of adjunctive HBOT use. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

For individuals with acute coronary syndrome who receive systemic HBOT, the evidence includes RCTs and a systematic review. Relevant outcomes are OS, symptoms, change in disease status, and functional outcomes. A Cochrane review identified 6 RCTs. There were 2 pooled analyses, 1 found significantly lower rates of death with HBOT and the other reported inconsistent results in left ventricular function. Additional RCT data are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with acute ischemic stroke with or without motor dysfunction associated with stroke who receive systemic HBOT, the evidence includes RCTs and a systematic review. Relevant outcomes are OS, symptoms, change in disease status, and functional outcomes. Cochrane reviewers could only pool data for a single outcome for acute ischemic stroke (mortality at 3 to 6 months), and for that outcome, there was no significant difference between active and sham HBOT treatments. An RCT found that HBOT combined with DAPT may enhance neurological recovery, and support daily living activities in elderly acute cerebral infarction patients; larger, multicenter studies with standardized HBOT protocols and extended follow-up are needed to confirm these findings. For motor dysfunction, an RCT, which used a crossover design, found better outcomes with HBOT at 2 months than with delayed treatment. However, the trial had a number of methodologic limitations (eg, lack of patient blinding, heterogeneous population, high dropout rate) that make it difficult to evaluate the efficacy of HBOT. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with motor dysfunction associated with stroke who receive systemic HBOT, the evidence includes an RCT. Relevant outcomes are symptoms and functional outcomes. The RCT, which used a crossover design, found better outcomes with HBOT at 2 months than with delayed treatment. However, the trial had a number of methodologic limitations (eg, lack of patient blinding, heterogeneous population, high dropout rate) that make it difficult to evaluate the efficacy of HBOT. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with Bell palsy who receive systemic HBOT, the evidence includes a systematic review. Relevant outcomes are symptoms, change in disease status, and functional outcomes. A Cochrane review did not identify any RCTs meeting selection criteria; the single RCT found did not have a blinded outcome assessment. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with traumatic brain injury who receive systemic HBOT, the evidence includes RCTs and systematic reviews. Relevant outcomes are OS, symptoms, change in disease status, and functional outcomes. RCTs were heterogeneous regarding intervention protocols, patient populations, and outcomes reported. Systematic reviews conducted pooled analyses only on a minority of the published RCTs, and these findings were inconsistent. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with inflammatory bowel disease who receive systemic HBOT, the evidence includes RCTs, observational studies, and a systematic reviews. Relevant outcomes are symptoms,

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

change in disease status and functional outcomes. RCTs have reported mixed findings in patients with ulcerative colitis, with one study terminated early due to futility. A systematic review including the RCT and observational studies found a high rate of bias in the literature due to attrition and reporting bias. Another systematic review found that HBOT appears to be a safe and effective adjunctive therapy for UC, with potential benefits for CD requiring further investigation. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with idiopathic sudden sensorineural hearing loss who receive systemic HBOT, the evidence includes systematic reviews. Relevant outcomes are symptoms, change in disease status, and functional outcomes. A Cochrane review of RCTs had mixed findings from studies that included individuals with tinnitus. Some outcomes (ie, improvement in hearing of all frequencies, >25% return of hearing) were better with HBOT than with a control intervention, but more than 50% return of hearing did not differ significantly between groups. There was important variability in the patients enrolled in the studies. A subsequent systematic review had similarly limited conclusions due to the inclusion of non-randomized studies. A third review found a higher proportion of patients with hearing recovery with HBOT compared to medical treatment alone, but the analysis was limited to 2 RCTs with methodological limitations. One RCT published subsequent to the systematic reviews found a positive effect of HBOT plus steroid combination therapy on measures of auditory function compared to either HBOT or steroids alone, but other outcomes were not reported and the study had numerous relevance, design, and conduct limitations. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with delayed-onset muscle soreness who receive systemic HBOT, the evidence includes RCTs and a systematic review. Relevant outcomes are symptoms and functional outcomes. A Cochrane review of RCTs found worse short-term pain outcomes with HBOT than with control and no difference in longer-term pain or other outcomes (eg, swelling). The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with autism spectrum disorder who receive systemic HBOT, the evidence includes an RCT and systematic reviews. Relevant outcomes are symptoms and functional outcomes. A systematic review reported that HBOT appears promising for reducing core ASD symptoms and enhancing functional outcomes, but further high-quality, multicenter trials are needed to confirm efficacy and establish optimal treatment parameters. A Cochrane review identified a single RCT on HBOT for autism spectrum disorder and this trial did not find significantly better parental-assessed or clinician-assessed outcomes with HBOT compared with sham. A subsequent controlled trial reached the same conclusion. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with cerebral palsy who receive systemic HBOT, the evidence includes 2 RCTs and an observational study. Relevant outcomes are symptoms and functional outcomes. One RCT was stopped early due to futility, and the other did not find significantly better outcomes with HBOT than with a sham intervention. The observational study focused on sleep disorders in children with

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

cerebral palsy and reported improvements with the HBOT treatment. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with vascular dementia who receive systemic HBOT, the evidence includes an RCT and a systematic review. Relevant outcomes are symptoms and functional outcomes. The Cochrane review identified only a single RCT with methodologic limitations. Well-conducted controlled trials are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with radiotherapy adverse events who receive systemic HBOT, the evidence includes RCTs, nonrandomized comparator trials, case series, and systematic reviews. Relevant outcomes are symptoms and functional outcomes. Two systematic reviews included few RCTs and provide limited evidence on the effect of HBOT. Two RCTs had inconsistent findings. One reported no short-term benefit with HBOT, but some benefits 12 months after radiotherapy; the other did not find a significant benefit of HBOT at 12-month follow-up. Another RCT assessed HBOT for radiation-induced cystitis and found significant benefit by some measures but not others. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with idiopathic femoral neck necrosis who receive systemic HBOT, the evidence includes an RCT. Relevant outcomes are symptoms, change in disease status, and functional outcomes. The RCT, which had a small sample, only reported short-term (ie, 6-week) outcomes. Larger well-conducted RCTs reporting longer-term outcomes are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with a migraine who receive systemic HBOT, the evidence includes RCTs and a systematic review. Relevant outcomes are symptoms, change in disease status, and functional outcomes. The Cochrane review conducted a pooled analysis including 3 of the 11 trials. Meta-analysis of these 3 RCTs found significantly greater relief of migraine symptoms with HBOT than with a comparator intervention within 45 minutes of treatment. Longer-term data are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with herpes zoster who receive systemic HBOT, the evidence includes a RCT. Relevant outcomes are symptoms and change in disease status. The RCT was unblinded and only reported short-term (ie, 6-week) outcomes. Additional well-conducted RCTs with longer follow-up are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with fibromyalgia who receive systemic HBOT, the evidence includes RCTs. Relevant outcomes are symptoms, change in disease status, and functional outcomes. Only 2 RCTs were identified, and both reported positive effects of HBOT on tender points and pain. However, the trials had relatively small samples and methodologic limitations (eg, quasi-randomization, no or uncertain sham control for a condition with subjective outcomes susceptible to a placebo effect).

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

Moreover, the HBOT protocols varied. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with multiple sclerosis who receive systemic HBOT, the evidence includes RCTs and a systematic review. Relevant outcomes are symptoms and functional outcomes. A Cochrane review of RCTs did not find a significant difference in Expanded Disability Status Scale scores when patients with multiple sclerosis were treated with HBOT versus a comparator intervention. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with cancer and are undergoing chemotherapy who receive systemic HBOT, the evidence includes an RCT and a systematic review. Relevant outcomes are OS and change in disease status. While the systematic review reported improvements in tumor control in patients with head and neck cancer who received HBOT, the adverse events accompanying the treatment (eg, radiation tissue injury, seizures) were significant. The single RCT did not find a significant difference in survival for cancer patients who received HBOT before chemotherapy compared with usual care. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Supplemental Information

Clinical Input From Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

2024 Input

Clinical input was sought to help determine whether the use of systemic HBOT in individuals with necrotizing soft tissue infections, idiopathic sudden sensorineural hearing loss, central retinal artery occlusion, or acute peripheral artery insufficiency would provide a clinically meaningful improvement in net health outcome and whether the use is consistent with generally accepted medical practice. In response to requests, clinical input was received from 2 respondents, including 2 specialty society-level responses.

For individuals with necrotizing soft tissue infections, idiopathic sudden sensorineural hearing loss, central retinal artery occlusion, or acute peripheral artery insufficiency who receive HBOT, clinical input supports this use provides a clinically meaningful improvement in net health outcomes and indicates this use is consistent with generally accepted medical practice.

2023 Input

Clinical input was sought to help determine whether the use of systemic hyperbaric oxygen therapy (HBOT) in individuals with acute surgical or traumatic wounds and compromised skin grafts or flaps would provide a clinically meaningful improvement in net health outcome and whether the use is

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

consistent with generally accepted medical practice. In response to requests, clinical input was received from 2 respondents, including 2 specialty society-level responses.

For individuals with acute surgical or traumatic wounds and compromised skin grafts or flaps who receive systemic HBOT, clinical input supports this use provides a clinically meaningful improvement in net health outcomes and indicates this use is consistent with generally accepted medical practice.

2010 Input

In response to requests, input was received from 6 physician specialty societies and 5 academic medical centers while this policy was under review in 2010. Clinical input varied by condition. There was consensus that topical hyperbaric oxygen therapy (HBOT) and systemic HBOT for autism spectrum disorder and headache/migraine are investigational. There was also wide support for adding acute carbon monoxide poisoning, compromised skin grafts or flaps, chronic refractory osteomyelitis, and necrotizing soft tissue infections to the list of medically necessary indications for HBOT. Several reviewers acknowledged that there is a paucity of clinical trials on HBOT for compromised skin grafts/flaps, necrotizing soft tissue infections, and chronic refractory osteomyelitis. These reviewers commented on the support from basic science, animal studies, and retrospective case series, as well as lack of effective alternative treatments for these conditions. Based on the available evidence and clinical input, acute carbon monoxide poisoning and chronic refractory osteomyelitis were changed in 2010 to medically necessary indications for HBOT. However, despite the clinical input and given the limited published evidence, compromised skin grafts and flaps and necrotizing soft tissue infections are still considered investigational.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American Academy of Otolaryngology - Head and Neck Surgery

In 2019, the American Academy of Otolaryngology-Head and Neck Surgery updated clinical guidelines on the treatment of sudden sensorineural hearing loss (SSNHL). They give the following options regarding HBOT:

"Clinicians may offer, or refer to a physician who can offer, hyperbaric oxygen therapy (HBOT) combined with steroid therapy within 2 weeks of onset of SSNHL."

"Clinicians may offer, or refer to a physician who can offer, hyperbaric oxygen therapy (HBOT) combined with steroid therapy as salvage within 1 months of onset of SSNHL."

The guideline provided a comprehensive list of evidence gaps and future research needs on the use of HBOT for SSNHL. These included, among others, the need for a standardized, evidence-based

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

definition of SSNHL, the assessment of the prevalence of SSNHL, and the need for the development of standardized HBOT treatment protocols and standardized outcome assessments.

American College of Cardiology/American Heart Association

In 2024, the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines published a Guideline for the Management of Lower Extremity PAD. The Guideline was developed in collaboration with and endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation, American Podiatric Medical Association, Association of Black Cardiologists, Society for Cardiovascular Angiography and Interventions, Society for Vascular Medicine, Society for Vascular Nursing, Society for Vascular Surgery, Society of Interventional Radiology, and Vascular & Endovascular Surgery Society. The Guideline included the following statements relevant to this evidence review:

"Beyond wound care, hyperbaric oxygen therapy has been studied in the context of wound healing for CLTI as an adjunctive therapy to revascularization and may have a limited role in this population."

"Hyperbaric oxygen therapy may be considered as an adjunctive therapy to revascularization for wound healing in the context of CLTI (chronic limb threatening ischemia) and diabetic foot ulcers."

Autism spectrum disorder

American Academy of Pediatrics

In 2020, the American Academy of Pediatrics published a clinical report on the management of children with autism spectrum disorder, which was reaffirmed in 2025. Regarding hyperbaric oxygen therapy (HBOT), the report states "Clinical trials do not support the use of antifungal agents, immunotherapy, or hyperbaric oxygen treatment, and concern for safety, in addition to lack of supporting data, cautions against chelation therapy for children with ASD [autism spectrum disorder]".

Inflammatory Bowel Disease

American Gastroenterological Association

In 2024, the American Gastroenterological Association (AGA) released a clinical guideline update on the management of moderate-to-severe ulcerative colitis in adults; the guidelines did not mention use of HBOT.

In 2025, the AGA released a clinical guideline update on the management on the management of moderate-to-severe Crohn's disease; the guidelines did not mention use of HBOT.

American College of Gastroenterology

In 2025, the American College of Gastroenterology (ACG) released a clinical guideline update on the management of ulcerative colitis in adults; the guidelines did not mention use of HBOT.

In 2025, the ACG released a clinical guideline update on the management of Crohn's disease in adults; the guidelines did not mention use of HBOT.

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

Stroke

American College of Cardiology/American Stroke Association

In 2019 the American Heart Association and American Stroke Association updated the guidelines for early management of acute ischemic stroke. The guidelines were endorsed by the Society for Academic Emergency Medicine, the Neurocritical Care Society, the American Association of Neurological Surgeons, and the Congress of Neurological Surgeons. The Guideline included the following statements relevant to this evidence review:

"The limited data available on the utility of HBO therapy for acute ischemic stroke (not related to cerebral air embolism) show no benefit. HBO therapy is associated with claustrophobia and middle ear barotrauma, as well as an increased risk of seizures. Given the confines of HBO chambers, the ability to closely/adequately monitor patients may also be compromised. HBO thus should be offered only in the context of a clinical trial or to individuals with cerebral air embolism."

The 2021 American Heart Association and American Stroke Association scientific statement on the primary care of adult patients after stroke did not mention use of HBOT.

Department of Veterans Affairs and Department of Defense

In 2021, the U.S. Department of Veterans Affairs (VA) and U.S. Department of Defense (DoD) released a clinical practice guideline on the management of stroke rehabilitation; the guidelines did not mention use of HBOT.

Traumatic Brain Injury

American College of Surgeons

In 2024, the American College of Surgeons released best practice guidelines on the management of traumatic brain injury (TBI). Utilization of HBOT is not mentioned.

Brain Trauma Foundation

In 2016, the Brain Trauma Foundation published guidelines for the management of severe traumatic brain injury. Hyperbaric oxygen therapy was listed as an excluded intervention for the guideline and thus not included within its evidence synthesis.

Department of Veterans Affairs and Department of Defense

In 2021, the U.S. VA/DoD released a clinical practice guideline on the management of post-acute mild TBI. Regarding HBOT, the guideline states "We recommend against the use of hyperbaric oxygen therapy for the treatment of patients with symptoms attributed to mild traumatic brain injury."

Diabetic Foot Ulcers

In 2016, the Society of Vascular Surgery in collaboration with the American Podiatric Medical Association and the Society for Vascular Medicine published guidelines on the management of the diabetic foot. According to the guidelines, for diabetic foot ulcers that fail to demonstrate improvement (>50% wound area reduction) after a minimum of 4 weeks of standard wound therapy, adjunctive therapy such as HBOT is recommended (grade 1B). Also, for diabetic foot ulcers with

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

adequate perfusion that fail to respond to 4 to 6 weeks of conservative management, HBOT is suggested (grade 2B).

In 2015, the Undersea and Hyperbaric Medical Society (UHMS) published guidelines on the use of HBOT for treating diabetic foot ulcers. Recommendations in the current version include:

- Suggest against using HBOT in patients with "Wagner Grade 2 or lower diabetic foot ulcers..."
- Suggest adding HBOT in patients with "Wagner Grade 3 or higher diabetic foot ulcers that have not shown significant improvement after 30 days of [standard of care] therapy..."
- Suggest "adding acute post-operative hyperbaric oxygen therapy to the standard of care" in patients with "Wagner Grade 3 or higher diabetic foot ulcers" who have just had foot surgery related to their diabetic ulcers.

Additional Conditions

The 2023 UHMS Hyperbaric Oxygen Therapy Indications (15th edition) included the following indications as recommended:

1. Air or Gas Embolism
2. Arterial insufficiencies: Central Retinal Artery Occlusion; Hyperbaric oxygen Therapy for Selected Problem Wounds
3. Carbon Monoxide Poisoning and carbon monoxide complicated by cyanide poisoning
4. Clostridial Myonecrosis (Gas Gangrene)
5. Acute Traumatic Ischemias
6. Decompression Sickness
7. Severe anemia
8. Intracranial abscess
9. Necrotizing soft tissue infections
10. Refractory osteomyelitis
11. Delayed radiation injury (soft tissue and bony necrosis)
12. Compromised grafts and flaps
13. Acute thermal burn injury
14. Sudden Sensorineural hearing loss
15. Avascular Necrosis (Aseptic Osteonecrosis).

Recommended treatment dose and number of treatment sessions per the UHMS Hyperbaric Oxygen Therapy Committee (15th edition, 2023) include:

- **Acute traumatic ischemia** – there are 3 stages of wound healing. Treatment recommended varies based on stage, and ranges from 2-3 times per day for 2-3 days for acute inflammatory stage, 14 days for repair stage, and up to a month for remodeling.
- **Carbon monoxide poisoning** – Use up to 3 ATA for 1 to 3 sessions or to clinical plateau.
- **Central Retinal Artery Occlusion (CRAO)**– Recommend 2 to 2.8 ATA or U.S. Navy Table 6 or equivalent. Treat twice daily to clinical plateau, which typically occurs in less than a week, plus 3 days. Hyperbaric treatments 2-3 times daily may be necessary until the angiogram normalizes or the patient has no further improvement for 3 treatments.

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

- **Clostridial myositis, Clostridial myonecrosis (Gas gangrene)**- Recommend 3 ATA pressure for 90 minutes, 2-3 times in the first 24 hours, and then 2 times daily for the next 2-5 days. Review is indicated after 10 treatments.
- **Chronic refractory osteomyelitis** – Typically, once daily, 5-7 days per week for 90-120 minutes using 2-3 ATA and continued for 4-6 weeks. 20-40 sessions are typically needed, although might be situations where up to 60 sessions are needed. Patients with refractory stage 3 or 4 osteomyelitis are most likely to benefit from adjunctive hyperbaric oxygen therapy, especially when complicated by adverse local or systemic factors.
- **Compartment syndrome** – Use 2 to 2.4 ATA, usually twice a day for 2 days but sometimes might need 3 times a day. After fasciotomy, twice a day for 7-14 days.
- **Compromised skin grafts and flaps** – Use 2 to 2.5 ATA twice daily for up to 20 sessions.
- **Crush injury** – similar to acute traumatic ischemia above. The UHMS notes that HBOT should be started as close as possible to the time of injury; 3 or more treatments during the first 24 to 72 hours are recommended; 1-2 times per day for 14 days if in the repair phase; daily use for 3-6 weeks during remodeling.
- **Cyanide poisoning** – Patients with cyanide poisoning frequently present with simultaneous carbon monoxide poisoning. Treatment protocol recommended is the same as for carbon monoxide poisoning.
- **Decompression sickness** – Use U.S. Navy Treatment Table 6 or equivalent, typically up to 2.8 ATA, for 1 session up to a clinical plateau. Typically, no more than 1 to 2 treatment sessions are needed.
- **Diabetic lower extremity wounds, selected individuals and healing of other problem wounds** – Use 2 to 2.5 ATA daily, should see effects by 2-3 weeks; course of outpatient therapy is typically 30 sessions but might require up to 40 sessions. For HBOT to continue, reevaluation at 30-day intervals must show continued progress in healing.
- **Necrotizing soft-tissue infections** – Use 2 to 2.5 ATA twice daily until stabilization occurs, often occurs within 7-10 treatments. If differential diagnosis includes the possibility of Clostridial myositis and/or myonecrosis and/or remains unclear, 2.8-3 ATA pressures are warranted with 3 treatments in the first 24-48 hours. Avoidance of premature cessation of HBOT is advised, and once extension of necrosis has been halted then once daily treatments over an extended period until the infection is well controlled is recommended. This might require 30 treatments. Review after 30 treatments.
- **Radiation Necrosis** – Most treatments range from 2-2.5 ATA for 40-60 treatments, and review should occur after 60 treatments.
 1. **Mandibular osteoradionecrosis, laryngeal necrosis, other soft tissue head and neck, chest wall necrosis, radiation cystitis, radiation proctitis, miscellaneous abdominal pelvic injuries, cutaneous necrosis** – 2 to 2.4 ATA daily for 90 minutes.
 2. **Neoadjuvant hyperbaric oxygen therapy before dental extractions** – 2 to 2.4 ATA, typically 20 treatments before extraction and 10 treatments after.
- **Sudden sensorineural hearing loss** – Recommend 2 to 2.5 ATA for 10 to 20 sessions.
- **Severe Anemia** – Use 2 to 3 ATA for 3 or 4 times a day until there is replacement of red blood cells by regeneration or transfusion.

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

In 2003 (updated in 2017), the Centers for Medicare & Medicaid added Medicare coverage of HBOT for diabetic wounds of the lower extremities meeting certain criteria. As of the current coverage statement, Medicare coverage is provided for HBOT administered in a chamber for the following conditions:

1. "Acute carbon monoxide intoxication,
2. Decompression illness,
3. Gas embolism,
4. Gas gangrene,
5. Acute traumatic peripheral ischemia. HBO therapy is a valuable adjunctive treatment to be used in combination with accepted standard therapeutic measures when loss of function, limb, or life is threatened.
6. Crush injuries and suturing of severed limbs. As in the previous conditions, HBO therapy would be an adjunctive treatment when loss of function, limb, or life is threatened.
7. Progressive necrotizing infections (necrotizing fasciitis),
8. Acute peripheral arterial insufficiency,
9. Preparation and preservation of compromised skin grafts (not for primary management of wounds),
10. Chronic refractory osteomyelitis, unresponsive to conventional medical and surgical management,
11. Osteoradionecrosis as an adjunct to conventional treatment,
12. Soft tissue radionecrosis as an adjunct to conventional treatment,
13. Cyanide poisoning,
14. Actinomycosis, only as an adjunct to conventional therapy when the disease process is refractory to antibiotics and surgical treatment,
15. Diabetic wounds of the lower extremities in patients who meet the following 3 criteria:
 - a. You have type I or type II diabetes and has a lower extremity wound that is due to diabetes;
 - b. You have a wound classified as Wagner grade III or higher; and
 - c. You have failed an adequate course of standard wound therapy."

The use of HBO therapy is covered as adjunctive therapy only after there are no measurable signs of healing for at least 30-days of treatment with standard wound therapy and must be used in addition to standard wound care. Standard wound care in patients with diabetic wounds includes: assessment of a patient's vascular status and correction of any vascular problems in the affected limb if possible, optimization of nutritional status, optimization of glucose control, debridement by any means to remove devitalized tissue, maintenance of a clean, moist bed of granulation tissue with appropriate moist dressings, appropriate off-loading, and necessary treatment to resolve any infection that might be present. Failure to respond to standard wound care occurs when there are no measurable signs of healing for at least 30 consecutive days. Wounds must be evaluated at least every 30 days during the

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

administration of HBO therapy. Continued treatment with HBO therapy is not covered if measurable signs of healing have not been demonstrated within any 30-day period of treatment.”

Systemic HBOT for other indications is not covered, nor is topical HBOT for any indication.

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 1.

Table 1. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT02407028	Hyperbaric Oxygen Brain Injury Treatment (HOBIT) Trial	200	Jun 2029
NCT06581003	Hyperbaric Oxygen Treatment for Veterans with Traumatic Brain Injury	420	Aug 2029
NCT07214415	Evaluating the Effects of Hyperbaric Oxygen Therapy on Cognitive Function, Mental Health, and Quality of Life in Veterans	54	Sept 2026
NCT06109961	Hyperbaric Oxygen Therapy for Perianal Fistulizing Crohn's Disease Not Responsive or Intolerant to Conventional Care: HYPNOTIC Pilot Study	40	Mar 2026
NCT05987852	Hyperbaric Oxygen Therapy for Ulcerative Colitis Patients Hospitalized for Moderate to Severe Flares: A Multi-Center, Randomized, Double-Blind, Sham-Controlled Trial	126	Sept 2027
NCT06148285	Hyperbaric Oxygen Therapy in Acute Ischemic Stroke Recovery	120	Dec 2026

NCT: national clinical trial.

References

1. Undersea and Hyperbaric Medical Society. Hyperbaric Oxygen Therapy Indications, 15th Edition. Best Publishing Company (North Palm Beach, FL). 2023.
2. Sadri RA, Cooper JS. Hyperbaric, complications. NCBI Bookshelf 2025; <https://www.ncbi.nlm.nih.gov/books/NBK459191/>.
3. U.S. Food and Drug Administration. Hyperbaric Oxygen Therapy: Get the Facts. 2021; <https://www.talkingaboutthescience.com/studies/FDA2013.pdf>.
4. de Smet GHJ, Kroese LF, Menon AG, et al. Oxygen therapies and their effects on wound healing. Wound Repair Regen. Aug 2017; 25(4): 591-608. PMID 28783878

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

5. Sharma R, Sharma SK, Mudgal SK, et al. Efficacy of hyperbaric oxygen therapy for diabetic foot ulcer, a systematic review and meta-analysis of controlled clinical trials. *Sci Rep*. Jan 26 2021; 11(1): 2189. PMID 33500533
6. Kranke P, Bennett MH, Martyn-St James M, et al. Hyperbaric oxygen therapy for chronic wounds. *Cochrane Database Syst Rev*. Jun 24 2015; 2015(6): CD004123. PMID 26106870
7. Elraiyah T, Tsapas A, Prutsky G, et al. A systematic review and meta-analysis of adjunctive therapies in diabetic foot ulcers. *J Vasc Surg*. Feb 2016; 63(2 Suppl): 46S-58S.e1-2. PMID 26804368
8. Buckley NA, Juurlink DN, Isbister G, et al. Hyperbaric oxygen for carbon monoxide poisoning. *Cochrane Database Syst Rev*. Apr 13 2011; 2011(4): CD002041. PMID 21491385
9. Nakajima M, Aso S, Matsui H, et al. Hyperbaric oxygen therapy and mortality from carbon monoxide poisoning: A nationwide observational study. *Am J Emerg Med*. Feb 2020; 38(2): 225-230. PMID 30797609
10. Bennett MH, Feldmeier J, Hampson NB, et al. Hyperbaric oxygen therapy for late radiation tissue injury. *Cochrane Database Syst Rev*. Apr 28 2016; 4(4): CD005005. PMID 27123955
11. Borab Z, Mirmanesh MD, Gantz M, et al. Systematic review of hyperbaric oxygen therapy for the treatment of radiation-induced skin necrosis. *J Plast Reconstr Aesthet Surg*. Apr 2017; 70(4): 529-538. PMID 28081957
12. Ravi P, Vaishnavi D, Gnanam A, et al. The role of hyperbaric oxygen therapy in the prevention and management of radiation-induced complications of the head and neck - a systematic review of literature. *J Stomatol Oral Maxillofac Surg*. Dec 2017; 118(6): 359-362. PMID 28838774
13. Savvidou OD, Kaspiris A, Bolia IK, et al. Effectiveness of Hyperbaric Oxygen Therapy for the Management of Chronic Osteomyelitis: A Systematic Review of the Literature. *Orthopedics*. Jul 01 2018; 41(4): 193-199. PMID 30035798
14. Maynor ML, Moon RE, Camporesi EM, et al. Chronic osteomyelitis of the tibia: treatment with hyperbaric oxygen and autogenous microsurgical muscle transplantation. *J South Orthop Assoc*. 1998; 7(1): 43-57. PMID 9570731
15. Davis JC, Heckman JD, DeLee JC, et al. Chronic non-hematogenous osteomyelitis treated with adjuvant hyperbaric oxygen. *J Bone Joint Surg Am*. Oct 1986; 68(8): 1210-7. PMID 3771602
16. Chen CE, Ko JY, Fu TH, et al. Results of chronic osteomyelitis of the femur treated with hyperbaric oxygen: a preliminary report. *Chang Gung Med J*. Feb 2004; 27(2): 91-7. PMID 15095953
17. Chen CE, Shih ST, Fu TH, et al. Hyperbaric oxygen therapy in the treatment of chronic refractory osteomyelitis: a preliminary report. *Chang Gung Med J*. Feb 2003; 26(2): 114-21. PMID 12718388
18. Chen CY, Lee SS, Chan YS, et al. Chronic refractory tibia osteomyelitis treated with adjuvant hyperbaric oxygen: a preliminary report. *Changgeng Yi Xue Za Zhi*. Jun 1998; 21(2): 165-71. PMID 9729650
19. Villanueva E, Bennett MH, Wasiak J, et al. Hyperbaric oxygen therapy for thermal burns. *Cochrane Database Syst Rev*. 2004; 2004(3): CD004727. PMID 15266540
20. Eskes A, Vermeulen H, Lucas C, et al. Hyperbaric oxygen therapy for treating acute surgical and traumatic wounds. *Cochrane Database Syst Rev*. Dec 16 2013; 2013(12): CD008059. PMID 24343585

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

21. Dauwe PB, Pulikkottil BJ, Lavery L, et al. Does hyperbaric oxygen therapy work in facilitating acute wound healing: a systematic review. *Plast Reconstr Surg*. Feb 2014; 133(2): 208e-215e. PMID 24469192
22. Zhou YY, Liu W, Yang YJ, et al. Use of hyperbaric oxygen on flaps and grafts in China: analysis of studies in the past 20 years. *Undersea Hyperb Med*. 2014; 41(3): 209-16. PMID 24984315
23. Freiburger JJ, Padilla-Burgos R, McGraw T, et al. What is the role of hyperbaric oxygen in the management of bisphosphonate-related osteonecrosis of the jaw: a randomized controlled trial of hyperbaric oxygen as an adjunct to surgery and antibiotics. *J Oral Maxillofac Surg*. Jul 2012; 70(7): 1573-83. PMID 22698292
24. Levett D, Bennett MH, Millar I. Adjunctive hyperbaric oxygen for necrotizing fasciitis. *Cochrane Database Syst Rev*. Jan 15 2015; 1(1): CD007937. PMID 25879088
25. Hedetoft M, Bennett MH, Hyldegaard O. Adjunctive hyperbaric oxygen treatment for necrotising soft-tissue infections: A systematic review and meta-analysis. *Diving Hyperb Med*. Mar 31 2021; 51(1): 34-43. PMID 33761539
26. Bennett MH, Lehm JP, Jepson N. Hyperbaric oxygen therapy for acute coronary syndrome. *Cochrane Database Syst Rev*. Jul 23 2015; 2015(7): CD004818. PMID 26202854
27. Bennett MH, Weibel S, Wasiak J, et al. Hyperbaric oxygen therapy for acute ischaemic stroke. *Cochrane Database Syst Rev*. Nov 12 2014; 2014(11): CD004954. PMID 25387992
28. Wang B, Yu K, Wang J, et al. Efficacy of hyperbaric oxygen combined with dual antiplatelet therapy in elderly patients with acute cerebral infarction and its impact on nerve factors. *Thromb J*. Oct 08 2025; 23(1): 93. PMID 41063286
29. Efrati S, Fishlev G, Bechor Y, et al. Hyperbaric oxygen induces late neuroplasticity in post stroke patients--randomized, prospective trial. *PLoS One*. 2013; 8(1): e53716. PMID 23335971
30. Holland NJ, Bernstein JM, Hamilton JW. Hyperbaric oxygen therapy for Bell's palsy. *Cochrane Database Syst Rev*. Feb 15 2012; 2012(2): CD007288. PMID 22336830
31. Wang F, Wang Y, Sun T, et al. Hyperbaric oxygen therapy for the treatment of traumatic brain injury: a meta-analysis. *Neurol Sci*. May 2016; 37(5): 693-701. PMID 26746238
32. Crawford C, Teo L, Yang E, et al. Is Hyperbaric Oxygen Therapy Effective for Traumatic Brain Injury? A Rapid Evidence Assessment of the Literature and Recommendations for the Field. *J Head Trauma Rehabil*. 2017; 32(3): E27-E37. PMID 27603765
33. Bennett MH, Trytko B, Jonker B. Hyperbaric oxygen therapy for the adjunctive treatment of traumatic brain injury. *Cochrane Database Syst Rev*. Dec 12 2012; 12(12): CD004609. PMID 23235612
34. Hart BB, Weaver LK, Gupta A, et al. Hyperbaric oxygen for mTBI-associated PCS and PTSD: Pooled analysis of results from Department of Defense and other published studies. *Undersea Hyperb Med*. BIMA 2019; 46(3): 353-383. PMID 31394604
35. Shahid S, Saeed H, Ali M, et al. Hyperbaric oxygen therapy (HBOT) for neurocognitive deficits following traumatic brain injury: a systematic review and meta-analysis. *Ann Med Surg (Lond)*. Nov 2025; 87(11): 7490-7498. PMID 41180753
36. mTBI mechanisms of action of HBO2 for persistent post-concussive symptoms. U.S. National Library of Medicine. *ClinicalTrials.gov*. <https://clinicaltrials.gov/ct2/show/NCT01611194>.
37. Hart BB, Wilson SH, Churchill S, et al. Extended follow-up in a randomized trial of hyperbaric oxygen for persistent post-concussive symptoms. *Undersea Hyperb Med*. BIMA 2019; 46(3): 313-327. PMID 31394601

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

38. Churchill S, Deru K, Weaver LK, et al. Adverse events and blinding in two randomized trials of hyperbaric oxygen for persistent post-concussive symptoms. *Undersea Hyperb Med. BIMA* 2019; 46(3): 331-340. PMID 31394602
39. Weaver LK, Churchill S, Wilson SH, et al. A composite outcome for mild traumatic brain injury in trials of hyperbaric oxygen. *Undersea Hyperb Med. BIMA* 2019; 46(3): 341-352. PMID 31394603
40. Hyperbaric oxygen therapy (HBO2) for persistent post-concussive symptoms after mild traumatic brain injury (mTBI) (HOPPS). U.S. National Library of Medicine. ClinicalTrials.gov. <https://clinicaltrials.gov/ct2/show/NCT01306968>. Updated September 5, 2014.
41. Harch PG, Andrews SR, Rowe CJ, et al. Hyperbaric oxygen therapy for mild traumatic brain injury persistent postconcussion syndrome: a randomized controlled trial. *Med Gas Res.* 2020; 10(1): 8-20. PMID 32189664
42. Hadanny A, Catalogna M, Yaniv S, et al. Hyperbaric oxygen therapy in children with post-concussion syndrome improves cognitive and behavioral function: a randomized controlled trial. *Sci Rep.* Sep 23 2022; 12(1): 15233. PMID 36151105
43. Zhang Z, Li Z, Li S, et al. Hyperbaric oxygen for moderate-to-severe traumatic brain injury: outcomes 5-8 years after injury. *Med Gas Res.* Mar 01 2025; 15(1): 156-163. PMID 39324933
44. You JH, Jiang JL, He WB, et al. Addition of hyperbaric oxygen therapy versus usual care alone for inflammatory bowel disease: A systematic review and meta-analysis. *Heliyon.* Oct 2022; 8(10): e11007. PMID 36276722
45. McCurdy J, Siw KCK, Kandel R, et al. The Effectiveness and Safety of Hyperbaric Oxygen Therapy in Various Phenotypes of Inflammatory Bowel Disease: Systematic Review With Meta-analysis. *Inflamm Bowel Dis.* Mar 30 2022; 28(4): 611-621. PMID 34003289
46. Dulai PS, Buckey JC, Raffals LE, et al. Hyperbaric oxygen therapy is well tolerated and effective for ulcerative colitis patients hospitalized for moderate-severe flares: a phase 2A pilot multi-center, randomized, double-blind, sham-controlled trial. *Am J Gastroenterol.* Oct 2018; 113(10): 1516-1523. PMID 29453383
47. Dulai PS, Raffals LE, Hudesman D, et al. A phase 2B randomised trial of hyperbaric oxygen therapy for ulcerative colitis patients hospitalised for moderate to severe flares. *Aliment Pharmacol Ther.* Sep 2020; 52(6): 955-963. PMID 32745306
48. Pagoldh M, Hultgren E, Arnell P, et al. Hyperbaric oxygen therapy does not improve the effects of standardized treatment in a severe attack of ulcerative colitis: a prospective randomized study. *Scand J Gastroenterol.* Sep 2013; 48(9): 1033-40. PMID 23879825
49. Bennett MH, Kertesz T, Perleth M, et al. Hyperbaric oxygen for idiopathic sudden sensorineural hearing loss and tinnitus. *Cochrane Database Syst Rev.* Oct 17 2012; 10(10): CD004739. PMID 23076907
50. Rhee TM, Hwang D, Lee JS, et al. Addition of Hyperbaric Oxygen Therapy vs Medical Therapy Alone for Idiopathic Sudden Sensorineural Hearing Loss: A Systematic Review and Meta-analysis. *JAMA Otolaryngol Head Neck Surg.* Dec 01 2018; 144(12): 1153-1161. PMID 30267033
51. Joshua TG, Ayub A, Wijesinghe P, et al. Hyperbaric Oxygen Therapy for Patients With Sudden Sensorineural Hearing Loss: A Systematic Review and Meta-analysis. *JAMA Otolaryngol Head Neck Surg.* Jan 01 2022; 148(1): 5-11. PMID 34709348

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

52. Eryigit B, Ziylan F, Yaz F, et al. The effectiveness of hyperbaric oxygen in patients with idiopathic sudden sensorineural hearing loss: a systematic review. *Eur Arch Otorhinolaryngol.* Dec 2018; 275(12): 2893-2904. PMID 30324404
53. Cavaliere M, De Luca P, Scarpa A, et al. Combination of Hyperbaric Oxygen Therapy and Oral Steroids for the Treatment of Sudden Sensorineural Hearing Loss: Early or Late?. *Medicina (Kaunas).* Oct 10 2022; 58(10). PMID 36295581
54. Bennett M, Best TM, Babul S, et al. Hyperbaric oxygen therapy for delayed onset muscle soreness and closed soft tissue injury. *Cochrane Database Syst Rev.* Oct 19 2005; 2005(4): CD004713. PMID 16235376
55. Tu P, Halili X, Zhang S, et al. The effectiveness of hyperbaric oxygen therapy in children and adolescents and with autism spectrum disorders: A systematic review and meta-analysis. *Prog Neuropsychopharmacol Biol Psychiatry.* Mar 20 2025; 137: 111257. PMID 39826608
56. Xiong T, Chen H, Luo R, et al. Hyperbaric oxygen therapy for people with autism spectrum disorder (ASD). *Cochrane Database Syst Rev.* Oct 13 2016; 10(10): CD010922. PMID 27737490
57. Sampanthavivat M, Singkhwa W, Chaiyakul T, et al. Hyperbaric oxygen in the treatment of childhood autism: a randomised controlled trial. *Diving Hyperb Med.* Sep 2012; 42(3): 128-33. PMID 22987458
58. Rizzato A, D'Alessandro N, Berenci E, et al. Effect of mild hyperbaric oxygen therapy on children diagnosed with autism. *Undersea Hyperb Med.* 2018; 45(6): 639-645. PMID 31158930
59. Lacey DJ, Stolfi A, Pilati LE. Effects of hyperbaric oxygen on motor function in children with cerebral palsy. *Ann Neurol.* Nov 2012; 72(5): 695-703. PMID 23071074
60. Collet JP, Vanasse M, Marois P, et al. Hyperbaric oxygen for children with cerebral palsy: a randomised multicentre trial. HBO-CP Research Group. *Lancet.* Feb 24 2001; 357(9256): 582-6. PMID 11558483
61. Long Y, Tan J, Nie Y, et al. Hyperbaric oxygen therapy is safe and effective for the treatment of sleep disorders in children with cerebral palsy. *Neurol Res.* Mar 2017; 39(3): 239-247. PMID 28079475
62. Xiao Y, Wang J, Jiang S, et al. Hyperbaric oxygen therapy for vascular dementia. *Cochrane Database Syst Rev.* Jul 11 2012; 2012(7): CD009425. PMID 22786527
63. Villeirs L, Tailly T, Ost P, et al. Hyperbaric oxygen therapy for radiation cystitis after pelvic radiotherapy: Systematic review of the recent literature. *Int J Urol.* Feb 2020; 27(2): 98-107. PMID 31617263
64. Gothard L, Haviland J, Bryson P, et al. Randomised phase II trial of hyperbaric oxygen therapy in patients with chronic arm lymphoedema after radiotherapy for cancer. *Radiother Oncol.* Oct 2010; 97(1): 101-7. PMID 20605648
65. Oscarsson N, Müller B, Rosén A, et al. Radiation-induced cystitis treated with hyperbaric oxygen therapy (RICH-ART): a randomised, controlled, phase 2-3 trial. *Lancet Oncol.* Nov 2019; 20(11): 1602-1614. PMID 31537473
66. Camporesi EM, Vezzani G, Bosco G, et al. Hyperbaric oxygen therapy in femoral head necrosis. *J Arthroplasty.* Sep 2010; 25(6 Suppl): 118-23. PMID 20637561
67. Bennett MH, French C, Schnabel A, et al. Normobaric and hyperbaric oxygen therapy for the treatment and prevention of migraine and cluster headache. *Cochrane Database Syst Rev.* Dec 28 2015; 2015(12): CD005219. PMID 26709672

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

68. Peng Z, Wang S, Huang X, et al. Effect of hyperbaric oxygen therapy on patients with herpes zoster. *Undersea Hyperb Med.* 2012; 39(6): 1083-7. PMID 23342765
69. Efrati S, Golan H, Bechor Y, et al. Hyperbaric oxygen therapy can diminish fibromyalgia syndrome--prospective clinical trial. *PLoS One.* 2015; 10(5): e0127012. PMID 26010952
70. Yildiz S, Kiralp MZ, Akin A, et al. A new treatment modality for fibromyalgia syndrome: hyperbaric oxygen therapy. *J Int Med Res.* 2004; 32(3): 263-7. PMID 15174219
71. Bennett M, Heard R. Hyperbaric oxygen therapy for multiple sclerosis. *CNS Neurosci Ther.* Apr 2010; 16(2): 115-24. PMID 20415839
72. Bennett M, Feldmeier J, Smee R, et al. Hyperbaric oxygenation for tumour sensitisation to radiotherapy. *Cochrane Database Syst Rev.* Oct 19 2005; (4): CD005007. PMID 16235387
73. Bennett MH, Feldmeier J, Smee R, et al. Hyperbaric oxygenation for tumour sensitisation to radiotherapy. *Cochrane Database Syst Rev.* Apr 18 2012; 2012(4): CD005007. PMID 22513926
74. Heys SD, Smith IC, Ross JA, et al. A pilot study with long term follow up of hyperbaric oxygen pretreatment in patients with locally advanced breast cancer undergoing neo-adjuvant chemotherapy. *Undersea Hyperb Med.* 2006; 33(1): 33-43. PMID 16602255
75. Hyman SL, Levy SE, Myers SM, et al. Identification, Evaluation, and Management of Children With Autism Spectrum Disorder. *Pediatrics.* Jan 2020; 145(1). PMID 31843864
76. Singh S, Loftus EV, Limketkai BN, et al. AGA Living Clinical Practice Guideline on Pharmacological Management of Moderate-to-Severe Ulcerative Colitis. *Gastroenterology.* Dec 2024; 167(7): 1307-1343. PMID 39572132
77. Scott FI, Ananthakrishnan AN, Click B, et al. AGA Living Clinical Practice Guideline on the Pharmacologic Management of Moderate-to-Severe Crohn's Disease. *Gastroenterology.* Dec 2025; 169(7): 1397-1448. PMID 41274746
78. Rubin DT, Ananthakrishnan AN, Siegel CA, et al. ACG Clinical Guideline Update: Ulcerative Colitis in Adults. *Am J Gastroenterol.* Jun 03 2025; 120(6): 1187-1224. PMID 40701556
79. Lichtenstein GR, Loftus EV, Afzali A, et al. ACG Clinical Guideline: Management of Crohn's Disease in Adults. *Am J Gastroenterol.* Jun 03 2025; 120(6): 1225-1264. PMID 40701562
80. Powers WJ, Rabinstein AA, Ackerson T, et al. Guidelines for the Early Management of Patients With Acute Ischemic Stroke: 2019 Update to the 2018 Guidelines for the Early Management of Acute Ischemic Stroke: A Guideline for Healthcare Professionals From the American Heart Association/American Stroke Association. *Stroke.* Dec 2019; 50(12): e344-e418. PMID 31662037
81. Kernan WN, Viera AJ, Billinger SA, et al. Primary Care of Adult Patients After Stroke: A Scientific Statement From the American Heart Association/American Stroke Association. *Stroke.* Aug 2021; 52(9): e558-e571. PMID 34261351
82. Department of Veterans Affairs. VA/DoD clinical practice guideline for the management of stroke rehabilitation. 2024. https://www.healthquality.va.gov/HEALTHQUALITY/guidelines/Rehab/stroke/VADOD-2024-Stroke-Rehab-CPG-Full-CPG_final_508.pdf
83. ACS Committee on Trauma. Best Practice Guidelines: The Management Of Traumatic Brain Injury. 2024. <https://www.facs.org/media/vgfgjpfk/best-practices-guidelines-traumatic-brain-injury.pdf>.
84. Carney N, Totten AM, O'Reilly C, et al. Guidelines for the management of severe traumatic brain injury, fourth edition. *Neurosurgery.* 2017;80(1):6-15.

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

85. Department of Veterans Affairs. VA/DoD clinical practice guideline for the management and rehabilitation of post-acute mild traumatic brain injury. 2021. https://www.healthquality.va.gov/HEALTHQUALITY/guidelines/Rehab/mtbi/VADODmTBI_CPGFinal508.pdf.
86. Hingorani A, LaMuraglia GM, Henke P, et al. The management of diabetic foot: A clinical practice guideline by the Society for Vascular Surgery in collaboration with the American Podiatric Medical Association and the Society for Vascular Medicine. J Vasc Surg. Feb 2016; 63(2 Suppl): 3S-21S. PMID 26804367
87. Huang ET, Mansouri J, Murad MH, et al. A clinical practice guideline for the use of hyperbaric oxygen therapy in the treatment of diabetic foot ulcers. Undersea Hyperb Med. 2015; 42(3): 205-47. PMID 26152105
88. Centers for Medicare and Medicaid Services (CMS). National Coverage Determination (NCD) for Hyperbaric Oxygen Therapy (20.29). 2017; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?ncid=12&ver=3>.
89. U.S. Food & Drug Administration. Hyperbaric Oxygen Therapy: Get the Facts. October 1, 2024.

Policy History

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

08/19/2003	Medical Policy Committee review
08/25/2003	Managed Care Advisory Council approval
08/10/2004	Medical Director review
08/17/2004	Medical Policy Committee review
08/31/2004	Medical Director review
09/21/2004	Medical Policy Committee review. Format revision and the following changes to coverage eligibility: Retinal artery insufficiency deleted from list of covered conditions. Prophylactic pre- and post-treatment for patients undergoing dental surgery of a radiated jaw added to the list of covered conditions.
09/27/2004	Managed Care Advisory Council approval
10/10/2005	Medical Director review
10/18/2005	Medical Policy Committee review. Format revision. Clinical criteria revision. HBO2 for acute coronary syndromes and as an adjunct to percutaneous coronary interventions added to investigational indications. Coverage eligibility changes. Refractory mycoses, mucomycosis, actinomycosis and candidiobolus coronato changed from eligible for coverage to investigational. <i>Effective date of policy will reflect 60 day period following the notification of providers that coverage eligibility has changed.</i>
10/27/2005	Managed Care Advisory Council approval
01/10/2007	Medical Director review
01/17/2007	Medical Policy Committee approval. Format revision. Coverage eligibility unchanged.
12/05/2007	Medical Director review

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

12/19/2007	Medical Policy Committee approval. Coverage eligibility unchanged. Added autism as investigational.
12/03/2008	Medical Director review
12/17/2008	Medical Policy Committee approval. No change to coverage.
07/02/2009	Medical Director review
07/22/2009	Medical Policy Committee approval. No change to coverage eligibility.
07/01/2010	Medical Policy Committee approval
07/21/2010	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
07/07/2011	Medical Policy Committee review
07/20/2011	Medical Policy Implementation Committee approval. Changed chronic refractory osteomyelitis from investigational to eligible for coverage.
06/28/2012	Medical Policy Committee review
07/27/2012	Medical Policy Implementation Committee approval. Acute surgical and traumatic wounds, idiopathic femoral neck necrosis, chronic arm lymphedema following radiotherapy for cancer, radiation-induced injury in the head and neck added as investigational. Changed chronic diabetic wounds to chronic non-diabetic wounds as an investigational indication, since chronic diabetic wounds are covered.
08/01/2013	Medical Policy Committee review
08/21/2013	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
08/07/2014	Medical Policy Committee review
08/20/2014	Medical Policy Implementation Committee approval. Added vascular dementia, herpes zoster, motor dysfunction associated with stroke, and bisphosphonate-related osteonecrosis of the jaw as investigational.
10/02/2014	Medical Policy Committee review
10/15/2014	Medical Policy Implementation Committee approval. Clarified soft tissue radiation necrosis. Radiation myelitis, cystitis, enteritis or proctitis was removed from investigational section.
01/01/2015	Coding Update
08/03/2015	Coding update: ICD10 Diagnosis code section added; ICD9 Procedure code section removed.
12/03/2015	Medical Policy Committee review
12/16/2015	Medical Policy Implementation Committee approval. Indications added to INV statement: Fibromyalgia, mental illness (ie, posttraumatic stress disorder, generalized anxiety disorder or depression), and Inflammatory bowel disease (Crohn disease or ulcerative colitis).
12/01/2016	Medical Policy Committee review
12/21/2016	Medical Policy Implementation Committee approval. No change to coverage.
01/01/2017	Coding update: Removing ICD-9 Diagnosis Codes
12/07/2017	Medical Policy Committee review
12/20/2017	Medical Policy Implementation Committee approval. No change to coverage.
12/06/2018	Medical Policy Committee review
12/19/2018	Medical Policy Implementation Committee approval. No change to coverage.

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

01/01/2019	Coding update
12/05/2019	Medical Policy Committee review
12/11/2019	Medical Policy Implementation Committee approval. Added coverage for central retinal artery occlusion when treatment is initiated within 24 hours after initial vision loss and acute gas embolism. Brown recluse spider bites changed to investigational. Coding update.
04/02/2020	Medical Policy Committee review
04/08/2020	Medical Policy Implementation Committee approval. Removed radiation-induced injury in the head and neck, except as noted earlier in the medically necessary statement from the investigational section.
04/01/2021	Medical Policy Committee review
04/14/2021	Medical Policy Implementation Committee approval. No change to coverage.
04/07/2022	Medical Policy Committee review
04/13/2022	Medical Policy Implementation Committee approval. No change to coverage.
04/06/2023	Medical Policy Committee review
04/12/2023	Medical Policy Implementation Committee approval. Title changed from “Hyperbaric Oxygen Pressurization (HBO) to “Hyperbaric Oxygen Therapy”. Policy statements changed from hyperbaric oxygen pressurization to hyperbaric oxygen therapy. Replaced the heading and coverage statements from “When Services Are Eligible for Coverage” to “When Services May Be Eligible for Coverage” to include criteria. Revised criteria for compromised skin grafts and added information for compromised skin flaps to the criteria. Added information on idiopathic sudden sensorial hearing loss to the criteria. Removed idiopathic sudden sensorial hearing loss from the investigational conditions.
04/04/2024	Medical Policy Committee review
04/10/2024	Medical Policy Implementation Committee approval. Added “not meeting criteria specified in the eligible for coverage statement” to the Acute surgical and traumatic wounds investigational statement. Added “flaps” and additional bullet “visible ischemic changes such as pallor, mottling, or frank necrosis of the overlying skin” to patient selection criteria. Added to deny investigational when criteria are not met.
11/07/2024	Medical Policy Committee review
11/13/2024	Medical Policy Implementation Committee approval. Central retinal artery occlusion added to eligible for coverage statement. Also changed Idiopathic sudden sensorineural hearing loss (ISSHL) in individuals with a loss of 30 decibels or more or hearing loss at least 61 decibels of pure tone thresholds when HBOT is combined with steroid therapy (systemic or intra-tympanic) and initiated within 1 month of onset of SSNHL. Added “Peripheral arterial insufficiency related to non-healing arterial insufficiency ulcers with no measurable signs of healing after 30 days of an adequate course of standard wound therapy” as eligible. Under the investigational section, “Acute peripheral artery insufficiency (outside of other listed medically necessary indications involving arterial insufficiency) (see Policy Guidelines)” was added and “Retinal artery insufficiency, acute” was deleted.
04/03/2025	Medical Policy Committee review
04/09/2025	Medical Policy Implementation Committee approval. FDA information updated.

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

04/02/2026 Medical Policy Committee review

04/10/2026 Medical Policy Implementation Committee approval. Some of policy guidelines pulled into criteria section.

Added “Treatment of infection (if present)” to the coverage criteria.

The following was changed for clarification: “Idiopathic sudden sensorineural hearing loss (ISSHL) in individuals with one of the following:

- Sudden hearing loss of 30 decibels or more; OR
- Hearing loss at least 61 decibels of pure tone thresholds when HBOT is combined with steroid therapy (systemic or intra-tympanic) and initiated within 1 month of onset of SSNHL”

Next Scheduled Review Date: 04/2027

Coding

The five character codes included in the Louisiana Blue Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)†, copyright 2025 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Louisiana Blue Medical Policy Coverage Guidelines is with Louisiana Blue and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Louisiana Blue Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Louisiana Blue Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	99183
HCPCS	A4575, G0277
ICD-10 Diagnosis	All related Diagnoses

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
 1. Consultation with technology evaluation center(s);
 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
 3. Reference to federal regulations.

****Medically Necessary (or “Medical Necessity”)** - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

For these purposes, “nationally accepted standards of medical practice” means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: If the Patient’s health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

Hyperbaric Oxygen Therapy (HBO)

Policy # 00070

Original Effective Date: 08/25/2003

Current Effective Date: 05/01/2026

NOTICE: Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage.

NOTICE: If an authorization for an ongoing course of treatment has been provided to a member and the member changes from one health plan to another health plan (e.g., a member moves from carrier A to Louisiana Blue), Louisiana Blue may honor the previous health plan's authorization for the same service under the same type of in-network benefit for a 90-day transition period. Documentation of the authorization for the ongoing course of treatment from the previous health plan must be provided to us by the member or their provider and the services provided for the course of treatment must otherwise be a covered service under the Louisiana Blue health plan. This provision does not apply to medications covered under the plan's pharmacy benefit.