

Image-Guided Minimally Invasive Decompression for Spinal Stenosis

Policy # 00278

Original Effective Date: 11/16/2010

Current Effective Date: 12/01/2025

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Note: Interspinous and Interlaminar Stabilization/Distraction Devices (Spacers) is addressed separately in medical policy 00221.

Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers image-guided minimally invasive spinal decompression to be **investigational**.*

Background/Overview

Spinal Stenosis

In spinal stenosis, the space around the spinal cord narrows, compressing the spinal cord and its nerve roots. The goal of surgical treatment is to "decompress" the spinal cord and/or nerve roots.

The most common symptoms of lumbar spinal stenosis are back pain with neurogenic claudication (ie, pain, numbness, weakness) in the legs that worsens with standing or walking and is alleviated by sitting or leaning forward. Compression of neural elements generally occurs from a combination of degenerative changes, including ligamentum flavum hypertrophy, bulging of the intervertebral disc, and facet thickening with arthropathy. Spinal stenosis is often linked to age-related changes in disc height and arthritis of the facet joints. Lumbar spinal stenosis is among the most common reasons for back surgery and the most common reason for lumbar spine surgery in adults over the age of 65.

The most common symptoms of cervical/thoracic spinal stenosis are neck pain and radiculopathy of the shoulder and arm. The most common cause of cervical radiculopathy is degenerative changes, including disc herniation.

Image-Guided Minimally Invasive Decompression for Spinal Stenosis

Policy # 00278

Original Effective Date: 11/16/2010

Current Effective Date: 12/01/2025

Treatment

Conventional Posterior Decompression Surgery

For patients with lumbar spinal stenosis, surgical laminectomy has established benefits in reducing pain and improving quality of life.

For patients with cervical or thoracic stenosis, surgical treatment includes discectomy or foraminal decompression.

A systematic review by Chou et al (2009) assessed surgery for back pain; it was commissioned by the American Pain Society and conducted by an evidence-based center. Four higher quality randomized trials were reviewed; they compared surgery with nonsurgical therapy for spinal stenosis, including 2 studies from the multicenter Spine Patient Outcomes Research Trial that evaluated laminectomy for spinal stenosis (specifically with or without degenerative spondylolisthesis). All 4 studies found that initial decompressive surgery (laminectomy) was slightly to moderately superior to initial nonsurgical therapy (eg, average 8- to 18-point differences on the 36-Item Short-Form Health Survey and Oswestry Disability Index). However, there was insufficient evidence to determine the optimal adjunctive surgical methods for laminectomy (ie, with or without fusion, instrumented vs noninstrumented fusion) in patients with or without degenerative spondylolisthesis. The Spine Patient Outcomes Research Trial continues to be referenced as the highest quality evidence published on decompressive surgery.

Less invasive surgical procedures include open laminotomy and microendoscopic laminotomy. In general, the literature comparing surgical procedures is limited. The literature has suggested that less invasive surgical decompression may reduce perioperative morbidity without impairing long-term outcomes when performed in appropriately selected patients. Posterior decompressive surgical procedures include: decompressive laminectomy, hemilaminotomy and laminotomy, and microendoscopic decompressive laminotomy.

Decompressive laminectomy, the classic treatment for lumbar spinal stenosis, unroofs the spinal canal by extensive resection of posterior spinal elements, including the lamina, spinous processes, portions of the facet joints, ligamentum flavum, and the interspinous ligaments. Wide muscular dissection and retraction is needed to achieve adequate surgical visualization. The extensive resection and injury to the posterior spine and supporting musculature can lead to instability with significant morbidity, both postoperatively and longer term. Spinal fusion, performed at the same time as laminectomy or after symptoms have developed, may be required to reduce resultant instability. Laminectomy may also be used for extensive multilevel decompression.

Hemilaminotomy and laminotomy, sometimes termed laminoforaminotomy, are less invasive than laminectomy. These procedures focus on the interlaminar space, where most of the pathologic changes are concentrated, minimizing resection of the stabilizing posterior spine. A laminotomy

Image-Guided Minimally Invasive Decompression for Spinal Stenosis

Policy # 00278

Original Effective Date: 11/16/2010

Current Effective Date: 12/01/2025

typically removes the inferior aspect of the cranial lamina, superior aspect of the subjacent lamina, ligamentum flavum, and the medial aspect of the facet joint. Unlike laminectomy, laminotomy does not disrupt the facet joints, supra- and interspinous ligaments, a major portion of the lamina, or the muscular attachments. Muscular dissection and retraction are required to achieve adequate surgical visualization.

Microendoscopic decompressive laminotomy, similar to laminotomy, uses endoscopic visualization. The position of the tubular working channel is confirmed by fluoroscopic guidance, and serial dilators are used to dilate the musculature and expand the fascia. For microendoscopic decompressive laminotomy, an endoscopic curette, rongeur, and drill are used for the laminotomy, facetectomy, and foraminotomy. The working channel may be repositioned from a single incision for multilevel and bilateral dissections.

Image-Guided Minimally Invasive Spinal Decompression

Posterior decompression for spinal stenosis has been evolving toward increasingly minimally invasive procedures in an attempt to reduce postoperative morbidity and spinal instability. Unlike conventional surgical decompression, the percutaneous mild^{®‡} decompressive procedure is performed solely under fluoroscopic guidance (eg, without endoscopic or microscopic visualization of the work area). This procedure is indicated for central stenosis only, without the capability of addressing nerve root compression or disc herniation, should either be required.

Percutaneous image-guided minimally invasive spinal decompression using a specially designed tool kit (mild^{®‡}) has been proposed as an ultra-minimally invasive treatment of central lumbar spinal stenosis. In this procedure, the epidural space is filled with contrast medium under fluoroscopic guidance. Using a 6-gauge cannula clamped in place with a back plate, single-use tools (portal cannula, surgical guide, bone rongeur, tissue sculpter, trocar) are used to resect thickened ligamentum flavum and small pieces of lamina. The tissue and bone sculpting is conducted entirely under fluoroscopic guidance, with contrast media added throughout the procedure to aid visualization of the decompression. The process is repeated on the opposite side for bilateral decompression of the central canal. The devices are not intended for use near the lateral neural elements and are contraindicated for disc procedures.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

In 2006, the X-Sten MILD Tool Kit (now the mild^{®‡} device kit, X-Sten Corp. renamed Vertos Medical) was cleared for marketing by the U.S. Food and Drug Administration through the 510(k) process for treatment of various spinal conditions. This set of specialized surgical instruments is used to perform percutaneous lumbar decompressive procedures.

Image-Guided Minimally Invasive Decompression for Spinal Stenosis

Policy # 00278

Original Effective Date: 11/16/2010

Current Effective Date: 12/01/2025

Vertos's mild[®]‡ instructions state that the device is not intended for disc procedures but rather for tissue resection at the perilaminar space, within the interlaminar space, and at the ventral aspect of the lamina. The device is not intended for use near the lateral neural elements and remains dorsal to the dura using image guidance and anatomic landmarks.

Food and Drug Administration product code: HRX.

Rationale/Source

This medical policy was developed through consideration of peer-reviewed medical literature generally recognized by the relevant medical community, U.S. Food and Drug Administration approval status, nationally accepted standards of medical practice and accepted standards of medical practice in this community, technology evaluation centers, reference to regulations, other plan medical policies, and accredited national guidelines.

Image-guided minimally invasive decompression describes a percutaneous procedure for decompression of the central spinal canal in patients with spinal stenosis and hypertrophy of the ligamentum flavum. In this procedure, a specialized cannula and surgical tools (mild[®])‡ are used under fluoroscopic guidance for bone and tissue sculpting near the spinal canal. Image-guided minimally invasive lumbar decompression is proposed as an alternative to existing posterior decompression procedures.

Summary of Evidence

For individuals who have lumbar spinal stenosis who receive image-guided minimally invasive lumbar decompression (MILD), the evidence includes a large, randomized controlled trial (RCT) (N=302), a second RCT (N=138) comparing MILD to non-surgical conventional medical management (CMM), a systematic review that included a small RCT (N=38), and a number of prospective and retrospective cohort studies and case series. Relevant outcomes are symptoms, functional outcomes, health status measures, and treatment-related morbidity. The largest RCT (MIDAS Encore) compared image-guided MILD with epidural steroid injections (control) in patients who had ligamentum flavum hypertrophy and who failed conservative therapy. Results suggested reductions in pain and improvements in function scores in the image-guided minimally invasive lumbar decompression group versus the control group. The trial was unblinded, and there is evidence of differing expectations and follow-up in the 2 groups, suggesting a high risk of bias. The MOTION RCT compared MILD as first-line therapy in combination with nonsurgical CMM to CMM alone in 138 individuals with lumbar spinal stenosis. At 1-year follow-up, patients in the MILD + CMM group experienced a 16.1-point composite Oswestry Disability Index (ODI) mean improvement (the primary outcome), compared with a 2.0-point mean improvement for participants in the CMM-alone arm (p<.001). A major limitation of this trial was the wide variation in CMM interventions received by individuals in both the intervention and control groups; for example, 38.7% of individuals in the CMM alone group received no interventional therapy. Lack of blinding

Image-Guided Minimally Invasive Decompression for Spinal Stenosis

Policy # 00278

Original Effective Date: 11/16/2010

Current Effective Date: 12/01/2025

and follow-up for only 12 months were additional limitations. The available evidence is insufficient to determine the efficacy of MILD compared with placebo, open decompression, or conservative treatment. Well-designed and conducted trials with relevant control groups could provide greater certainty on the risks and benefits of this procedure. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have cervical or thoracic spinal stenosis who receive image-guided minimally invasive spinal decompression, no evidence was identified. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Supplemental Information

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

Lumbar Spinal Stenosis Consensus Group

In 2018, the Lumbar Spinal Stenosis Consensus Group, composed of a panel of nationally recognized spine experts, convened to evaluate the available literature and develop guidelines for minimally invasive spine treatment (MIST Guidelines). Based on a systematic review of the available literature on percutaneous image-guided lumbar decompression, the consensus committee determined there is sufficient support to warrant Level I evidence (Grade A, Level I, Consensus strong). Grade A evidence is defined as "extremely recommendable (good evidence that the measure is effective and that benefits outweigh the harms.)"

North American Spine Society

In 2011, the North American Spine Society revised clinical practice guidelines on the diagnosis and treatment of degenerative lumbar spinal stenosis. Treatment recommendations included:

- Interlaminar epidural steroid injection for short-term (6 weeks to 6 months) symptom relief in patients with neurogenic claudication or radiculopathy; however, there is conflicting evidence regarding long-term efficacy. (Grade of Recommendation: B)
- A multiple injection regimen of radiographically-guided transforaminal epidural steroid injection or caudal injection for medium-term relief of pain. (Grade of Recommendation: C)
- Decompressive surgery to improve outcomes in patients with moderate to severe symptoms of lumbar spinal stenosis. (Grade of Recommendation: B)

No specific recommendations on percutaneous image-guided lumbar decompression were provided.

Image-Guided Minimally Invasive Decompression for Spinal Stenosis

Policy # 00278

Original Effective Date: 11/16/2010

Current Effective Date: 12/01/2025

2. Chou R, Loeser JD, Owens DK, et al. Interventional therapies, surgery, and interdisciplinary rehabilitation for low back pain: an evidence-based clinical practice guideline from the American Pain Society. *Spine (Phila Pa 1976)*. May 01 2009; 34(10): 1066-77. PMID 19363457
3. Weinstein JN, Lurie JD, Tosteson TD, et al. Surgical versus nonsurgical treatment for lumbar degenerative spondylolisthesis. *N Engl J Med*. May 31 2007; 356(22): 2257-70. PMID 17538085
4. Weinstein JN, Tosteson TD, Lurie JD, et al. Surgical versus nonsurgical therapy for lumbar spinal stenosis. *N Engl J Med*. Feb 21 2008; 358(8): 794-810. PMID 18287602
5. Staats PS, Benyamin RM, McDonnell F, et al. MiDAS ENCORE: Randomized Controlled Clinical Trial Report of 6-Month Results. *Pain Physician*. Feb 2016; 19(2): 25-38. PMID 26815247
6. Benyamin RM, Staats PS, MiDAS Encore I. MILD® Is an Effective Treatment for Lumbar Spinal Stenosis with Neurogenic Claudication: MiDAS ENCORE Randomized Controlled Trial. *Pain Physician*. May 2016; 19(4): 229-42. PMID 27228511
7. Staats PS, Chafin TB, Golovac S, et al. Long-Term Safety and Efficacy of Minimally Invasive Lumbar Decompression Procedure for the Treatment of Lumbar Spinal Stenosis With Neurogenic Claudication: 2-Year Results of MiDAS ENCORE. *Reg Anesth Pain Med*. Oct 2018; 43(7): 789-794. PMID 30199512
8. Deer TR, Costandi SJ, Washabaugh E, et al. The MOTION Study: A Randomized Controlled Trial with Objective Real-World Outcomes for Lumbar Spinal Stenosis Patients Treated with the mild® Procedure: One-Year Results. *Pain Med*. Apr 08 2022; 23(4): 625-634. PMID 35167700
9. Deer TR, Chafin TB, Costandi SJ, et al. The MOTION study: Two-year results of a real-world randomized controlled trial of the mild® procedure for treatment of lumbar spinal stenosis. *Pain Pract*. Jan 2024; 24(1): 109-119. PMID 37661347
10. Kreiner DS, MacVicar J, Duszynski B, et al. The mild® procedure: a systematic review of the current literature. *Pain Med*. Feb 2014; 15(2): 196-205. PMID 24308292
11. Brown LL. A double-blind, randomized, prospective study of epidural steroid injection vs. the mild® procedure in patients with symptomatic lumbar spinal stenosis. *Pain Pract*. Jun 2012; 12(5): 333-41. PMID 22272730
12. Chopko BW. Long-term results of percutaneous lumbar decompression for LSS: two-year outcomes. *Clin J Pain*. Nov 2013; 29(11): 939-43. PMID 23446067
13. Deer TR, Grider JS, Pope JE, et al. The MIST Guidelines: The Lumbar Spinal Stenosis Consensus Group Guidelines for Minimally Invasive Spine Treatment. *Pain Pract*. Mar 2019; 19(3): 250-274. PMID 30369003
14. North American Spine Society (NASS). Evidence-Based Clinical Guidelines for Multidisciplinary Spine Care: Diagnosis and Treatment of Degenerative Lumbar Spinal Stenosis. 2011; <https://www.spine.org/Portals/0/Assets/Downloads/ResearchClinicalCare/Guidelines/LumbarStenosis.pdf>.

Image-Guided Minimally Invasive Decompression for Spinal Stenosis

Policy # 00278

Original Effective Date: 11/16/2010

Current Effective Date: 12/01/2025

15. Centers for Medicare & Medicaid Services. National Coverage Determination (NCD) for Percutaneous Image-guided Lumbar Decompression for Lumbar Spinal Stenosis (2020)
<https://www.cms.gov/medicare-coverage-database/view/ncd.aspx?ncdid=358&ncdver=2>.

Policy History

Original Effective Date: 11/16/2010

Current Effective Date: 12/01/2025

11/04/2010	Medical Policy Committee review
11/16/2010	Medical Policy Implementation Committee approval. New policy.
11/03/2011	Medical Policy Committee review.
11/16/2011	Medical Policy Implementation Committee.
12/06/2011	Posted as Archived policy.
03/05/2012	Re-posted as active policy.
11/01/2012	Medical Policy Committee review.
11/28/2012	Medical Policy Implementation Committee. No change to coverage.
11/07/2013	Medical Policy Committee review.
11/20/2013	Medical Policy Implementation Committee. No change to coverage.
11/06/2014	Medical Policy Committee review.
11/21/2014	Medical Policy Implementation Committee. No change to coverage.
08/03/2015	Coding update: ICD10 Diagnosis code section added; ICD9 Procedure code section removed.
10/29/2015	Medical Policy Committee review.
11/16/2015	Medical Policy Implementation Committee. No change to coverage.
11/03/2016	Medical Policy Committee review.
11/16/2016	Medical Policy Implementation Committee. No change to coverage.
11/02/2017	Medical Policy Committee review.
11/15/2017	Medical Policy Implementation Committee. Policy statement changed from “lumbar” to “spinal” to include cervical/thoracic decompression. “Lumbar” removed from title.
11/08/2018	Medical Policy Committee review.
11/21/2018	Medical Policy Implementation Committee. No change to coverage.
11/07/2019	Medical Policy Committee review.
11/13/2019	Medical Policy Implementation Committee. No change to coverage.
11/05/2020	Medical Policy Committee review.
11/11/2020	Medical Policy Implementation Committee. No change to coverage.
11/04/2021	Medical Policy Committee review.
11/10/2021	Medical Policy Implementation Committee. No change to coverage.
11/03/2022	Medical Policy Committee review.
11/09/2022	Medical Policy Implementation Committee. No change to coverage.
11/02/2023	Medical Policy Committee review.

Image-Guided Minimally Invasive Decompression for Spinal Stenosis

Policy # 00278

Original Effective Date: 11/16/2010

Current Effective Date: 12/01/2025

11/08/2023 Medical Policy Implementation Committee. No change to coverage.
11/07/2024 Medical Policy Committee review.
11/13/2024 Medical Policy Implementation Committee. Coverage eligibility unchanged.
11/06/2025 Medical Policy Committee review.
11/12/2025 Medical Policy Implementation Committee. Coverage eligibility unchanged.
01/01/2026 Coding update
Next Scheduled Review Date: 11/2026

Coding

The five character codes included in the Louisiana Blue Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)†, copyright 2024 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Louisiana Blue Medical Policy Coverage Guidelines is with Louisiana Blue and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Louisiana Blue Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Louisiana Blue Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	0274T Delete code effective 01/01/2026: 0275T Add codes effective 01/01/2026: 62330, 62331
HCPCS	G0276
ICD-10 Diagnosis	All Related Diagnoses

Image-Guided Minimally Invasive Decompression for Spinal Stenosis

Policy # 00278

Original Effective Date: 11/16/2010

Current Effective Date: 12/01/2025

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
 - 1. Consultation with technology evaluation center(s);
 - 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
 - 3. Reference to federal regulations.

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: If the Patient's health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

NOTICE: Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage.