

Lung and Lobar Lung Transplant

Policy # 00414

Original Effective Date: 05/21/2014

Current Effective Date: 05/01/2026

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the “Company”), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Investigational or experimental services are not covered. This includes any drug, device, procedure, or service provided under the investigational arm of a clinical trial or clinical study. These services are excluded from coverage under benefits.

When Services Are Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member’s contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of available data, the Company may consider lung transplantation for carefully selected individuals **eligible for coverage**** when all coverage criteria are met:

- Irreversible, progressively disabling, end-stage pulmonary disease including, but not limited to, the following conditions:
 - Bilateral bronchiectasis
 - Alpha-1 antitrypsin deficiency
 - Primary pulmonary hypertension
 - Pulmonary hypertension due to cardiac disease
 - Cystic fibrosis (both lungs to be transplanted)
 - Bronchopulmonary dysplasia
 - Post inflammatory pulmonary fibrosis
 - Idiopathic/ interstitial pulmonary fibrosis
 - Sarcoidosis
 - Scleroderma/ systemic sclerosis
 - Lymphangiomyomatosis
 - Eosinophilic granuloma (Langerhans cell histiocytosis)
 - Bronchiolitis obliterans
 - Chronic obstructive pulmonary disease (COPD), e.g., emphysema
 - Recurrent pulmonary embolism
 - Eisenmenger’s syndrome
 - Other restrictive lung disease (e.g., allergic alveolitis, asbestosis, collagen vascular disease, desquamative interstitial fibrosis, post-chemotherapy); **AND**
- Unresponsive to maximum medical or surgical therapy; **AND**
- Must meet United Network for Organ Sharing (UNOS) guidelines for a Lung Allocation Score greater than zero; **AND**

Lung and Lobar Lung Transplant

Policy # 00414

Original Effective Date: 05/21/2014

Current Effective Date: 05/01/2026

- None of the contraindications are present (subject to the judgment of the transplant center):
 - Active malignancy, including metastatic cancer
 - Recent malignancy with moderate to high risk of recurrence (assessment of risk of recurrence is documented by the transplant team)
 - Untreated systemic infection making immunosuppression unsafe, including chronic infection (pulmonary or non-pulmonary)
 - Other irreversible end-stage diseases not attributed to lung disease
 - Systemic disease that could be exacerbated by immunosuppression
 - Psychosocial conditions or chemical dependency affecting ability to adhere to therapy
 - Coronary artery disease not amenable to percutaneous intervention or bypass grafting, or associated with significant impairment of left ventricular function
 - Colonization with highly resistant or highly virulent bacteria, fungi, or mycobacteria
 - Acquired immune deficiency syndrome (AIDS) (diagnosis based on Centers for Disease Control and Prevention [CDC] definition of CD4 count, 200 cells/mm³), unless the following are documented:
 - CD4 count greater than 200 cells/mm³ for greater than 6 months;
 - HIV-1 RNA (viral load) undetectable for at least 6 months;
 - On stable anti-retroviral therapy greater than 3 months;
 - No other complications from AIDS (e.g., opportunistic infection, including aspergillus, tuberculosis, coccidioidomycosis, resistant fungal infections, Kaposi's sarcoma or other neoplasm); AND
 - All other criteria for transplantation are met.

Note:

Bilateral lung transplantation is typically required when chronic lung infection and disease is present (i.e., associated with cystic fibrosis and bronchiectasis). Some, but not all, cases of pulmonary hypertension will require bilateral lung transplantation.

Based on review of available data, the Company may consider a lobar lung transplant from a living or deceased donor for carefully selected individuals with end-stage pulmonary disease who meet criteria for lung transplantation to be **eligible for coverage.****

Based on review of available data, the Company may consider lung or lobar lung retransplantation after a failed lung or lobar lung transplant in individuals who meet criteria for lung transplantation to be **eligible for coverage.****

Note:

Bronchiolitis obliterans is associated with chronic lung transplant rejection and thus may be the etiology of a request for lung retransplantation.

Lung and Lobar Lung Transplant

Policy # 00414

Original Effective Date: 05/21/2014

Current Effective Date: 05/01/2026

When Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers lung or lobar lung transplantation in all other situations to be **investigational**.*

Policy Guidelines

Contraindications

The factors below are potential contraindications subject to the judgment of the transplant center:

- Known current malignancy, including metastatic cancer
- Recent malignancy with high risk of recurrence
- Untreated systemic infection making immunosuppression unsafe, including chronic infection
- Other irreversible end-stage diseases not attributed to lung disease
- History of cancer with a moderate risk of recurrence
- Systemic disease that could be exacerbated by immunosuppression
- Psychosocial conditions or chemical dependency affecting ability to adhere to therapy

Policy specific:

- Coronary artery disease not amenable to percutaneous intervention or bypass grafting, or associated with significant impairment of left ventricular function^a; or
- Colonization with highly resistant or highly virulent bacteria, fungi, or mycobacteria.

^a Some individuals may be candidates for combined heart and lung transplantation.

Individuals must meet United Network for Organ Sharing guidelines for a Lung Allocation Score greater than zero.

Lung-Specific Guidelines

Bilateral lung transplantation is typically required when chronic lung infection and disease is present (ie, associated with cystic fibrosis and bronchiectasis). Some, but not all, cases of pulmonary hypertension will require bilateral lung transplantation.

Bronchiolitis obliterans is associated with chronic lung transplant rejection, and thus may be the etiology of a request for lung retransplantation.

Background/Overview

Solid organ transplantation offers a treatment option for patients with different types of endstage organ failure that can be lifesaving or provide significant improvements to a patient's quality of life. Many advances have been made in the last several decades to reduce perioperative complications. Available data supports improvement in long-term survival as well as improved

Lung and Lobar Lung Transplant

Policy # 00414

Original Effective Date: 05/21/2014

Current Effective Date: 05/01/2026

quality of life particularly for liver, kidney, pancreas, heart, and lung transplants. Allograft rejection remains a key early and late complication risk for any organ transplantation. Transplant recipients require life-long immunosuppression to prevent rejection. Patients are prioritized for transplant by mortality risk and severity of illness criteria developed by the Organ Procurement and Transplantation Network (OPTN) and United Network of Organ Sharing.

Lung Transplant

In 2024, 48,136 transplants were performed in the United States procured from more than 41,000 deceased donors and 6900 living donors. Lung transplants were the fourth most common procedure with 3,339 transplants performed from both deceased and living donors in 2024.

End-stage lung disease may derive from different etiologies. The most common indications for lung transplantation are chronic obstructive pulmonary disease, idiopathic pulmonary fibrosis, cystic fibrosis, α_1 -antitrypsin deficiency, and idiopathic pulmonary arterial hypertension. Before consideration for transplant, patients should be receiving maximal medical therapy, including oxygen supplementation, or surgical options, such as lung volume reduction surgery for chronic obstructive pulmonary disease. Lung or lobar lung transplantation is an option for patients with end-stage lung disease despite these measures.

A lung transplant refers to single-lung or double-lung replacement. In a single-lung transplant, only 1 lung from a deceased donor is provided to the recipient. In a double-lung transplant, both the recipient's lungs are removed and replaced by the donor's lungs. In a lobar transplant, a lobe of the donor's lung is excised, sized appropriately for the recipient's thoracic dimensions, and transplanted. Donors for lobar transplant have primarily been living-related donors, with 1 lobe obtained from each of 2 donors (generally friends or family members) in cases for which bilateral transplantation is required. There are also cases of cadaver lobe transplants.

Potential recipients who are 12 years of age and older are ranked according to the Lung Allocation Score. A score may range between 0 and 100 and incorporates predicted survival after transplantation and predicted survival on the waiting list; the Lung Allocation Score takes into consideration the patient's disease and clinical parameters. The waiting list incorporates the Lung Allocation Score, geography, and blood type classifications. Children younger than 12 years old receive priority for lung allocation. Under this system, children younger than 12 years old with respiratory lung failure and/or pulmonary hypertension who meet criteria are considered "priority 1", and all other candidates in the age group are considered "priority 2". A lung review board has the authority to adjust scores on appeal for adults and children.

Potential Contraindications to Transplantation

Malignancy

Malignancies are common after lung transplantation, with 21% and 40% of patients reporting 1 or more malignancies at 5 and 10 years posttransplantation, respectively. Skin cancer occurred most frequently, and lymphoproliferative disorders were the malignancies most associated with morbidity posttransplantation.

Lung and Lobar Lung Transplant

Policy # 00414

Original Effective Date: 05/21/2014

Current Effective Date: 05/01/2026

Human Immunodeficiency Virus Infection

Current OPTN policy permits human immunodeficiency virus (HIV)-positive transplant candidates. The 2020 US Public Health Service guideline also allows for transplantations in HIV-positive recipients with proper screenings and effective regimens for HIV infections; it recommended that all transplant candidates receive HIV, hepatitis b virus (HBV), and hepatitis C virus (HCV) testing during hospital admission for transplant surgery. In 2022, the US Public Health Service published updated guidance for testing transplant candidates aged less than 12 years of age. They recommended that children less than 12 years of age who have received postnatal infectious disease testing are exempt from repeat pretransplant HIV, HBV, and HCV testing during hospital admission for transplant surgery.

The British HIV Association and the British Transplantation Society (2017) updated their guidelines on kidney transplantation in patients with HIV disease. These criteria for adding a patient to the waitlist may be extrapolated to other organs:

- Adherent with treatment, particularly antiretroviral therapy;
- Cluster of Differentiation 4 count greater than 100 cells/mL (ideally >200 cells/mL) for at least 3 months;
- Undetectable HIV viremia (<50 HIV-1 RNA copies/mL) for at least 6 months;
- No opportunistic infections for at least 6 months;
- No history of progressive multifocal leukoencephalopathy, chronic intestinal cryptosporidiosis, or lymphoma.

Other Infections

Infection with *Burkholderia cenocepacia* is associated with increased mortality in some transplant centers, a factor that may be considered when evaluating the overall risk of transplant survival. Two articles have evaluated the impact of infection with various species of *Burkholderia* on outcomes for lung transplantation for cystic fibrosis. In a study by Murray et al (2008), multivariate Cox survival models were applied to 1026 lung transplant candidates and 528 transplant recipients. Of the transplant recipients, 88 were infected with *Burkholderia*. Among transplant recipients infected with *B. cenocepacia*, only those infected with nonepidemic strains (n=11) had significantly greater posttransplant mortality than uninfected patients (hazard ratio [HR], 2.52; 95% confidence interval [CI], 1.04 to 6.12; p=.04). Transplant recipients infected with *Burkholderia gladioli* (n=14) also had significantly greater post transplant mortality than uninfected patients (HR, 2.23; 95% CI, 1.05 to 4.74; p=.04). When adjustments for specific species or strains were included, the Lung Allocation Scores of *Burkholderia multivorans*-infected transplant candidates were comparable with uninfected candidate scores, and scores for patients infected with nonepidemic *B. cenocepacia* or *B. gladioli* were lower. In a smaller study of 22 patients colonized with *Burkholderia cepacia* complex who underwent lung transplantation in 2 French centers, Boussaud et al (2008) reported that the risk of death by univariate analysis was significantly higher for the 8 patients infected with *B. cenocepacia* than for the other 14 colonized patients (11 of whom had *B. multivorans*).

An analysis of international registry data by Yusen et al (2016) found that non-cytomegalovirus (CMV) infection is a major cause of mortality within 30 days of a lung transplant in adults. A total

Lung and Lobar Lung Transplant

Policy # 00414

Original Effective Date: 05/21/2014

Current Effective Date: 05/01/2026

of 655 (19%) of 3424 deaths after transplants between 1990 and 2015 were due to non-CMV infection. Only 3 (0.1%) of the deaths were due to CMV infection.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

Solid organ transplants are a surgical procedure and, as such, are not subject to regulation by the U.S. Food and Drug Administration (FDA).

The FDA regulates human cells and tissues intended for implantation, transplantation, or infusion through the Center for Biologics Evaluation and Research, under Code of Federal Regulation Title 21, parts 1270 and 1271. Solid organs used for transplantation are subject to these regulations.

Rationale/Source

This medical policy was developed through consideration of peer-reviewed medical literature generally recognized by the relevant medical community, U.S. Food and Drug Administration approval status, nationally accepted standards of medical practice and accepted standards of medical practice in this community, technology evaluation centers, reference to regulations, other plan medical policies, and accredited national guidelines.

A lung transplant consists of replacing all or part of diseased lungs with healthy lung(s) or lobes. Transplantation is an option for individuals with end-stage lung disease.

Summary of Evidence

For individuals who have end-stage pulmonary disease who receive a lung transplant, the evidence includes case series and registry studies. Relevant outcomes are overall survival (OS), change in disease status, and treatment-related mortality and morbidity. International registry data on a large number of patients receiving lung transplantation (>50,000) found relatively high patient survival rates, especially among those who survived the first year posttransplant. After adjusting for potential confounding factors, survival did not differ significantly after single- or double-lung transplant. Lung transplantation may be the only option for some patients with end-stage lung disease. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have end-stage pulmonary disease who receive a lobar lung transplant, the evidence includes case series and systematic reviews. Relevant outcomes are OS, change in disease status, and treatment-related mortality and morbidity. There are less data on lung lobar transplants than on whole-lung transplants, but several case series have reported reasonably similar survival outcomes between the procedures, and lung lobar transplants may be the only option for patients unable to wait for a whole-lung transplant. A 2017 systematic review found 1-year survival rates in available published studies ranging from 50% to 100%. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Lung and Lobar Lung Transplant

Policy # 00414

Original Effective Date: 05/21/2014

Current Effective Date: 05/01/2026

For individuals with a prior lung or lobar transplant who meet criteria for a lung transplant and receive a lung or lobar lung retransplant, the evidence includes case series and registry studies. Relevant outcomes are OS, change in disease status, and treatment-related mortality and morbidity. Data from registries and case series have found favorable outcomes with lung retransplantation in patients who meet criteria for initial lung transplantation. Given the exceedingly poor survival prognosis without retransplantation of patients who have exhausted other treatments, the evidence of a moderate level of posttransplant survival may be considered sufficient in this patient population. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Supplemental Information

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American Thoracic Society et al

Evidence-based recommendations from the American Thoracic Society and 3 international cardiac societies were published in 2011 for the diagnosis and management of patients with idiopathic fibrosis. For appropriately selected patients with idiopathic pulmonary fibrosis, the international guideline panel recommended lung transplantation (strong recommendation, low-quality evidence). An update to this document was published in 2015 in which the committee did not make a recommendation regarding single versus bilateral lung transplantation in patients with idiopathic fibrosis. The committee stated that "it is unclear whether single or bilateral lung transplantation is preferential for long-term outcomes."

In 2022, the American Thoracic Society along with the 3 other international cardiac societies published updated guidance on diagnosis and management of idiopathic pulmonary fibrosis and progressive pulmonary fibrosis. In terms of treatment considerations, the committee stated that "patients at increased risk of mortality should be referred for lung transplantation at diagnosis."

In 2014, the American Thoracic Society published guidelines on the management of bronchiolitis obliterans syndrome in lung transplant recipients in conjunction with the International Society for Heart and Lung Transplantation and the European Respiratory Society. The guideline recommends referral to a transplant surgeon to be evaluated for retransplantation for end-stage bronchial obliterans syndrome that is refractory to other therapies.

In 2025, the American Thoracic Society published guidelines on interventional strategies for children with progressive pulmonary hypertension. Recommendations made regarding lung transplantation are below:

Lung and Lobar Lung Transplant

Policy # 00414

Original Effective Date: 05/21/2014

Current Effective Date: 05/01/2026

- "We suggest lung transplantation in children with progressive PH [pulmonary hypertension] and RV [right ventricular] failure despite optimal therapy (conditional recommendation, very low certainty of evidence)."
- "We suggest lung transplantation in children with progressive PH and RV failure despite optimal therapy who are on ECMO [extracorporeal membrane oxygenation] support without reversible cause (conditional recommendation, very low certainty of evidence)."

International Society for Heart and Lung Transplantation

Initial Transplant

In 2021, the International Society for Heart and Lung Transplantation published updated consensus-based guidelines on the selection of lung transplant candidates. The guidelines states that:

- "Lung transplantation should be considered for adults with chronic, end-stage lung disease who meet all the following general criteria:
 1. High (>50%) risk of death from lung disease within 2 years if lung transplantation is not performed.
 2. High (>80%) likelihood of 5-year post-transplant survival from a general medical perspective provided that there is adequate graft function."

The guideline also notes risk factors to be considered in the evaluation of transplant candidates, along with pediatric and disease-specific considerations.

Retransplant

The 2021 guideline update briefly addressed lung retransplantation, with the consensus statement noting that "The outcomes after re-transplants are inferior compared to first lung transplants, particularly if the re-transplant is done within the first year after the original transplant or for patients with restrictive allograft syndrome (RAS) [...] In the pre-transplant evaluation of such patients, particular emphasis should be focused on understanding the possible reasons for the graft failure, such as alloimmunization, poor adherence, gastroesophageal reflux, or repeated infections."

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

Lung transplantation is covered under Medicare when performed in a facility approved by Medicare as meeting institutional coverage criteria. The Centers for Medicare & Medicaid Services have stated that, under certain limited cases, exceptions to the facility-related criteria may be warranted if there is justification and the facility ensures safety and efficacy objectives.

Ongoing and Unpublished Clinical Trials

Some currently ongoing and unpublished trials that might influence this review are listed in Table 1.

Lung and Lobar Lung Transplant

Policy # 00414

Original Effective Date: 05/21/2014

Current Effective Date: 05/01/2026

Table 1. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT00177918	Prospective Evaluations of Infectious Complication in Lung Transplant Recipients	600	Dec 2025

NCT: national clinical trial.

References

1. Black CK, Termanini KM, Aguirre O, et al. Solid organ transplantation in the 21 st century. *Ann Transl Med.* Oct 2018; 6(20): 409. PMID 30498736
2. Transplant trends. United Network for Organ Sharing website. Updated July 28, 2025. <https://unos.org/data/transplant-trends/>.
3. Organ Procurement and Transplantation Network (OPTN). Policy 10: Allocation of Lungs. Updated June 26, 2025; https://optn.transplant.hrsa.gov/media/1200/optn_policies.pdf.
4. Yusen RD, Christie JD, Edwards LB, et al. The Registry of the International Society for Heart and Lung Transplantation: Thirtieth Adult Lung and Heart-Lung Transplant Report--2013; focus theme: age. *J Heart Lung Transplant.* Oct 2013; 32(10): 965-78. PMID 24054805
5. Jones JM, Kracalik I, Levi ME, et al. Assessing Solid Organ Donors and Monitoring Transplant Recipients for Human Immunodeficiency Virus, Hepatitis B Virus, and Hepatitis C Virus Infection - U.S. Public Health Service Guideline, 2020. *MMWR Recomm Rep.* Jun 26 2020; 69(4): 1-16. PMID 32584804
6. Free RJ, Levi ME, Bowman JS, et al. Updated U.S. Public Health Service Guideline for testing of transplant candidates aged 12 years for infection with HIV, hepatitis B virus, and hepatitis C virus - United States, 2022. *Am J Transplant.* Sep 2022; 22(9): 2269-2272. PMID 36039545
7. Working Party of the British Transplantation Society. Kidney and Pancreas Transplantation in Patients with HIV. Second Edition (Revised). British Transplantation Society Guidelines. Macclesfield, UK: British Transplantation Society; 2017.
8. Alexander BD, Petzold EW, Reller LB, et al. Survival after lung transplantation of cystic fibrosis patients infected with *Burkholderia cepacia* complex. *Am J Transplant.* May 2008; 8(5): 1025-30. PMID 18318775
9. Murray S, Charbeneau J, Marshall BC, et al. Impact of *burkholderia* infection on lung transplantation in cystic fibrosis. *Am J Respir Crit Care Med.* Aug 15 2008; 178(4): 363-71. PMID 18535253
10. Boussaud V, Guillemain R, Grenet D, et al. Clinical outcome following lung transplantation in patients with cystic fibrosis colonised with *Burkholderia cepacia* complex: results from two French centres. *Thorax.* Aug 2008; 63(8): 732-7. PMID 18408050
11. Yusen RD, Edwards LB, Dipchand AI, et al. The Registry of the International Society for Heart and Lung Transplantation: Thirty-third Adult Lung and Heart-Lung Transplant Report-2016; Focus Theme: Primary Diagnostic Indications for Transplant. *J Heart Lung Transplant.* Oct 2016; 35(10): 1170-1184. PMID 27772669

Lung and Lobar Lung Transplant

Policy # 00414

Original Effective Date: 05/21/2014

Current Effective Date: 05/01/2026

12. Paraskeva MA, Edwards LB, Levvey B, et al. Outcomes of adolescent recipients after lung transplantation: An analysis of the International Society for Heart and Lung Transplantation Registry. *J Heart Lung Transplant*. Mar 2018; 37(3): 323-331. PMID 28320631
13. Goldfarb SB, Levvey BJ, Edwards LB, et al. The Registry of the International Society for Heart and Lung Transplantation: Nineteenth Pediatric Lung and Heart-Lung Transplantation Report-2016; Focus Theme: Primary Diagnostic Indications for Transplant. *J Heart Lung Transplant*. Oct 2016; 35(10): 1196-1205. PMID 27772671
14. Thabut G, Christie JD, Kremers WK, et al. Survival differences following lung transplantation among US transplant centers. *JAMA*. Jul 07 2010; 304(1): 53-60. PMID 20606149
15. Black MC, Trivedi J, Schumer EM, et al. Double lung transplants have significantly improved survival compared with single lung transplants in high lung allocation score patients. *Ann Thorac Surg*. Nov 2014; 98(5): 1737-41. PMID 25110334
16. Yu H, Bian T, Yu Z, et al. Bilateral Lung Transplantation Provides Better Long-term Survival and Pulmonary Function Than Single Lung Transplantation: A Systematic Review and Meta-analysis. *Transplantation*. Dec 2019; 103(12): 2634-2644. PMID 31283687
17. Yusef RD, Shearon TH, Qian Y, et al. Lung transplantation in the United States, 1999-2008. *Am J Transplant*. Apr 2010; 10(4 Pt 2): 1047-68. PMID 20420652
18. Shafii AE, Mason DP, Brown CR, et al. Too high for transplantation? Single-center analysis of the lung allocation score. *Ann Thorac Surg*. Nov 2014; 98(5): 1730-6. PMID 25218678
19. Date H. Update on living-donor lobar lung transplantation. *Curr Opin Organ Transplant*. Oct 2011; 16(5): 453-7. PMID 21836512
20. Eberlein M, Reed RM, Chahla M, et al. Lobar lung transplantation from deceased donors: A systematic review. *World J Transplant*. Feb 24 2017; 7(1): 70-80. PMID 28280698
21. Barr ML, Schenkel FA, Bowdish ME, et al. Living donor lobar lung transplantation: current status and future directions. *Transplant Proc*. Nov 2005; 37(9): 3983-6. PMID 16386604
22. Date H, Sato M, Aoyama A, et al. Living-donor lobar lung transplantation provides similar survival to cadaveric lung transplantation even for very ill patients†. *Eur J Cardiothorac Surg*. Jun 2015; 47(6): 967-72; discussion 972-3. PMID 25228745
23. Date H, Shiraishi T, Sugimoto S, et al. Outcome of living-donor lobar lung transplantation using a single donor. *J Thorac Cardiovasc Surg*. Sep 2012; 144(3): 710-5. PMID 22717276
24. Slama A, Ghanim B, Klikovits T, et al. Lobar lung transplantation--is it comparable with standard lung transplantation?. *Transpl Int*. Sep 2014; 27(9): 909-16. PMID 24810771
25. Kilic A, Beaty CA, Merlo CA, et al. Functional status is highly predictive of outcomes after redo lung transplantation: an analysis of 390 cases in the modern era. *Ann Thorac Surg*. Nov 2013; 96(5): 1804-11; discussion 1811. PMID 23968759
26. Kawut SM. Lung retransplantation. *Clin Chest Med*. Jun 2011; 32(2): 367-77. PMID 21511096
27. Biswas Roy S, Panchanathan R, Walia R, et al. Lung Retransplantation for Chronic Rejection: A Single-Center Experience. *Ann Thorac Surg*. Jan 2018; 105(1): 221-227. PMID 29100649
28. Organ Procurement and Transplantation Network (OPTN). National Data. n.d.; <https://optn.transplant.hrsa.gov/data/view-data-reports/national-data/>.
29. Raghu G, Collard HR, Egan JJ, et al. An official ATS/ERS/JRS/ALAT statement: idiopathic pulmonary fibrosis: evidence-based guidelines for diagnosis and management. *Am J Respir Crit Care Med*. Mar 15 2011; 183(6): 788-824. PMID 21471066

Lung and Lobar Lung Transplant

Policy # 00414

Original Effective Date: 05/21/2014

Current Effective Date: 05/01/2026

30. Raghu G, Rochwerg B, Zhang Y, et al. An Official ATS/ERS/JRS/ALAT Clinical Practice Guideline: Treatment of Idiopathic Pulmonary Fibrosis. An Update of the 2011 Clinical Practice Guideline. *Am J Respir Crit Care Med*. Jul 15 2015; 192(2): e3-19. PMID 26177183
31. Raghu G, Remy-Jardin M, Richeldi L, et al. Idiopathic Pulmonary Fibrosis (an Update) and Progressive Pulmonary Fibrosis in Adults: An Official ATS/ERS/JRS/ALAT Clinical Practice Guideline. *Am J Respir Crit Care Med*. May 01 2022; 205(9): e18-e47. PMID 35486072
32. Meyer KC, Raghu G, Verleden GM, et al. An international ISHLT/ATS/ERS clinical practice guideline: diagnosis and management of bronchiolitis obliterans syndrome. *Eur Respir J*. Dec 2014; 44(6): 1479-503. PMID 25359357
33. Hayes D, Jennerich AL, Coleman RD, et al. Interventional Strategies for Children with Progressive Pulmonary Hypertension Despite Optimal Therapy: An Official American Thoracic Society Clinical Practice Guideline. *Am J Respir Crit Care Med*. Feb 2025; 211(2): 157-173. PMID 39531626
34. Leard LE, Holm AM, Valapour M, et al. Consensus document for the selection of lung transplant candidates: An update from the International Society for Heart and Lung Transplantation. *J Heart Lung Transplant*. Nov 2021; 40(11): 1349-1379. PMID 34419372
35. Centers for Medicare & Medicaid. Transplant. Updated June 9, 2025; <https://www.cms.gov/Medicare/Provider-Enrollment-and-Certification/CertificationandCompliance/Transplant.html>.

Policy History

Original Effective Date: 05/21/2014

Current Effective Date: 05/01/2026

05/01/2014	Medical Policy Committee review
05/21/2014	Medical Policy Implementation Committee approval. New policy.
08/06/2015	Medical Policy Committee review
08/19/2015	Medical Policy Implementation Committee approval. No change to coverage.
08/04/2016	Medical Policy Committee review
08/17/2016	Medical Policy Implementation Committee approval. No change to coverage.
01/01/2017	Coding update: Removing ICD-9 Diagnosis Codes
08/03/2017	Medical Policy Committee review
08/23/2017	Medical Policy Implementation Committee approval. No change to coverage.
10/04/2018	Medical Policy Committee review
10/17/2018	Medical Policy Implementation Committee approval. No change to coverage. Policy reformatted.
10/03/2019	Medical Policy Committee review
10/09/2019	Medical Policy Implementation Committee approval. No change to coverage.
10/01/2020	Medical Policy Committee review
10/07/2020	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
11/04/2021	Medical Policy Committee review

Lung and Lobar Lung Transplant

Policy # 00414

Original Effective Date: 05/21/2014

Current Effective Date: 05/01/2026

11/10/2021	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
11/03/2022	Medical Policy Committee review
11/09/2022	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
11/02/2023	Medical Policy Committee review
11/08/2023	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
11/07/2024	Medical Policy Committee review
11/13/2024	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
11/06/2025	Medical Policy Committee review
11/12/2025	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
04/02/2026	Medical Policy Committee review
04/10/2026	Medical Policy Implementation Committee approval. Added examples of conditions eligible for lung transplant- associated with irreversible, progressively disabling, end-stage- pulmonary disease (list is not exhaustive). Added criteria for human immunodeficiency virus (HIV) positive recipients.

Next Scheduled Review Date: 04/2027

Coding

The five character codes included in the Louisiana Blue Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)[‡], copyright 2025 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Louisiana Blue Medical Policy Coverage Guidelines is with Louisiana Blue and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Louisiana Blue Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Louisiana Blue Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Lung and Lobar Lung Transplant

Policy # 00414

Original Effective Date: 05/21/2014

Current Effective Date: 05/01/2026

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	32850, 32851, 32852, 32853, 32854, 32855, 32856
HCPCS	S2060, S2061
ICD-10 Diagnosis	All related Diagnoses

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
 1. Consultation with technology evaluation center(s);
 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
 3. Reference to federal regulations.

**Medically Necessary (or “Medical Necessity”) - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

For these purposes, “nationally accepted standards of medical practice” means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

Lung and Lobar Lung Transplant

Policy # 00414

Original Effective Date: 05/21/2014

Current Effective Date: 05/01/2026

NOTICE: If the Patient's health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

NOTICE: Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage.

NOTICE: If an authorization for an ongoing course of treatment has been provided to a member and the member changes from one health plan to another health plan (e.g., a member moves from carrier A to Louisiana Blue), Louisiana Blue may honor the previous health plan's authorization for the same service under the same type of in-network benefit for a 90-day transition period. Documentation of the authorization for the ongoing course of treatment from the previous health plan must be provided to us by the member or their provider and the services provided for the course of treatment must otherwise be a covered service under the Louisiana Blue health plan. This provision does not apply to medications covered under the plan's pharmacy benefit.