

Repository Corticotropin Injection

Policy # 00230

Original Effective Date: 07/16/2008

Current Effective Date: 01/01/2026

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Note: Vigabatrin Products are addressed separately in medical policy 00244.

When Services May Be Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of available data, the Company may consider repository corticotropin injection (Acthar[®] Gel)[†] for the treatment of infantile spasms (West syndrome) to be **eligible for coverage**.**

Patient Selection Criteria

Coverage eligibility for the use of repository corticotropin injection (Acthar Gel) will be considered when the following criteria are met:

- Patient has a diagnosis of infantile spasms AND is less than 2 years of age.

When Services Are Not Covered

Based on review of available data, the Company considers the use of repository corticotropin injection (Acthar Gel) EXCEPT for treating infantile spasms to be **not covered**.**

Based on review of available data, the Company considers the use of repository corticotropin injection (Cortrophin[™] Gel)[‡] to be **not covered**.**

Note: Repository corticotropin injection (Acthar Gel) for all conditions EXCEPT infantile spasms is considered an exclusion in most member contracts. Repository corticotropin injection (Cortrophin Gel) for all FDA-approved indications is considered an exclusion in most member contracts.

Background/Overview

Repository corticotropin injection (Acthar Gel, Cortrophin Gel) is a purified, sterile preparation of the natural form of adrenocorticotrophic hormone (ACTH) in gelatin to provide a prolonged release after intramuscular or subcutaneous injection. ACTH works by stimulating the adrenal cortex to produce cortisol, corticosterone, and a number of other hormones.

Repository Corticotropin Injection

Policy # 00230

Original Effective Date: 07/16/2008

Current Effective Date: 01/01/2026

Acthar Gel is an ACTH analogue indicated as monotherapy for the treatment of infantile spasms in infants and children less than 2 years of age. According to the package insert, Acthar gel may also be used for the following disorders and diseases: rheumatic, collagen, dermatologic, allergic states, ophthalmic, respiratory and edematous states. However, the drug was approved by the Food and Drug Administration (FDA) in 1952 prior to the requirement that companies provide evidence of clinical efficacy.

Cortrophin gel is an ACTH analogue indicated for specific rheumatic disorders, collagen diseases, dermatologic diseases, allergic states, ophthalmic diseases, respiratory diseases, edematous states, and multiple sclerosis exacerbations. This product was also initially approved by the FDA before the requirement that companies provide evidence of clinical efficacy. Unlike Acthar Gel, Cortrophin Gel is not approved for the treatment of infantile spasms.

Contraindications for use of these agents include scleroderma, osteoporosis, systemic fungal infections, ocular herpes simplex, recent surgery, history of or the presence of a peptic ulcer, congestive heart failure, uncontrolled hypertension, or sensitivity to proteins of porcine origin.

West Syndrome/Infantile Spasms

West syndrome is a rare epileptic disorder of early infancy (90% of cases are diagnosed the first year of life) consisting of three main characteristics: infantile spasm, mental retardation, and hypsarrhythmia, a specific abnormal pattern on EEG. Often the term infantile spasms is used synonymously with West syndrome. Infantile spasms are characterized by an initial contraction phase followed by a more sustained tonic phase.

Another treatment option for infantile spasms is vigabatrin (Sabril®, Vigafyde™, generics)[‡] which is available in various oral dosage forms and brand and generic formulations. Vigabatrin is indicated as monotherapy for pediatric patients with infantile spasms for whom the potential benefits outweigh the potential risk of vision loss.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

Acthar is currently approved in the U.S. for the treatment of infantile spasms, multiple sclerosis exacerbations, rheumatic disorders, collagen diseases, dermatologic diseases, allergic states, ophthalmic diseases, symptomatic sarcoidosis, and edematous states.

Cortrophin Gel is currently approved for the following disorders:

- Rheumatic disorders: As adjunctive therapy for short-term administration (to tide the patient over an acute episode or exacerbation) in
 - Psoriatic arthritis
 - Rheumatoid arthritis, including juvenile rheumatoid arthritis (selected cases may require low-dose maintenance therapy)
 - Ankylosing spondylitis
 - Acute gouty arthritis

Repository Corticotropin Injection

Policy # 00230

Original Effective Date: 07/16/2008

Current Effective Date: 01/01/2026

- Collagen diseases: during an exacerbation or as maintenance therapy in selected cases of
 - Systemic lupus erythematosus
 - Systemic dermatomyositis (polymyositis)
- Dermatologic diseases:
 - Severe erythema multiforme (Stevens-Johnson syndrome)
 - Severe psoriasis
- Allergic states:
 - Atopic dermatitis
 - Serum sickness
- Ophthalmic diseases: severe acute and chronic allergic and inflammatory processes involving the eye and its adnexa such as:
 - Allergic conjunctivitis
 - Keratitis
 - Iritis and iridocyclitis
 - Diffuse posterior uveitis and choroiditis
 - Optic neuritis
 - Chorioretinitis
 - Anterior segment inflammation
- Respiratory diseases:
 - Symptomatic sarcoidosis
- Nervous system:
 - Acute exacerbations of multiple sclerosis

Rationale/Source

This medical policy was developed through consideration of peer-reviewed medical literature generally recognized by the relevant medical community, U.S. Food and Drug Administration approval status, nationally accepted standards of medical practice and accepted standards of medical practice in this community, technology evaluation centers, reference to regulations, other plan medical policies, and accredited national guidelines.

Infantile spasms

The effectiveness of Acthar Gel as a treatment for infantile spasms was demonstrated in a single blinded (video EEG Interpreter-blinded) clinical trial in which patients were randomized to receive either a 2 week course of treatment with Acthar Gel (75 U/m²) intramuscular twice daily or prednisone 1 mg/kg by mouth twice daily. The primary outcome was a comparison of the number of patients in each group who were treatment responders, defined as a patient having complete suppression of both clinical spasms and hypsarrhythmia on a full sleep cycle video EEG performed 2 weeks following treatment initiation, rated by an investigator blinded to treatment. Thirteen of 15 patients (86.7%) responded to Acthar Gel as compared to 4 of 14 patients (28.6%) given prednisone ($p < 0.002$). The 2-week treatment was followed by a 2-week period of taper. Nonresponders to the prednisone treatment were eligible to receive Acthar Gel treatment. Seven of 8 patients (87.5%)

Repository Corticotropin Injection

Policy # 00230

Original Effective Date: 07/16/2008

Current Effective Date: 01/01/2026

responded to Acthar Gel after not responding to prednisone. Similarly, the 2 nonresponder patients from the Acthar Gel treatment were eligible to receive treatment with prednisone. One of the 2 patients (50%) responded to the prednisone treatment after not responding to Acthar Gel.

A supportive single-blind, randomized clinical trial comparing high-dose, long-duration treatment (150U/m² once daily for 3 weeks, n = 30) of Acthar Gel with low-dose, short-duration treatment (20U once daily for 2 weeks, n = 29) for the treatment of infantile spasms was also evaluated in infants and children less than 2 years of age. Nonresponders (defined as in the previously described study) in the low-dose group received a dose escalation at 2 weeks to 30U once daily. Nominal statistical superiority of the high dose treatment, as compared to the low dose treatment, was observed for cessation of spasms but not for the resolution of hypsarrhythmia.

In 2012 the American Academy of Neurology (AAN) and the Child Neurology Society updated the evidence-based guideline for the medical treatment of infantile spasms. The guidelines note that ACTH is a first-line agent for the short-term treatment of infantile spasms. The Infantile Spasms Working Group (ISWG) published a US consensus report on infantile spasms in 2010. Data regarding ACTH use in infantile spasms were detailed and it was determined that ACTH is an effective first-line therapy for infantile spasms.

Acute Exacerbations of Multiple Sclerosis

Although Acthar is indicated for the treatment of exacerbations of MS in adults, more recent studies evaluating multiple sclerosis have demonstrated that intravenous corticosteroids are at least as effective, or more effective than repository corticotropin. Acthar has been studied in patients with acute exacerbations of MS and short-term use, usually given IM or SC for 14 or fewer days, led to benefits in signs and symptoms of MS. A double-blind, randomized controlled trial found that ACTH given IM over 14 days had similar efficacy in acute exacerbations of MS as methylprednisolone given as 1 gram IV daily for 3 days.

Other potential uses of repository corticotrophin injection

There are a limited number of small case series reporting on the use of repository corticotropin injection for other corticosteroid-responsive conditions. For example, in 2011, Bomback et al published a retrospective case series in 21 patients with idiopathic, nondiabetic nephrotic syndrome who were treated with repository corticotropin. Repository corticotropin was used as a primary therapy in 3 patients; the other 18 patients had failed a mean of 2.3 immunosuppressive regimens before using repository corticotropin. Four (19%) of the 21 patients were in complete remission, defined as stable or improved renal function with final proteinuria falling to less than 500 mg/d. An additional 7 (33%) of 21 patients had a partial remission (at least a 50% reduction in proteinuria and final proteinuria 500-3500 mg/d). Regarding these other uses of repository corticotropin injection, data and guidelines do not suggest a substantial role in therapy. Further data are needed before use in other areas can be recommended.

Policy # 00230

Original Effective Date: 07/16/2008

Current Effective Date: 01/01/2026

References

1. Mackay MT, Weiss SK, Adams-Webber T et al. Practice Parameter: Medical treatment of Infantile Spasms: Report of the American Academy of Neurology and the Child Neurology Society. *Neurology* 2004; 62(10):1668-81. (Also available online at <http://www.neurology.org/cgi/content/full/62/10/1668>).
2. McElhaney JL. Repository corticotropin injection as an adjunct to smoking cessation during the initial nicotine withdrawal period: results from a family practice clinic. *Clin Ther* 1989; 11(6):846-53.
3. Underwood M. Gout. *Clin Evid (Online)* 2008; 2008, pii: 1120.
4. Janssens HJ, Lucassen PL, Van de Laar FA et al. Systemic corticosteroids for acute gout. *Cochrane Database Syst Rev* 2008; (2):CD005521.
5. Schlesinger N. Overview of the management of acute gout and the role of adrenocorticotrophic hormone. *Drugs* 2008; 68(4):407-15.
6. Gayatri NA, Ferrie CD, Cross H. Corticosteroids including ACTH for childhood epilepsy other than epileptic spasms. *Cochrane Database Syst Rev* 2007; (1):CD005222.
7. Hancock E, Osborne JP, Milner P. Treatment of infantile spasms. *Cochrane Database Syst Rev* 2002; (2):CD001770. Update in: *Cochrane Database Syst Rev* 2008; (4):CD001770.
8. Gettig J, Cummings JP, Matuszewski K. H.P. Acthar gel and cosyntropin review. *P T* 2009; 34(5):250-7. Available at <http://www.ncbi.nlm.nih.gov/>.
9. Bomback AS, Tumlin JA, Baranski J et al. Treatment of nephritic syndrome with adrenocorticotrophic hormone (ACTH) gel. *Drug Design Dev Ther* 2011; 5: 147-153.
10. Go CY, Mackay MT, Weiss SK, et al. Evidence-based guideline update: medical treatment of infantile spasms: Report of the guideline development subcommittee of the American Academy of Neurology and the Practice Committee of the Child Neurology Society. *Neurology*. 2012;78:1974-1980.
11. Pellock JM, Hrachovy R, Shinnar S, et al. Infantile spasms: a US consensus report. *Epilepsia*. 2010;51(10):2175-2189.
12. National Clinical Advisory Board of the National Multiple Sclerosis Society. Expert Opinion Paper. Treatment recommendations for physicians. Recommendations regarding corticosteroids in the management of multiple sclerosis. 2008. Available at <http://www.nationalmssociety.org/ms-clinical-care-network/clinical-resources-and-tools/publications/expert-opinion-papers/index.aspx>.
13. National Multiple Sclerosis Society. HP Acthar Gel (adrenocorticotrophic hormone [ACTH]). Available at <http://www.nationalmssociety.org/about-multiple-sclerosis/what-we-know-about-ms/treatments/medications/acth/index.aspx>.
14. Kidney Disease: Improving Global Outcomes (KDIGO) glomerulonephritis Work Group. KDIGO Clinical Practice Guidelines for Glomerulonephritis. *Kidney Int. Suppl.* 2012;2:139-274. Available at http://www.kdigo.org/clinical_practice_guidelines/pdf/KDIGO-GN-Guideline.pdf.
15. Gipson DS, Massengill SF, Yao L, et al. Management of childhood onset nephrotic syndrome. *Pediatrics*. 2009;124:747-757.
16. Waldman M, Austin HA. Treatment of idiopathic membranous nephropathy. *J Am Soc Nephrol*. 2012;23:1617-1630.

Repository Corticotropin Injection

Policy # 00230

Original Effective Date: 07/16/2008

Current Effective Date: 01/01/2026

17. Baram TZ, Mitchell WG, Tournay A, et al. High-dose corticotrophin (ACTH) versus prednisone for infantile spasms: a prospective, randomized, blinded study. *Pediatrics*. 1996;97(3):375-379.
18. Hrachovy RA, Frost JD, Glaze DG. High-dose, long-duration versus low-dose, short-duration corticotropin therapy for infantile spasms. *J Pediatr*. 1994;124(5 Pt 1):803-806.
19. Rose AS, Kuzma JW, Kurtzke JF, et al. Cooperative study in the evaluation of therapy in multiple sclerosis: ACTH vs. placebo—final report. *Neurology*. 1970;20(5):1-59.
20. Sibley WA. Spotlight series: pivotal trials through today's knowledge – adrenocorticotrophic hormone. *Int MS J*. 2009;16:42-26.
21. Filippini G, Brusaferri F, Sibley WA, et al. Corticosteroids or ACTH for acute exacerbations in multiple sclerosis (Review). *Cochrane Database Sys Rev*. 2000;(4):CD001331.
22. Simsarian J, Saunders C, Smith DM. Five-day regimen of intramuscular or subcutaneous self-administered adrenocorticotropin hormone gel for acute exacerbations of multiple sclerosis: a prospective, randomized, open-label pilot trial. *Drug Design Devel Ther*. 2011;5:381-389.
23. Thompson AJ, Kennard D, Swash M, et al. Relative efficacy of intravenous methylprednisolone and ACTH in the treatment of acute relapse in MS. *Neurology*. 1989;39:969-971.
24. Berkovich R, Fernandez M, Subhani D. Monthly pulse adrenocorticotrophic hormone (ACTH) or methylprednisolone therapy for long-term treatment of multiple sclerosis as an add-on therapy to beta-interferons: interim results from a pilot study [poster P576]. Presented at: European Committee for Treatment and Research in Multiple Sclerosis (ECTRIMS) (with ACTRIMS [Americas Committee for Treatment and Research in Multiple Sclerosis]). Amsterdam, The Netherlands, October 19-22, 2011. Available at <http://www.posters2view.com/ECTRIMS2011/view.php?nu=489>.
25. Tumlin JA, Galphin CM, Rovin BH. Advanced diabetic nephropathy with nephrotic range proteinuria: long-term efficacy of subcutaneous adrenocorticotrophic hormone (ACTH) therapy on proteinuria and urinary vascular endothelial growth factor (VEGF) levels [abstract TH-OR073]. Presented at: the American Society of Nephrology Renal Week; Philadelphia, PA; November 8-13, 2011. Available at http://www.abstracts2view.com/asn_2011/view.php?nu=20482&terms=&type=abstract.
26. Food and Drug Administration. Center for Drug Evaluation and Research. Pre Decisional Agency Memorandum. Subject: Acthar Gel (NDA 022432). September 27, 2010. Application number: 022432Orig1s000. Available at: http://www.accessdata.fda.gov/drugsatfda_docs/nda/2010/022432Orig1s000OtherR.pdf
27. Bomback AS, Tumlin JA, Baranski J, et al. Treatment of nephrotic syndrome with adrenocorticotrophic hormone (ACTH) gel. *Drug Des Devel Ther*. 2011;5:147-153.
28. Bomback AS, Canetta PA, Beck LH, et al. Treatment of resistant glomerular diseases with adrenocorticotrophic hormone gel: a prospective trial. *Am J Nephrol*. 2012;36:58-67.
29. Berg AL, Arnadottir M. ACTH-induced improvement in the nephrotic syndrome in patients with a variety of diagnoses. *Nephrol Dial Transplant*. 2004;19:1305-1307.
30. Ponticelli C, Passerini P, Salvadori M, et al. A randomized pilot trial comparing methylprednisolone plus a cytotoxic agent versus synthetic adrenocorticotrophic hormone in idiopathic membranous nephropathy. *Am J Kidney Dis*. 2006;47(2):223-240.
31. Rauen T, Michaelis A, Floege J, Mertens PR. Case series of idiopathic membranous nephropathy with long-term beneficial effects of ACTH peptide 1-24. *Clin Nephrol*. 2009;71(6):637-642.

Repository Corticotropin Injection

Policy # 00230

Original Effective Date: 07/16/2008

Current Effective Date: 01/01/2026

32. Berg AL, Nilsson-Ehle P, Arnadottir M. Beneficial effects of ACTH on the serum lipoprotein profile and glomerular function in patients with membranous nephropathy. *Kidney Int.* 1999;56(4):1534-1543.
33. Hahn BH, McMahon MA, Wilkinson A, et al. American College of Rheumatology guidelines for screening, treatment, and management of lupus nephritis. *Arthritis Care Res (Hoboken).* 2012;64(6):797-808.
34. Levine T. Treating refractory dermatomyositis or polymyositis with adrenocorticotrophic hormone gel: a retrospective case series. *Drug Des Devel Ther.* 2012;6:133-139.
35. Ernste FC, Reed AM. Idiopathic inflammatory myopathies: current trends in pathogenesis, clinical features, and up-to-date treatment recommendations. *Mayo Clin Proc.* 2013;88(1):83-105.
36. Purified Cortrophin Gel [package insert]. ANI Pharmaceuticals, Inc. Baudette, MN. Updated November 2021.

Policy History

Original Effective Date: 07/16/2008

Current Effective Date: 01/01/2026

07/02/2008	Medical Director review
07/16/2008	Medical Policy Committee approval. New policy.
07/02/2009	Medical Director review
07/22/2009	Medical Policy Committee approval. No change to coverage.
12/04/2009	Medical Policy Committee approval
12/16/2009	Medical Policy Implementation Committee approval. Title changed from "ACTH Gel (Adrenocorticotrophic Hormone)" to "Repository Corticotropin Injection (ACTH Gel, H.P. Acthar Gel)". No change to coverage eligibility.
07/01/2010	Medical Policy Committee approval
07/21/2010	Medical Policy Implementation Committee approval. Acute gout and childhood epilepsy added as investigational conditions.
12/01/2010	Medical Policy Committee review
12/15/2010	Medical Policy Implementation Committee approval. Added "in infants and children less than two years of age" to repository corticotropin injection for the treatment of infantile spasms (West's syndrome) coverage eligibility statement.
12/08/2011	Medical Policy Committee review
12/21/2011	Medical Policy Implementation Committee approval. No change to coverage.
12/06/2012	Medical Policy Committee review
12/19/2012	Medical Policy Implementation Committee approval. No change to coverage.
09/05/2013	Medical Policy Committee review
09/18/2013	Medical Policy Implementation Committee approval. Policy now states that covered indications include infantile spasms and multiple sclerosis only. Updated background, source, rationale, investigational, and not medically necessary sections.
09/04/2014	Medical Policy Committee review
09/17/2014	Medical Policy Implementation Committee approval. No change to coverage.

Repository Corticotropin Injection

Policy # 00230

Original Effective Date: 07/16/2008

Current Effective Date: 01/01/2026

09/03/2015	Medical Policy Committee review
09/23/2015	Medical Policy Implementation Committee approval. No change to coverage.
09/08/2016	Medical Policy Committee review
09/21/2016	Medical Policy Implementation Committee approval. No change to coverage.
01/01/2017	Coding update: Removing ICD-9 Diagnosis Codes
09/07/2017	Medical Policy Committee review
09/20/2017	Medical Policy Implementation Committee approval. No change to coverage.
06/07/2018	Medical Policy Committee review
06/20/2018	Medical Policy Implementation Committee approval. Policy now states that repository corticotropin is considered not medically necessary for exacerbations of multiple sclerosis. Updated background information, rationale, and references.
06/06/2019	Medical Policy Committee review
06/19/2019	Medical Policy Implementation Committee approval. No change to coverage.
06/04/2020	Medical Policy Committee review
06/10/2020	Medical Policy Implementation Committee approval. No change to coverage.
06/03/2021	Medical Policy Committee review
06/09/2021	Medical Policy Implementation Committee approval. No change to coverage.
06/02/2022	Medical Policy Committee review
06/08/2022	Medical Policy Implementation Committee approval. Added new product, Cortrophin Gel, to policy. Updated title and background information to remove reference to obsolete products.
06/01/2023	Medical Policy Committee review
06/14/2023	Medical Policy Implementation Committee approval. No change to coverage.
09/18/2023	Coding update
06/06/2024	Medical Policy Committee review
06/12/2024	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
06/05/2025	Medical Policy Committee review
06/11/2025	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
11/06/2025	Medical Policy Committee review
11/12/2025	Medical Policy Implementation Committee approval. Removed “When Services are Considered Not Medically Necessary” and “When Services are Considered Investigational” sections. Added “When Services are Not Covered” section to reflect contract exclusion for these products (except for the diagnosis of infantile spasms).

Next Scheduled Review Date: 11/2026

Repository Corticotropin Injection

Policy # 00230

Original Effective Date: 07/16/2008

Current Effective Date: 01/01/2026

Coding

The five character codes included in the Louisiana Blue Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)†, copyright 2024 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Louisiana Blue Medical Policy Coverage Guidelines is with Louisiana Blue and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Louisiana Blue Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Louisiana Blue Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	No codes
HCPCS	J0801, J0802 Delete code effective 07/01/2025: J3490, J3590
ICD-10 Diagnosis	All related diagnoses

****Medically Necessary (or “Medical Necessity”)** - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

Repository Corticotropin Injection

Policy # 00230

Original Effective Date: 07/16/2008

Current Effective Date: 01/01/2026

For these purposes, “nationally accepted standards of medical practice” means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: If the Patient’s health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

NOTICE: Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage.