

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Investigational or experimental services are not covered. This includes any drug, device, procedure, or service provided under the investigational arm of a clinical trial or clinical study. These services are excluded from coverage under benefits.

When Services May Be Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Implantable Ventricular Assist Devices

Destination Therapy

Based on review of available data, the Company may consider implantable ventricular assist devices (VADs) as a destination therapy **eligible for coverage**** when all of the following criteria are met:

- Implantable VAD used in accordance with Food and Drug Administration (FDA) approval; **AND**
- Adult individual with end-stage heart failure; **AND**
- Documented New York Heart Association (NYHA) Class III heart failure with dyspnea upon mild physical activity or NYHA Class IV; **AND**
- Left ventricular ejection fraction less than or equal to 25%; **AND**
- Inotrope-dependent **OR** cardiac index less than 2.2 liters/min/m² while not on inotropes **AND** also meeting one of the following:
 - On optimal medical management, based on current heart failure practice guidelines for at least 45 of the last 60 days and are failing to respond; **OR**
 - Advanced heart failure for at least 14 days and dependent on intra-aortic balloon pump for greater than or equal to 7 days.

Bridge to Transplantation

Based on review of available data, the Company may consider implantable ventricular assist devices (VADs) as a bridge to heart transplantation for adults **eligible for coverage**** when both coverage criteria are met:

- Implantable VAD used in accordance with FDA approval; **AND**

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

- Adult individual is currently listed as a heart transplantation candidate **OR** is undergoing evaluation to determine candidacy for heart transplantation **AND** is not expected to survive until a donor heart can be obtained.

Based on review of available data, the Company may consider implantable VADs as a bridge to heart transplantation for children **eligible for coverage**** when all of the following criteria are met:

- Implantable VAD used in accordance with FDA approval (including humanitarian device exemption approval); **AND**
- Request is for a child 16 years old or younger; **AND**
- Individual is currently listed as a heart transplantation candidate **OR** is undergoing evaluation to determine candidacy for heart transplantation **AND** is not expected to survive until a donor heart can be obtained.

Post cardiectomy Setting/Bridge to Recovery

Based on review of available data, the Company considers implantable VADs with FDA approval or clearance in the post cardiectomy setting in individuals who are unable to be weaned off cardiopulmonary bypass to be **eligible for coverage****.

Total Artificial Hearts

Based on review of available data, the Company may consider total artificial hearts (TAHs) as a bridge to heart transplantation **eligible for coverage**** when all of the following criteria are met:

- TAH is used in accordance with FDA approval; **AND**
- Individual has biventricular heart failure; **AND**
- There are no other reasonable medical or surgical treatment options; **AND**
- Ineligible for other univentricular or biventricular support devices; **AND**
- Currently listed as heart transplantation candidate **OR** is undergoing evaluation to determine candidacy for heart transplantation **AND** is not expected to survive until a donor heart can be obtained; **AND**
- Absence of any of the contraindications:
 - Chronic irreversible hepatic, renal, or respiratory failure
 - Active systemic infection
 - Coagulation disorders and inability to tolerate anticoagulation therapy.

Percutaneous Ventricular Assist Devices

Based on review of available data, the Company may consider left heart percutaneous ventricular assist devices (pVADs) for the treatment of cardiogenic shock **eligible for coverage**** when all of the following criteria are met:

- Left heart pVAD is FDA approved; **AND**
- Request is for an adult individual (18 years or older); **AND**
- Acute myocardial infarction of less than 36 hours duration with one of the following findings:

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

- ST- segment elevation myocardial infarction (STEMI) with new onset ST-segment elevation; **OR**
 - New onset ST segment depression or left bundle branch block (LBBB) and emergency angiography demonstrating acute proximal occlusion of coronary artery; **AND**
- Cardiogenic shock of less than 24 hours duration confirmed by hypotension and all of the following:
 - Systolic blood pressure less than 100 mmHg or ongoing vasopressor support, e.g., dopamine, norepinephrine or epinephrine; **AND**
 - End-organ hypoperfusion with arterial lactate greater than or equal to 2.5 mmol/L; **AND**
 - Left ventricle ejection fraction (LVEF) less than 45%; **AND**
- Absence of ANY one of the following:
 - Individual has been resuscitated from out-of-hospital cardiac arrest and remained comatose on arrival to the cardiac catheterization laboratory with persistent Glasgow coma scale less than 8 after return of spontaneous circulation;
Note: Cardiac arrest occurring in the ambulance or after arrival to the hospital is NOT an exclusion criterion.
 - Severe right ventricle failure by echocardiogram;
 - Shock duration greater than 24 hours;
 - Other causes of shock, e.g., hypovolemia, sepsis, pulmonary embolism, anaphylaxis;
 - Shock due to mechanical complication to myocardial infarction
 - Papillary muscle rupture
 - Rupture of ventricular septum or ventricular free wall;
 - Severe aortic valve regurgitation or stenosis;
 - Established diagnosis of severe peripheral arterial obstructive disease or abnormalities of the aorta that would preclude pVAD placement;
 - Presence of a mechanical aortic valve prosthesis;
 - Left ventricle thrombus;
 - Infective endocarditis.

When Services Are Considered Not Medically Necessary

Based on review of available data, the Company considers the use of percutaneous VADs when criteria are not met to be **not medically necessary**.**

Based on review of available data, the Company considers other applications of implantable ventricular assist devices (VADs) or total artificial hearts (TAHs) including, but not limited to, the use of TAHs as destination therapy to be **not medically necessary**.**

When Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

Based on review of available data, the Company considers the use of right heart percutaneous VADs to be **investigational**.*

Based on review of available data, the Company considers the use of non-FDA-approved or cleared implantable VADs or TAHs to be **investigational**.*

Policy Guidelines

The intent of treatment may evolve over the course of treatment; for example, there is not necessarily a strict delineation between bridge to transplant and destination therapy.

Some ventricular assist devices (VADs) have approval from the U.S. Food and Drug Administration (FDA) for the pediatric population. For example, the Berlin Heart EXCOR Pediatric VAD has FDA approval through the humanitarian device exemption process. This device is indicated for children with severe isolated left ventricular or biventricular dysfunction who are candidates for cardiac transplant and require circulatory support. The HeartMate3^{TM†} received approval for expanded approval for pediatric patients with advanced refractory left ventricular heart failure in 2020. As of April 15, 2024, the HeartMate 3 devices are under a Class I FDA recall due to the accumulation of biological material within the device - a serious complication that can lead to obstruction, posing significant risks of severe injury or death. In April 2025, Abbott removed HeartMate Mobile Power Unit (used with HeartMate II and HeartMate 3) due to instances of sudden power loss.

In general, candidates for bridge to transplant implantable VADs are those who are considered appropriate heart transplant candidates but who are unlikely to survive the waiting period until a human heart donor is available. Some studies have included the following hemodynamic selection criteria: either a left atrial pressure of 20 mm Hg or a cardiac index of less than 2.0 L/min/m while receiving maximal medical support. Individuals with VADs are classified by the United Network for Organ Sharing as status I (ie, persons who are most ill and are considered the highest priority for transplant).

The median duration for time on the device is between 20 and 120 days.

Contraindications for bridge to transplant VADs and total artificial hearts include conditions that would generally exclude individuals for heart transplant. Such conditions are chronic irreversible hepatic, renal, or respiratory failure; systemic infection; coagulation disorders; and inadequate psychosocial support. Due to potential problems with adequate function of the VAD or total artificial heart, implantation is also contraindicated in individuals with uncorrected valvular disease.

The Centers for Medicare and Medicaid Services requires that “Beneficiaries receiving a VAD must be managed by an explicitly identified, cohesive, multidisciplinary team of medical professionals with appropriate qualifications, training, and experience. The team embodies collaboration and dedication across medical specialties to offer optimal patient-centered care. Collectively, the team must ensure that patients and caregivers have the knowledge and support necessary to participate in

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

informed decision making. The team members must be based at the facility and must include individuals with experience working with patients before and after placement of a VAD.

The team must include, at a minimum:

- At least 1 physician with cardiothoracic surgery privileges and individual experience implanting at least 10 durable, intracorporeal, left ventricular assist devices over the course of the previous 36 months with activity in the last year.
- At least 1 cardiologist trained in advanced heart failure with clinical competence in medical- and device-based management including VADs, and clinical competence in the management of patients before and after placement of a VAD.
- A VAD program coordinator.
- A social worker.
- A palliative care specialist.”

Background/Overview

Heart Failure

According to a 2025 report from the American Heart Association and based on data collected from 2017 to 2020, roughly 6.7 million Americans ages 20 years or older had heart failure during that time frame. Prevalence of heart failure is projected to affect more than 8 million people 18 years of age and older by the year 2030. Between 2015 and 2018, the prevalence of heart failure was highest in non-Hispanic Black males. Based on data from the Multi-Ethnic Study of Atherosclerosis (MESA), in those without baseline cardiovascular disease, Black individuals had the highest risk of developing heart failure (4.6 per 1000 person-years), followed by Hispanic (3.5 per 1000 person-years), White (2.4 per 1000 person-years), and Chinese individuals (1.0 per 1000 person-years). Similar findings were demonstrated in the Atherosclerosis Risk in Communities (ARIC) Community Surveillance data, in which Black men and women had the highest burden of new-onset heart failure cases and the highest-age adjusted 30-day case fatality rate in comparison to White men and women. Higher risk reflected differential prevalence of hypertension, diabetes, and low socio-economic status.

Heart failure may be the consequence of a number of etiologies, including ischemic heart disease, cardiomyopathy, congenital heart defects, or rejection of a heart transplant. The reduction of cardiac output is considered to be severe when systemic circulation cannot meet the body's needs under minimal exertion. Heart transplantation improves quality of life and had a reported survival rate of nearly 92% or transplants performed in 2022. The number of candidates for transplants exceeds the supply of donor organs; thus the interest in the development of mechanical devices.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

A number of implantable ventricular assist devices (VADs) and artificial heart systems have been U.S. Food and Drug Administration (FDA) approved through a Humanitarian Device Exemption, 510(k), or premarket approval regulatory pathway. This section discusses currently marketed devices.

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

FDA maintains a list of recent device recalls at <https://www.fda.gov/medical-devices/medical-device-safety/medical-device-recalls>.

Ventricular Assist Devices

Implantable VADs are attached to the native heart, which may have enough residual capacity to withstand a device failure in the short term. In reversible heart failure conditions, the native heart may regain some function, and weaning and explanting of the mechanical support system after months of use has been described. VADs can be classified as internal or external, electrically or pneumatically powered, and pulsatile or continuous-flow. Initial devices were pulsatile, mimicking the action of a beating heart. More recent devices may use a pump, which provides continuous flow. Continuous devices may move blood in a rotary or axial flow.

Surgically implanted VADs represent a method of providing mechanical circulatory support for patients not expected to survive until a donor heart becomes available for transplant or for whom transplantation is contraindicated or unavailable. VADs are most commonly used to support the left ventricle but right ventricular and biventricular devices may be used. The device is larger than most native hearts, and therefore the size of the patient is an important consideration; the pump may be implanted in the thorax or abdomen or remain external to the body. Inflow to the device is attached to the apex of the failed ventricle, while outflow is attached to the corresponding great artery (aorta for the left ventricle, a pulmonary artery for the right ventricle). A small portion of the ventricular wall is removed for insertion of the outflow tube; extensive cardiectomy affecting the ventricular wall may preclude VAD use.

The intent of treatment may evolve over the course of treatment; for example, there is not necessarily a strict delineation between bridge to transplant and destination therapy, and transplant eligibility can change.

Table 1 lists the VADs currently available in the US. The HeartWare VAD System was discontinued in June 2021 due to evidence from observational studies demonstrating a higher frequency of neurological adverse events and mortality with the system compared to other commercially available left VADs.

- The DeBakey VAD Child received FDA Humanitarian Device Exemption (HDE) approval in 2004, offering a temporary lifeline for pediatric patients with end-stage heart failure as they awaited heart transplantation. However, this device was replaced by the HeartAssist 5 VAD which is not currently available for use as it was an investigational device that was discontinued.
- As of April 15, 2024, both the HeartMate II and HeartMate 3 devices have been placed under a Class I FDA recall in response to the accumulation of biological material within the devices, an issue that can result in serious obstructions and significantly increase the risk of severe injury or death. In April 2025, Abbott further removed the HeartMate Mobile Power Unit, which is used with both the HeartMate II and HeartMate 3, following reports of sudden and unexpected power loss.

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

- The Abbott CentriMag Circulatory Support System received FDA approval in 2019 to provide longer-term life support to critically ill patients. In 2023, it was approved for longer-term use in adults when extracorporeal membrane oxygenation.

Table 1. Available Ventricular Assist Devices

Device	Manufacturer	Approval Date	FDA Clearance	PMA, HDE, or 510(k) No.	Indication
HeartMate II	Thoratec (Abbott)	Apr 2008	PMA	P060040	Bridge to transplant and destination
CentriMag	Thoratec (Abbott)	Dec 2019	PMA	P170038	Postcardiotomy, bridge to decision
Berlin Heart EXCOR Pediatric VAD	Berlin	Jun 2017	PMA	P160035	Bridge to transplant or recovery
HeartMate 3 Left Ventricular Assist System	Thoratec (Abbott)	Aug 2017 Oct 2018	PMA PMA	P160054 P160054/S008	Bridge to transplant and destination

FDA: U.S. Food and Drug Administration; HDE: humanitarian device exemption; PMA: premarket approval; VAD: ventricular assist device.

Total Artificial Heart

The total artificial heart (TAH) is a biventricular device that completely replaces the function of the diseased heart. An internal battery requires frequent recharging from an external power source. Many systems use a percutaneous power line, but a transcutaneous power-transfer coil allows for a system without lines traversing the skin, possibly reducing the risk of infection. Because the native heart must be removed, failure of the device is synonymous with cardiac death.

Currently the Syncardia Temporary Total Artificial Heart (Syncardia Systems) is the only Total Artificial Heart available in the US (Table 2). The AbioCor Total Artificial Heart was FDA approved under the Humanitarian Device Exemption program in 2006, but is no longer being marketed or in development.

Table 2. Available Total Artificial Heart

Device	Manufacturer	Approval Date	FDA Clearance	PMA No.	Indication
SynCardia Temporary Total Artificial	SynCardia Systems	2004	510(k)	P030011	Bridge to transplant in cardiac transplant-eligible candidates at risk of imminent death from biventricular failure.

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

Heart (Formerly CardioWest Total Artificial Heart and Jarvik Total Artificial Heart)					
---	--	--	--	--	--

FDA: U.S. Food and Drug Administration; PMA: premarket approval.

Percutaneous Ventricular Assist Devices

Some circulatory assist devices are placed percutaneously (i.e., are not implanted). They may be referred to as percutaneous VADs (pVADs). Two different pVADs have been developed, the TandemHeart and the Impella device (Table 3).

In the TandemHeart System, a catheter is introduced through the femoral vein and passed into the left atrium via transseptal puncture. Oxygenated blood is then pumped from the left atrium into the arterial system via the femoral artery. LivaNova, the company that acquired CardiacAssist Inc. (the original developer of the TandemHeart), announced plans to wind down its Advanced Circulatory Support business, including the TandemHeart, in 2024. However, it is possible that some components, such as cannulas, may still be available for purchase.

The Impella device is introduced through a femoral artery catheter. In this device, a small pump is contained within the catheter placed into the left ventricle. Blood is pumped from the left ventricle, through the device, and into the ascending aorta. As of June 5, 2023, the FDA has issued a Class I recall for specific Impella 5.5 devices due to purge fluid leaks. On October 10, 2025, the FDA issued an alert on the automated Impella controller correction due to a cybersecurity issue.

Devices in which most of the system's components are external to the body are for short-term use (6 hours to 14 days) only, due to the increased risk of infection and need for careful, in-hospital monitoring. Adverse events associated with pVAD include access site complications such as bleeding, aneurysms, or leg ischemia. Cardiovascular complications can also occur, such as perforation, myocardial infarction, stroke, and arrhythmias.

Table 3. Available Percutaneous Ventricular Assist Devices

Device	Manufacturer	Approval Date	FDA Clearance	PMA, 510(k) No.	Indication
TandemHeart	Cardiac Assist (LivaNova)	Sep 2011	510(k)	K110493	Temporary left ventricular bypass of ≤6 h

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

Impella CP	Abiomed	Nov 2016	PMA	P140003	• Temporary (≤ 6 hours) ventricular support devices indicated for use during high-risk PCI • Temporary ventricular support for ≤ 4 days in cardiogenic shock
Impella 5.5	Abiomed	Nov 2016	PMA	P140003	Temporary ventricular support for ≤ 14 days in cardiogenic shock

FDA: U.S. Food and Drug Administration; PCI: percutaneous coronary intervention; PMA: premarket approval.

Rationale/Source

This medical policy was developed through consideration of peer-reviewed medical literature generally recognized by the relevant medical community, U.S. Food and Drug Administration approval status, nationally accepted standards of medical practice and accepted standards of medical practice in this community, technology evaluation centers, reference to regulations, other plan medical policies, and accredited national guidelines.

Description

A ventricular assist device (VAD) is mechanical support attached to the native heart and vessels to augment cardiac output. The total artificial heart (TAH) replaces the native ventricles and is attached to the pulmonary artery and aorta; the native heart is typically removed. Both the VAD and TAH may be used as a bridge to heart transplantation or as destination therapy. The VAD has also been used as a bridge to recovery in individuals with reversible conditions affecting cardiac output.

Summary of Evidence

Ventricular Assist Device

For individuals who have end-stage heart failure who receive a ventricular assist device (VAD) as a bridge to transplant, the evidence includes a randomized controlled trial (RCT), single-arm trials, and observational studies. Relevant outcomes are overall survival (OS), symptoms, functional outcomes, quality of life (QOL), and treatment-related mortality and morbidity. There is a substantial body of evidence from clinical trials and observational studies supporting implantable VADs as a bridge to transplant in patients with end-stage heart failure, possibly reducing mortality as well as improving QOL. These studies have reported that substantial numbers of patients have survived to transplant in situations in which survival would not be otherwise expected. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

For individuals who have end-stage heart failure who receive a VAD as destination therapy, the evidence includes RCTs and multiple single-arm studies. Relevant outcomes are OS, symptoms, functional outcomes, QOL, and treatment-related mortality and morbidity. A well-designed trial with 2 years of follow-up data has demonstrated an advantage of implantable VADs as destination therapy for patients ineligible for a heart transplant. Despite an increase in adverse events, both mortality and QOL appear to be improved for these patients. A more recent trial comparing VADs has broader inclusion criteria and supports that criteria move away from use of transplant ineligibility, as treatment may evolve over the course of treatment. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Total Artificial Heart

For individuals who have end-stage heart failure who receive a total artificial heart (TAH) as a bridge to transplant, the evidence includes case series. Relevant outcomes are OS, symptoms, functional outcomes, QOL, and treatment-related mortality and morbidity. Compared with VADs, the evidence for TAHs in these settings is less robust. However, given the lack of medical or surgical options for these patients and the evidence case series provide, TAH is likely to improve outcomes for a carefully selected population with end-stage biventricular heart failure awaiting transplant who are not appropriate candidates for a left VAD. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have end-stage heart failure who receive a TAH as destination therapy, the evidence includes 2 case series. Relevant outcomes are OS, symptoms, functional outcomes, QOL, and treatment-related mortality and morbidity. The body of evidence for TAHs as destination therapy is too limited to draw conclusions. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Percutaneous Ventricular Assist Device

For individuals with infarct-related cardiogenic shock who receive a percutaneous VAD (pVAD), the evidence includes three small, randomized trials that did not show a clinical benefit of microaxial flow pumps, registry studies that showed excess bleeding among patients who received a microaxial pump, and a more recent international, multicenter, randomized, open-label trial funded by the Danish Heart Foundation and Abiomed (DanGer Shock trial, NEJM 4/2024). DanGer Shock trial tested the hypothesis that routine use of a microaxial flow pump in addition to standard guideline-directed therapies in patients with STEMI-related cardiogenic shock results in a lower mortality than standard care alone. Patients with ST segment elevation myocardial infarction and cardiogenic shock were assigned to receive a microaxial flow pump (Impella CP) plus standard care or standard of care alone. The primary end point was death from any cause at 180 days (6 months). A composite safety end point was severe bleeding, limb ischemia, hemolysis, device failure, or worsening aortic regurgitation. From January 2013 through July 2023 a total of 1211 patients underwent screening. A total of 360 adult patients underwent randomization, 355 were included in final analysis (179 in the microaxial-flow-pump and 176 in the standard-care group). The median age of the patients was 67 years, and 79.2% were men. The median arterial lactate level was 4.5 mmol/L, the median systolic BP was 82 mm Hg, and the median LV EF was 25%. The trial was conducted in a more homogenous patient population (a mandatory elevation in arterial lactate level in the absence of a cardiac arrest

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

led to the identification of a patient population with profound LV failure). In the microaxial-flow-pump group, treatment was escalated to another mechanical circulatory support system in 28 patients (15.6%), and in the standard-care group in 37 patients (21%). Death from any cause occurred in 82 of 179 patients (45.8%) in the microaxial-flow-pump group and in 103 of 176 patients (58.5%) in the standard-care group (hazard ratio, 0.74; 95% confidence interval 0.55 to 0.99; $P=0.04$). The number needed to treat to avoid 1 death was 8. A composite safety end-point event occurred in 43 patients (24.0%) in the microaxial-flow-pump group and in 11 patients (6.2%) in the standard-care group (relative risk, 4.74; 95% CI, 2.36 to 9.55). In the microaxial-flow-pump group, the number needed to harm was 6. The relative risk (pump group vs. standard-care group) of moderate or severe bleeding was 2.06, of limb ischemia 5.15, and of sepsis with a positive blood culture 2.79. Renal-replacement therapy was administered to 75 patients (41.9%) in the pump group and to 47 patients (26.7%) in the standard-care group (relative risk, 1.98; 95% CI, 1.27 to 3.09). This considerably higher use of renal-replacement therapy is higher than that observed in the extracorporeal life support trial (ECLS-SHOCK trial). Authors noted that it remains a priority to address the prevention of serious adverse events that occur as a result of treatment with a microaxial flow pump. Trial limitations include strict inclusion and exclusion criteria; the results cannot be extrapolated to patients who remain comatose after cardiac arrest, patients without ST-segment elevation myocardial infarction, patients without an elevation in the arterial lactate level, and patients who have more prominent biventricular failure. The trial was conducted in a small number of centers in Denmark, Germany, and the United Kingdom, so results may differ in health care systems in other countries with more racial diversity. The trial was not blinded therefore authors could not exclude the possible effect on therapeutic decisions made by treating physicians. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome. Clinical input was obtained and supported coverage criteria that are based on the inclusion and exclusion criteria in the DanGer Shock trial, which is also consistent with the 2025 ACC/AHA Joint Committee on Clinical Practice Guidelines (Guideline for the management of patients with acute coronary syndromes).

For other individuals with cardiogenic shock who receive a percutaneous VAD (pVAD), the evidence includes RCTs, observational studies, and a systematic review. Relevant outcomes are OS, symptoms, morbid events, functional outcomes, QOL, and treatment-related mortality and morbidity. Four RCTs of pVAD versus intra-aortic balloon pump (IABP) for patients in cardiogenic shock failed to demonstrate a mortality benefit and reported higher complication rates with pVAD use. Comparative observational studies and a long-term follow-up study were consistent with the RCT evidence. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with cardiogenic shock refractory to IABP therapy who receive a pVAD, the evidence includes case series. Relevant outcomes are OS, symptoms, morbid events, functional outcomes, QOL, and treatment-related mortality and morbidity. Case series of patients with cardiogenic shock refractory to IABP have reported improved hemodynamic parameters following pVAD placement. However, these uncontrolled series do not provide evidence that pVADs improve mortality, and high rates of complications have been reported with pVAD use. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

For individuals who undergo high-risk cardiac procedures who receive a pVAD, the evidence includes RCTs, observational studies, and systematic reviews of these trials. Relevant outcomes are OS, symptoms, morbid events, functional outcomes, QOL, and treatment-related mortality and morbidity. Randomized controlled trials, controlled and uncontrolled observational studies, and systematic reviews of these studies have generally not demonstrated a benefit of pVAD used as ancillary support for patients undergoing high-risk cardiac procedures.

The PROTECT II trial (10/2012) led to the FDA approval of Impella as a temporary support for high-risk percutaneous cardiac interventions. Study included participants with PCI on an unprotected left main or last patent coronary vessel with LVEF $\leq 35\%$, or PCI for 3 vessel disease with LVEF $\leq 30\%$, and excluded patients with myocardial infarction, LV thrombus, PLT count $\leq 75,000/\text{mm}^3$, creatinine $\geq 4 \text{ mg/dL}$, or severe PVD precluding passage of the catheter. The primary endpoint of 30-day MAE in the PROTECT II study occurred in 34.3% of the per protocol (PP) patients in the Impella 2.5 arm compared with 42.2% of PP patients in the IABP arm ($P=0.092$). This study had important limitations. Because of the determination of futility, only 69% (452) of the planned enrollment occurred. A significant learning curve occurred in this trial with improvement in safety for Impella-supported patients in the last half of the trial. The trial did not enroll the number of patients it was powered for, therefore definitive statements concerning the primary end point are speculative. The trial could not be blinded of the presence of Impella support which led to a greater and more aggressive use of rotational atherectomy, and the differences confounded the analysis. Because the difference in 30-day major adverse events did not reach statistical significance for the entire study, the analysis of 90-day events remains exploratory.

Restore EF study (8/2022) was largest to date study looking at LV EF changes 30 days after high-risk PCI. It was an observational nonrandomized multicenter retrospective analysis of prospectively collected observational data set to assess 90-day LVEF in patients who survived with no additional procedure prior to 90-day assessment. 36 out of 495 screened patients (7.3%) who died before day 60 were excluded from assessment; 406 patients were evaluated. The study reported 30-day mortality of 6.9% (all screened patients), increasing to 9.5% at 180 days. Authors concluded that in an ideal cohort there is a signal that high-risk PCI affords improvement in 90-day LV EF (from median 35% to 45% with $P<.0001$, most pronounced in those with a baseline LVEF $\leq 20\%$, no change in patients with LVEF $> 45\%$ at baseline) along with significant relief of angina and heart failure symptoms, and that this hypothesis-generating finding merits assessment in large studies and randomized controlled trials. Important limitations include the lack of a comparator, investigator-reported data with no angiographic or echocardiographic core lab, or independent clinical events committee adjudication, and unknown impact on LVEF from optimal heart failure medical therapy (data on medications were not collected). A significant number of patients (61.8%) who were included in the final study population did not undergo 90-day follow-up LVEF assessment (due to patients missing appointment or physician not ordering LVEF assessment).

The PROTECT III study was sponsored by Abiomed. It was a prospective, single-arm, FDA post-marketing approval (PMA) study of Impella 2.5[®]† and Impella CP[®]‡ in high-risk PCI. Trial included 898 patients enrolled at 45 sites in the United States between 3/2017 and 7/2019. Using cVAD registry database data, PROTECT III demonstrates a reduction in the primary endpoint of death,

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

stroke, myocardial infarction and repeat procedures at 90 days with Impella-supported protected PCI, compared to PROTECT II.

Impella-supported PCI in high-risk patients with complex coronary artery disease and reduced LV function trial sponsored by Abiomed (the PROTECT IV trial, NCT04763200) is currently recruiting participants. It is an on-label, prospective, multi-center randomized controlled trial that is designed to provide the level of clinical evidence needed to achieve a global class I guideline recommendation (a strong recommendation, indicating that intervention is effective and should be performed in most patients in most circumstances) for Impella in high-risk PCI. The purpose of this study is to assess if using the Impella[®] CP (or Impella[®] 2.5)[‡] device during high-risk PCI in patients with reduced left-sided heart function will result in an improvement in symptoms, heart function and health outcomes after a PCI, compared to the current standard of care. Inclusion criteria: adult patients (18-90 years of age) with LVEF $\leq$ 40% in setting of chronic coronary syndrome or NSTEMI, or STEMI $\geq$ 24 hours and $<$ 30 days after symptom onset and LVEF $\leq$ 30%, and need of complex PCI (triple vessel disease, left main disease, ostial LAD and ostial LCX disease, or intervention of the last remaining vessel). Estimated study completion is 10/2027.

There is a lack of large-scale, high-quality RCTs demonstrating clear clinical benefits in the use of any of the mechanical circulatory support (MCS) devices during high-risk PCI. European Society of Cardiology guidelines did not make specific recommendations on the use of MCS during revascularization while the American College of Cardiology made only a class IIb recommendation (weak recommendation for which it is unknown whether benefit will outweigh the risk) for the use of hemodynamic support device as an adjunct to complex PCI for selected high-risk patients to prevent hemodynamic compromise. Ongoing and future research will shed light on unanswered questions including those around the definition of complex, high-risk and indicated PCI, patient selection, device selection, evaluation of new devices and timing of support initiation.

Additionally, 2 nonrandomized studies have compared ventricular tachycardia (VT) ablation with pVAD or IABP. Both studies demonstrated that patients who had pVAD support spent less time in unstable VT than patients without pVAD support. However, the current evidence does not support conclusions about the use of pVAD for VT ablation. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Supplemental Information

Clinical Input From Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American Association for Thoracic Surgery et al

In 2020, the American Association for Thoracic Surgery and the International Society for Heart and Lung Transplantation published guidelines on selected topics in mechanical circulatory support (MCS), including recommendations on the use of pVADs (Table 4). The guideline authors noted, "Compared with IABP [intraaortic balloon pump], contemporary percutaneous circulatory support devices provide a significant increase in cardiac index and mean arterial pressure; however, reported 30-day outcomes are similar."

Table 4. 2020 Guidelines on Mechanical Circulatory Support

Recommendation	COE	LOE
"Percutaneous LV to aorta pumps of appropriate size should be considered for cardiogenic shock from primary LV failure."	IIA	B

COE: class of evidence; LOE: level of evidence; LV: left ventricular.

American College of Cardiology Foundation et al

In 2017, the American College of Cardiology Foundation, American Heart Association (AHA), and Heart Failure Society of American published a focused update of the 2013 recommendations released by the American College of Cardiology Foundation and AHA. Left ventricular assist device was 1 of several treatment options recommended for patients with refractory New York Heart Association class III or IV heart failure (stage D). If symptoms were not improved after guideline-directed management and therapy, which included pharmacologic therapy, surgical management and/or other devices, then a left ventricular assist device would be an additional treatment option.

The 2017 update focused on changes in sections regarding biomarkers, comorbidities, and prevention of heart failure, while many of the previous recommendations remained unchanged. The American College of Cardiology Foundation and AHA (2013) released guidelines for the management of heart failure that included recommendations related to the use of MCS, including both durable and nondurable MCS devices. The guidelines categorized pVADs and extracorporeal ventricular assist devices (VADs) as nondurable MCS devices. Since the 2017 update, these guidelines have been updated regularly, with the most recent update occurring in 2022. Table 5 provides recommendations on MCS devices from the most recently updated guideline iteration.

Table 5. AHA/ACC/HFSA Guidelines on Mechanical Circulatory Support

Recommendation	COE ^a	LOE ^b
"In select patients with advanced HFrEF with NYHA class IV symptoms who are deemed to be dependent on continuous intravenous inotropes or temporary MCS, durable LVAD implantation is effective to improve functional status, QOL, and survival."	I	A

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

Recommendation	COE ^a	LOE ^b
"In select patients with advanced HFrEF who have NYHA class IV symptoms despite GDMT, durable MCS can be beneficial to improve symptoms, improve functional class, and reduce mortality."	IIA	B-R
"In patients with advanced HFrEF and hemodynamic compromise and shock, temporary MCS, including percutaneous and extracorporeal ventricular assist devices, are reasonable as a 'bridge to recovery' or 'bridge to decision'"	IIA	B-NR

ACC: American College of Cardiology; AHA: American Heart Association; COE: class of evidence; GDMT: guideline-directed medical therapy; HFrEF: heart failure with reduced ejection fraction; HFSA: Heart Failure Society of America; LOE: level of evidence; LVAD: left ventricular assist device; MCS: mechanical circulatory support; NYHA: New York Heart Association; QOL: quality of life; RCT: randomized controlled trial.

^aI: Strong; IIa: Moderate.

^bA: high quality evidence from more than 1 RCT; B-R: Moderate-quality evidence from 1 or more RCTs; B-NR: Moderate-quality evidence from 1 or more well-designed, well-executed nonrandomized studies, observational studies, or registry studies.

2025 ACC/AHA/ACEP/ NAEMSP/ SCAI Guideline for the Management of Patients With Acute Coronary Syndromes: A Report of the American College of Cardiology/ American Heart Association Joint Committee on Clinical Practice Guidelines (based on a comprehensive literature search from July 2023 to April 2024) included grade 2a recommendation (a moderate recommendation for which benefit most likely outweighs the risk): “In selected patients with STEMI and severe or refractory cardiogenic shock, insertion of a microaxial intravascular flow pump is reasonable to reduce death.” Based on the results of the DanGer-SHOCK trial, “use of a microaxial flow pump is reasonable to reduce mortality in patients with STEMI and cardiogenic shock who have clinical features consistent with the inclusion criteria of the DanGer-SHOCK trial. In particular, patients with STEMI who present with SCAI shock stages C, D, or E, who are noncomatose and have adequate peripheral vasculature to accommodate large-bore access are reasonable candidates for the microaxial flow pump.”

One of the recommendations for mechanical complications of ACS is related to mechanical circulatory support (MCS) devices (2a): “In patients with a mechanical complication of ACS, short-term MCS devices are reasonable for hemodynamic stabilization as a bridge to surgery.” It is noted that no RCTs have evaluated the role of MCS devices in improving clinical outcomes in the setting of mechanical complications. In patients with ventricular septal rupture, the use of an IABP has been shown to reduce left-to-right shunting and improve hemodynamics in patients with and without cardiogenic shock. Favorable hemodynamic effects with IABP are also noted with acute ischemic mitral regurgitation.”

American Heart Association

In 2012, the AHA published recommendations for the use of MCS. These guidelines defined nondurable MCS as IABPs, extracorporeal membrane oxygenation, extracorporeal VADs, and

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

pVADs. Table 6 lists recommendations made on indications for the use of MCS, including durable and nondurable devices.

Table 6. 2012 Guidelines on Mechanical Circulatory Support

Recommendation	COE	LOE
"MCS for BTT indication should be considered for transplant-eligible patients with end-stage HF who are failing optimal medical, surgical, and/or device therapies and at high risk of dying before receiving a heart transplantation."	I	B
"Implantation of MCS in patients before the development of advanced HF ... is associated with better outcomes. Therefore, early referral of HF patients is reasonable."	IIA	B
"MCS with a durable, implantable device for permanent therapy or DT is beneficial for patients with advanced HF, high 1-year mortality resulting from HF, and the absence of other life-limiting organ dysfunction; who are failing medical, surgical, and/or device therapies; and who are ineligible for heart transplantation."	I	B
"Elective rather than urgent implantation of DT can be beneficial when performed after optimization of medical therapy in advanced HF patients who are failing medical, surgical, and/or device therapies."	IIA	C
"Urgent nondurable MCS is reasonable in hemodynamically compromised HF patients with end-organ dysfunction and/or relative contraindications to heart transplantation/durable MCS that are expected to improve with time and restoration of an improved hemodynamic profile." "These patients should be referred to a center with expertise in the management of durable MCS and patients with advanced HF."	IIA I	C C
"Patients who are ineligible for heart transplantation because of pulmonary hypertension related to HF alone should be considered for bridge to potential transplant eligibility with durable, long-term MCS."	IIA	B

BTT: bridge to transplant; COE: class of evidence; DT: destination therapy; HF: heart failure; LOE: level of evidence; MCS: mechanical circulatory support.

International Society for Heart and Lung Transplantation

The International Society for Heart and Lung Transplantation and the Heart Failure Society of America released a guideline on acute MCS in 2023. The guideline focuses on timing, patient and device selection of acute MCS, and periprocedural and postprocedural care for cardiogenic and pulmonary shock. They provide specific recommendations depending on which MCS device is chosen. Table 7 summarizes relevant recommendations for timing of acute MCS made in the guidelines. Additional recommendations related to specific devices is related to procedural considerations.

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

Table 7. ISHLT/HFSA Guideline on Acute Mechanical Circulatory Support

Recommendation	COR	LOE
"Acute MCS should be initiated as soon as possible in patients with CS who fail to stabilize or continue to deteriorate despite initial interventions."	I	B
"The use of acute MCS should be considered in patients with multiorgan failure to allow successful optimization of clinical status and neurologic assessment before placement of durable MCS or organ transplantation."	II	C

COR: class of recommendation; CS: cardiogenic shock; HFSA: Heart Failure Society of America; ISHLT: International Society for Heart and Lung Transplantation; LOE: level of evidence; MCS: mechanical circulatory support

Society for Cardiovascular Angiography and Interventions et al

In 2015, the Society for Cardiovascular Angiography and Interventions, the Heart Failure Society of America, the Society of Thoracic Surgeons, and the American College of Cardiology published a joint clinical expert consensus statement on the use of percutaneous MCS devices in cardiovascular care. This statement addressed IABPs, left atrial-to-aorta assist device (eg, TandemHeart), left ventricle-to-aorta assist devices (eg, Impella), extracorporeal membrane oxygenation, and methods of right-sided support. Specific recommendations were not made, but the statement reviews the use of MCS in patients undergoing high-risk percutaneous intervention, those with cardiogenic shock, and those with acute decompensated heart failure.

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

Medicare has a national coverage determination (NCD) for VADs. The NCD mandates coverage for VADs for the following indications:

- For support of blood circulation in the post cardiectomy setting, defined as the period following open-heart surgery.
 - If the VAD has U.S. Food and Drug Administration (FDA) approval for that purpose and are used according to the FDA-labeled indication
- For short-term (e.g., bridge-to-recovery and bridge-to-transplant) or long-term (e.g., destination therapy) mechanical circulatory support for patients who meet the following criteria:
 - Have New York Heart Association (NYHA) Class IV heart failure; and
 - Have a left ventricular ejection fraction (LVEF) $\leq 25\%$; and
 - Are inotrope dependent

OR

have a cardiac index < 2.2 L/min/m², while not on inotropes, and also meet 1 of the following:

- Are on optimal medical management, based on current heart failure practice guidelines for at least 45 out of the last 60 days and are failing to respond;

OR

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

- Have advanced heart failure for at least 14 days and are dependent on an IABP or similar temporary mechanical circulatory support for at least 7 days.

"Beneficiaries receiving VADs for DT [destination therapy] must be managed by an explicitly identified cohesive, multidisciplinary team of medical professionals with the appropriate qualifications, training, and experience.... The team members must be based at the facility and must include individuals with experience working with patients before and after placement of a VAD."

"Facilities must be credentialed by an organization approved by the Centers for Medicare & Medicaid Services."

Effective December 1, 2020, Artificial Hearts has been removed from the NCD Manual. Coverage determinations for artificial hearts and related devices shall be made by the Medicare Administrative Contractors.

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 8.

Table 8. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT01627821 ^a	Evaluation of the Jarvik 2000 Left Ventricular Assist System With Post-Auricular Connector--Destination Therapy Study	350	Mar 2025 (last updated Mar 2024)
NCT02387112	Early Versus Emergency Left Ventricular Assist Device Implantation in Patients Awaiting Cardiac Transplantation	102	Dec 2024 (last updated Oct 2023)
NCT04768322	Left Ventricular Assist Device (LVAD) Versus Guideline Recommended Medical Therapy in Ambulatory Advanced Heart Failure Patients (GDMT)	92	Feb 2029
<i>Unpublished</i>			

NCT: national clinical trial.

^a Denotes industry-sponsored or cosponsored trial.

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

References

1. Martin SS, Aday AW, Allen NB, et al. 2025 Heart Disease and Stroke Statistics: A Report of US and Global Data From the American Heart Association. *Circulation*. Feb 25 2025; 151(8): e41-e660. PMID 39866113
2. Martin SS, Aday AW, Almarzooq ZI, et al. 2024 Heart Disease and Stroke Statistics: A Report of US and Global Data From the American Heart Association. *Circulation*. Feb 20 2024; 149(8): e347-e913. PMID 38264914
3. Lewsey SC, Breathett K. Racial and ethnic disparities in heart failure: current state and future directions. *Curr Opin Cardiol*. May 01 2021; 36(3): 320-328. PMID 33741769
4. Organ Procurement and Transplantation Network. National Heart Patient and Graft Survival as of June 21, 2024. <https://insights.unos.org/OPTN-metrics/>.
5. FDA. Heart Pump Accessory Removal: Abbott Removes HeartMate Mobile Power Unit due to Instances of Sudden Power Loss. April 24, 2025.
6. Abbott's Life Support System Receives New FDA Clearances to Help Doctors Treat More Critically Ill Patients. 2023.
7. LivaNova to Wind Down Advanced Circulatory Support Business Unit. January 8, 2024
8. Acharya D, Loyaga-Rendon RY, Pamboukian SV, et al. Ventricular Assist Device in Acute Myocardial Infarction. *J Am Coll Cardiol*. Apr 26 2016; 67(16): 1871-80. PMID 27102502
9. FDA. Medical Device Recalls and Early Alerts. Update September 29, 2025.
10. Goldstein DJ, Oz MC, Rose EA. Implantable left ventricular assist devices. *N Engl J Med*. Nov 19 1998; 339(21): 1522-33. PMID 9819452
11. Mehra MR, Uriel N, Naka Y, et al. A Fully Magnetically Levitated Left Ventricular Assist Device - Final Report. *N Engl J Med*. Apr 25 2019; 380(17): 1618-1627. PMID 30883052
12. Colombo PC, Mehra MR, Goldstein DJ, et al. Comprehensive Analysis of Stroke in the Long-Term Cohort of the MOMENTUM 3 Study. *Circulation*. Jan 08 2019; 139(2): 155-168. PMID 30586698
13. Cowger JA, Naka Y, Aaronson KD, et al. Quality of life and functional capacity outcomes in the MOMENTUM 3 trial at 6 months: A call for new metrics for left ventricular assist device patients. *J Heart Lung Transplant*. Jan 2018; 37(1): 15-24. PMID 29153637
14. Goldstein DJ, Naka Y, Horstmannshof D, et al. Association of Clinical Outcomes With Left Ventricular Assist Device Use by Bridge to Transplant or Destination Therapy Intent: The Multicenter Study of MagLev Technology in Patients Undergoing Mechanical Circulatory Support Therapy With HeartMate 3 (MOMENTUM 3) Randomized Clinical Trial. *JAMA Cardiol*. Apr 01 2020; 5(4): 411-419. PMID 31939996
15. Knosalla C, Färber G, Rieth AJ, et al. Rationale and design of the randomized 'early ventricular assist device'-Trial (VAD-DZHK3). *ESC Heart Fail*. Oct 2025; 12(5): 3731-3740. PMID 40685553
16. Frazier OH, Gemmato C, Myers TJ, et al. Initial clinical experience with the HeartMate II axial-flow left ventricular assist device. *Tex Heart Inst J*. 2007; 34(3): 275-81. PMID 17948075
17. John R, Kamdar F, Liao K, et al. Improved survival and decreasing incidence of adverse events with the HeartMate II left ventricular assist device as bridge-to-transplant therapy. *Ann Thorac Surg*. Oct 2008; 86(4): 1227-34; discussion 1234-5. PMID 18805167

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

18. Miller LW, Pagani FD, Russell SD, et al. Use of a continuous-flow device in patients awaiting heart transplantation. *N Engl J Med*. Aug 30 2007; 357(9): 885-96. PMID 17761592
19. Patel ND, Weiss ES, Schaffer J, et al. Right heart dysfunction after left ventricular assist device implantation: a comparison of the pulsatile HeartMate I and axial-flow HeartMate II devices. *Ann Thorac Surg*. Sep 2008; 86(3): 832-40; discussion 832-40. PMID 18721570
20. Strüber M, Sander K, Lahpor J, et al. HeartMate II left ventricular assist device; early European experience. *Eur J Cardiothorac Surg*. Aug 2008; 34(2): 289-94. PMID 18571932
21. Kirklin JK, Naftel DC, Stevenson LW, et al. INTERMACS database for durable devices for circulatory support: first annual report. *J Heart Lung Transplant*. Oct 2008; 27(10): 1065-72. PMID 18926395
22. Aissaoui N, Morshuis M, Maoulida H, et al. Management of end-stage heart failure patients with or without ventricular assist device: an observational comparison of clinical and economic outcomes. *Eur J Cardiothorac Surg*. Jan 01 2018; 53(1): 170-177. PMID 28950304
23. Schmitto JD, Pya Y, Zimpfer D, et al. Long-term evaluation of a fully magnetically levitated circulatory support device for advanced heart failure-two-year results from the HeartMate 3 CE Mark Study. *Eur J Heart Fail*. Jan 2019; 21(1): 90-97. PMID 30052304
24. Gustafsson F, Shaw S, Lavee J, et al. Six-month outcomes after treatment of advanced heart failure with a full magnetically levitated continuous flow left ventricular assist device: report from the ELEVATE registry. *Eur Heart J*. Oct 01 2018; 39(37): 3454-3460. PMID 30165521
25. Pagani FD, Mehra MR, Cowger JA, et al. Clinical outcomes and healthcare expenditures in the real world with left ventricular assist devices - The CLEAR-LVAD study. *J Heart Lung Transplant*. May 2021; 40(5): 323-333. PMID 33744086
26. Mehra MR, Cleveland JC, Uriel N, et al. Primary results of long-term outcomes in the MOMENTUM 3 pivotal trial and continued access protocol study phase: a study of 2200 HeartMate 3 left ventricular assist device implants. *Eur J Heart Fail*. Aug 2021; 23(8): 1392-1400. PMID 33932272
27. Khoufi EAA. Outcomes of Left Ventricular Assist Devices as Destination Therapy: A Systematic Review with Meta-Analysis. *Life (Basel)*. Jan 03 2025; 15(1). PMID 39859993
28. Rose EA, Gelijns AC, Moskowitz AJ, et al. Long-term use of a left ventricular assist device for end-stage heart failure. *N Engl J Med*. Nov 15 2001; 345(20): 1435-43. PMID 11794191
29. Park SJ, Tector A, Piccioni W, et al. Left ventricular assist devices as destination therapy: a new look at survival. *J Thorac Cardiovasc Surg*. Jan 2005; 129(1): 9-17. PMID 15632819
30. Long JW, Kfoury AG, Slaughter MS, et al. Long-term destination therapy with the HeartMate XVE left ventricular assist device: improved outcomes since the REMATCH study. *Congest Heart Fail*. 2005; 11(3): 133-8. PMID 15947534
31. Rogers JG, Pagani FD, Tatroles AJ, et al. Intrapericardial Left Ventricular Assist Device for Advanced Heart Failure. *N Engl J Med*. Feb 02 2017; 376(5): 451-460. PMID 28146651
32. Estep JD, Starling RC, Horstmanshof DA, et al. Risk Assessment and Comparative Effectiveness of Left Ventricular Assist Device and Medical Management in Ambulatory Heart Failure Patients: Results From the ROADMAP Study. *J Am Coll Cardiol*. Oct 20 2015; 66(16): 1747-1761. PMID 26483097
33. Starling RC, Estep JD, Horstmanshof DA, et al. Risk Assessment and Comparative Effectiveness of Left Ventricular Assist Device and Medical Management in Ambulatory Heart Failure

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

- Patients: The ROADMAP Study 2-Year Results. JACC Heart Fail. Jul 2017; 5(7): 518-527. PMID 28396040
34. Jorde UP, Kushwaha SS, Tatroles AJ, et al. Results of the destination therapy post-food and drug administration approval study with a continuous flow left ventricular assist device: a prospective study using the INTERMACS registry (Interagency Registry for Mechanically Assisted Circulatory Support). J Am Coll Cardiol. May 06 2014; 63(17): 1751-7. PMID 24613333
 35. Arnold SV, Jones PG, Allen LA, et al. Frequency of Poor Outcome (Death or Poor Quality of Life) After Left Ventricular Assist Device for Destination Therapy: Results From the INTERMACS Registry. Circ Heart Fail. Aug 2016; 9(8). PMID 27507111
 36. Mehra MR, Goldstein DJ, Cleveland JC, et al. Five-Year Outcomes in Patients With Fully Magnetically Levitated vs Axial-Flow Left Ventricular Assist Devices in the MOMENTUM 3 Randomized Trial. JAMA. Sep 27 2022; 328(12): 1233-1242. PMID 36074476
 37. Wever-Pinzon O, Drakos SG, McKellar SH, et al. Cardiac Recovery During Long-Term Left Ventricular Assist Device Support. J Am Coll Cardiol. Oct 04 2016; 68(14): 1540-53. PMID 27687196
 38. Topkara VK, Garan AR, Fine B, et al. Myocardial Recovery in Patients Receiving Contemporary Left Ventricular Assist Devices: Results From the Interagency Registry for Mechanically Assisted Circulatory Support (INTERMACS). Circ Heart Fail. Jul 2016; 9(7). PMID 27402861
 39. Maybaum S, Mancini D, Xydas S, et al. Cardiac improvement during mechanical circulatory support: a prospective multicenter study of the LVAD Working Group. Circulation. May 15 2007; 115(19): 2497-505. PMID 17485581
 40. Agrawal S, Garg L, Shah M, et al. Thirty-Day Readmissions After Left Ventricular Assist Device Implantation in the United States: Insights From the Nationwide Readmissions Database. Circ Heart Fail. Mar 2018; 11(3): e004628. PMID 29519902
 41. Dickstein K, Cohen-Solal A, Filippatos G, et al. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2008: the Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2008 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association of the ESC (HFA) and endorsed by the European Society of Intensive Care Medicine (ESICM). Eur Heart J. Oct 2008; 29(19): 2388-442. PMID 18799522
 42. Bulic A, Maeda K, Zhang Y, et al. Functional status of United States children supported with a left ventricular assist device at heart transplantation. J Heart Lung Transplant. Aug 2017; 36(8): 890-896. PMID 28363739
 43. Wehman B, Stafford KA, Bittle GJ, et al. Modern Outcomes of Mechanical Circulatory Support as a Bridge to Pediatric Heart Transplantation. Ann Thorac Surg. Jun 2016; 101(6): 2321-7. PMID 26912304
 44. Fraser CD, Jaquiss RD, Rosenthal DN, et al. Prospective trial of a pediatric ventricular assist device. N Engl J Med. Aug 09 2012; 367(6): 532-41. PMID 22873533
 45. Blume ED, Rosenthal DN, Rossano JW, et al. Outcomes of children implanted with ventricular assist devices in the United States: First analysis of the Pediatric Interagency Registry for Mechanical Circulatory Support (PediMACS). J Heart Lung Transplant. May 2016; 35(5): 578-84. PMID 27009673

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

46. Almond CS, Morales DL, Blackstone EH, et al. Berlin Heart EXCOR pediatric ventricular assist device for bridge to heart transplantation in US children. *Circulation*. Apr 23 2013; 127(16): 1702-11. PMID 23538380
47. Jordan LC, Ichord RN, Reinhartz O, et al. Neurological complications and outcomes in the Berlin Heart EXCOR® pediatric investigational device exemption trial. *J Am Heart Assoc*. Jan 22 2015; 4(1): e001429. PMID 25613996
48. Chen S, Lin A, Liu E, et al. Outpatient Outcomes of Pediatric Patients with Left Ventricular Assist Devices. *ASAIO J*. 2016; 62(2): 163-8. PMID 26720740
49. Conway J, Al-Aklabi M, Granoski D, et al. Supporting pediatric patients with short-term continuous-flow devices. *J Heart Lung Transplant*. May 2016; 35(5): 603-9. PMID 27009672
50. Aaronson KD, Eppinger MJ, Dyke DB, et al. Left ventricular assist device therapy improves utilization of donor hearts. *J Am Coll Cardiol*. Apr 17 2002; 39(8): 1247-54. PMID 11955839
51. Frazier OH, Rose EA, McCarthy P, et al. Improved mortality and rehabilitation of transplant candidates treated with a long-term implantable left ventricular assist system. *Ann Surg*. Sep 1995; 222(3): 327-36; discussion 336-8. PMID 7677462
52. Bank AJ, Mir SH, Nguyen DQ, et al. Effects of left ventricular assist devices on outcomes in patients undergoing heart transplantation. *Ann Thorac Surg*. May 2000; 69(5): 1369-74; discussion 1375. PMID 10881807
53. Shuhaiber JH, Hur K, Gibbons R. The influence of preoperative use of ventricular assist devices on survival after heart transplantation: propensity score matched analysis. *BMJ*. Feb 10 2010; 340: c392. PMID 20147346
54. Alba AC, McDonald M, Rao V, et al. The effect of ventricular assist devices on long-term post-transplant outcomes: a systematic review of observational studies. *Eur J Heart Fail*. Jul 2011; 13(7): 785-95. PMID 21551162
55. Deo SV, Sung K, Daly RC, et al. Cardiac transplantation after bridged therapy with continuous flow left ventricular assist devices. *Heart Lung Circ*. Mar 2014; 23(3): 224-8. PMID 23954004
56. Grimm JC, Sciortino CM, Magruder JT, et al. Outcomes in Patients Bridged With Univentricular and Biventricular Devices in the Modern Era of Heart Transplantation. *Ann Thorac Surg*. Jul 2016; 102(1): 102-8. PMID 27068177
57. Davies RR, Russo MJ, Hong KN, et al. The use of mechanical circulatory support as a bridge to transplantation in pediatric patients: an analysis of the United Network for Organ Sharing database. *J Thorac Cardiovasc Surg*. Feb 2008; 135(2): 421-7, 427.e1. PMID 18242279
58. Copeland JG, Smith RG, Arabia FA, et al. Cardiac replacement with a total artificial heart as a bridge to transplantation. *N Engl J Med*. Aug 26 2004; 351(9): 859-67. PMID 15329423
59. Copeland JG, Copeland H, Gustafson M, et al. Experience with more than 100 total artificial heart implants. *J Thorac Cardiovasc Surg*. Mar 2012; 143(3): 727-34. PMID 22245242
60. Food and Drug Administration. Summary of Safety and Probable Benefit - H040006: AbioCor Implantable Replacement Heart. 2006; https://www.accessdata.fda.gov/cdrh_docs/pdf4/H040006b.pdf.
61. Dowling RD, Gray LA, Etoch SW, et al. Initial experience with the AbioCor implantable replacement heart system. *J Thorac Cardiovasc Surg*. Jan 2004; 127(1): 131-41. PMID 14752423
62. Torregrossa G, Morshuis M, Varghese R, et al. Results with SynCardia total artificial heart beyond 1 year. *ASAIO J*. 2014; 60(6): 626-34. PMID 25158888

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

63. Tariq MD, Jain H, Khan AM, et al. Efficacy and safety of percutaneous mechanical circulatory support in patients with cardiogenic shock following acute myocardial infarction: A meta-analysis of randomized controlled trials. *Medicine (Baltimore)*. Nov 15 2024; 103(46): e40595. PMID 39560531
64. Ouweneel DM, Eriksen E, Sjauw KD, et al. Percutaneous Mechanical Circulatory Support Versus Intra-Aortic Balloon Pump in Cardiogenic Shock After Acute Myocardial Infarction. *J Am Coll Cardiol*. Jan 24 2017; 69(3): 278-287. PMID 27810347
65. Seyfarth M, Sibbing D, Bauer I, et al. A randomized clinical trial to evaluate the safety and efficacy of a percutaneous left ventricular assist device versus intra-aortic balloon pumping for treatment of cardiogenic shock caused by myocardial infarction. *J Am Coll Cardiol*. Nov 04 2008; 52(19): 1584-8. PMID 19007597
66. Bochaton T, Huot L, Elbaz M, et al. Mechanical circulatory support with the Impella® LP5.0 pump and an intra-aortic balloon pump for cardiogenic shock in acute myocardial infarction: The IMPELLA-STIC randomized study. *Arch Cardiovasc Dis*. Apr 2020; 113(4): 237-243. PMID 31740272
67. Møller JE, Engstrøm T, Jensen LO, et al. Microaxial Flow Pump or Standard Care in Infarct-Related Cardiogenic Shock. *N Engl J Med*. Apr 18 2024; 390(15): 1382-1393. PMID 38587239
68. Karami M, Eriksen E, Ouweneel DM, et al. Long-term 5-year outcome of the randomized IMPRESS in severe shock trial: percutaneous mechanical circulatory support vs. intra-aortic balloon pump in cardiogenic shock after acute myocardial infarction. *Eur Heart J Acute Cardiovasc Care*. Dec 06 2021; 10(9): 1009-1015. PMID 34327527
69. Ait Ichou J, Larivée N, Eisenberg MJ, et al. The effectiveness and safety of the Impella ventricular assist device for high-risk percutaneous coronary interventions: A systematic review. *Catheter Cardiovasc Interv*. Jun 2018; 91(7): 1250-1260. PMID 28941078
70. Iannaccone M, Barbero U, Franchin L, et al. Comparison of mid-term mortality after surgical, supported or unsupported percutaneous revascularization in patients with severely reduced ejection fraction: A direct and network meta-analysis of adjusted observational studies and randomized-controlled. *Int J Cardiol*. Feb 01 2024; 396: 131428. PMID 37820779
71. Briasoulis A, Telila T, Palla M, et al. Meta-Analysis of Usefulness of Percutaneous Left Ventricular Assist Devices for High-Risk Percutaneous Coronary Interventions. *Am J Cardiol*. Aug 01 2016; 118(3): 369-75. PMID 27265673
72. O'Neill WW, Kleiman NS, Moses J, et al. A prospective, randomized clinical trial of hemodynamic support with Impella 2.5 versus intra-aortic balloon pump in patients undergoing high-risk percutaneous coronary intervention: the PROTECT II study. *Circulation*. Oct 02 2012; 126(14): 1717-27. PMID 22935569
73. Ouweneel DM, Engstrom AE, Sjauw KD, et al. Experience from a randomized controlled trial with Impella 2.5 versus IABP in STEMI patients with cardiogenic pre-shock. Lessons learned from the IMPRESS in STEMI trial. *Int J Cardiol*. Jan 01 2016; 202: 894-6. PMID 26476989
74. Reddy YM, Chinitz L, Mansour M, et al. Percutaneous left ventricular assist devices in ventricular tachycardia ablation: multicenter experience. *Circ Arrhythm Electrophysiol*. Apr 2014; 7(2): 244-50. PMID 24532564
75. Aryana A, Gearoid O'Neill P, Gregory D, et al. Procedural and clinical outcomes after catheter ablation of unstable ventricular tachycardia supported by a percutaneous left ventricular assist device. *Heart Rhythm*. Jul 2014; 11(7): 1122-30. PMID 24732372

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

76. Kar B, Gregoric ID, Basra SS, et al. The percutaneous ventricular assist device in severe refractory cardiogenic shock. *J Am Coll Cardiol*. Feb 08 2011; 57(6): 688-96. PMID 20950980
77. Kirklin JK, Pagani FD, Goldstein DJ, et al. American Association for Thoracic Surgery/International Society for Heart and Lung Transplantation guidelines on selected topics in mechanical circulatory support. *J Heart Lung Transplant*. Mar 2020; 39(3): 187-219. PMID 31983666
78. Yancy CW, Jessup M, Bozkurt B, et al. 2017 ACC/AHA/HFSA Focused Update of the 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *Circulation*. Aug 08 2017; 136(6): e137-e161. PMID 28455343
79. Yancy CW, Jessup M, Bozkurt B, et al. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. Oct 15 2013; 62(16): e147-239. PMID 23747642
80. Heidenreich PA, Bozkurt B, Aguilar D, et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: Executive Summary: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. May 03 2022; 145(18): e876-e894. PMID 35363500
81. Peura JL, Colvin-Adams M, Francis GS, et al. Recommendations for the use of mechanical circulatory support: device strategies and patient selection: a scientific statement from the American Heart Association. *Circulation*. Nov 27 2012; 126(22): 2648-67. PMID 23109468
82. Bernhardt AM, Copeland H, Deswal A, et al. The International Society for Heart and Lung Transplantation/Heart Failure Society of America Guideline on Acute Mechanical Circulatory Support. *J Heart Lung Transplant*. Apr 2023; 42(4): e1-e64. PMID 36805198
83. Rihal CS, Naidu SS, Givertz MM, et al. 2015 SCAI/ACC/HFSA/STS Clinical Expert Consensus Statement on the Use of Percutaneous Mechanical Circulatory Support Devices in Cardiovascular Care: Endorsed by the American Heart Association, the Cardiological Society of India, and Sociedad Latino Americana de Cardiologia Intervencion; Affirmation of Value by the Canadian Association of Interventional Cardiology-Association Canadienne de Cardiologie d'intervention. *J Am Coll Cardiol*. May 19 2015; 65(19): e7-e26. PMID 25861963
84. Centers for Medicare & Medicaid Services. National Coverage Determination (NCD) for Ventricular Assist Devices (20.9.1). 2020; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?ncdid=360&ncdver=2&keyword=ventricular%20assist&keywordType=starts&areald=all&docType=NCD&contractOption=all&sortBy=relevance&bc=AAAAAAQAAAA&KeyWordLookUp=Doc&KeyWordSearchType=Exact>.
85. Jacob E.Moller et al. *The New England Journal of Medicine*. Microaxial Flow Pump or Standard Care in Infarct-Related Cardiogenic Shock. 2024;390:1382-1393.
86. Dharam J. Kumbhani, MD, SM, FACC. American College of Cardiology. Microaxial Flow Pump in Infarct-Related Cardiogenic Shock - DanGer Shock. December 23, 2024.
87. Bochaton T, Huot L, Elbaz M, et al. Mechanical circulatory support with the Impella LP5.0 pump and an intra-aortic balloon pump for cardiogenic shock in acute myocardial infarction: the IMPELLA-STIC randomized study. *Arch Cardiovasc Dis* 2020;113:237-243.

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

88. Ouweneel DM, Eriksen E, Sjauw KD, et al. Percutaneous mechanical circulatory support versus intra-aortic balloon pump in cardiogenic shock after acute myocardial infarction. *J Am Coll Cardiol* 2017;69:278-287.
89. Seyfarth M, Sibbing D, Bauer I, et al. A randomized clinical trial to evaluate the safety and efficacy of a percutaneous left ventricular assist device versus intra-aortic balloon pumping for treatment of cardiogenic shock caused by myocardial infarction. *J Am Coll Cardiol* 2008;52:1584-1588.
90. Bogerd M, Ten Berg S, Peters EJ, et al. Impella and venoarterial extracorporeal membrane oxygenation in cardiogenic shock complicating acute myocardial infarction. *Eur J Heart Fail* 2023;25:2021-2031.
91. Dhruva SS, Ross JS, Mortazavi BJ, et al. Use of mechanical circulatory support devices among patients with acute myocardial infarction complicated by cardiogenic shock. *JAMA Netw Open* 2021;4(2):e2037748-e2037748.
92. Schrage B, Ibrahim K, Loehn T, et al. Impella support for acute myocardial infarction complicated by cardiogenic shock. *Circulation* 2019;139:1249-1258.
93. Holgerthiele, M.D. et. al. Extracorporeal Life Support in Infarct-Related Cardiogenic Shock. *The New England Journal of Medicine*. VOL 389 NO.14. Published August 26, 2023.
94. William W. O'Neill M.D. et al. A Prospective, Randomized Clinical Trial of Hemodynamic Support With Impella 2.5 Versus Intra-Aortic Balloon Pump in Patients Undergoing High-Risk Percutaneous Coronary Intervention. *Circulation* Volume 126, Issue 14, 2 October 2012; Pages 1717-1727.
95. Jason Woolmuth et al. Ejection Fraction Improvement Following Contemporary High-Risk Percutaneous Coronary Intervention: RESTORE EF Study Results. *J Soc Cardiovasc Angiogr Interv*. 2022 Aug 13;1(5):100350. doi: 10.1016/j.jscv.2022.100350
96. <https://clinicaltrials.gov/study/NCT04763200><https://clinicaltrials.gov/study/NCT04763200>
97. Jonathan Gabriel Sung, Use of Mechanical Circulatory Support in Complex High-risk Percutaneous Coronary Intervention: A Review of Current Evidence. JAPSC. *Journal of Asian Pacific Society of Cardiology*.
98. Sunil V. Rao MD et al. 2025 ACC/AHA/ACEP/NAEMSP/SCAI Guideline for the Management of Patients With Acute Coronary Syndromes: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. Published February 27, 2025.
99. Thiele H, Lauer B, Hambrecht R, et al. Short and long-term hemodynamic effects of intra-aortic balloon support in ventricular septal defect complicating acute myocardial infarction. *Am J Cardiol*. 2003;92:450–454.

Policy History

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

01/07/2010 Medical Policy Committee approval

01/20/2010 Medical Policy Implementation Committee approval. New policy.

01/06/2011 Medical Policy Committee approval

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

01/19/2011	Medical Policy Implementation Committee approval. Title changed. Policy statements revised to address only implantable VADs and total artificial hearts.
04/12/2012	Medical Policy Committee approval
04/25/2012	Medical Policy Implementation Committee approval. Percutaneous VADs added to policy investigational statement and rationale.
04/04/2013	Medical Policy Committee review
04/24/2013	Medical Policy Implementation Committee approval. Added "Implantable" to the beginning of the 2nd coverage statement under Bridge to Transplant to make it consistent with the other coverage statements and the focus of the policy. Coverage statement on children amended; age range changed from 5-16 to 0-16, reflecting the approval of the BERLIN heart EXCOR device for pediatric patients aged 0-16. Clause added to coverage statement on total artificial hearts that says "...or are undergoing evaluation to determine candidacy for heart transplantation...".
08/07/2014	Medical Policy Committee review
08/20/2014	Medical Policy Implementation Committee approval. Coverage statement unchanged.
08/03/2015	Coding update: ICD10 Diagnosis code section added; ICD9 Procedure code section removed.
10/29/2015	Medical Policy Committee review
11/16/2015	Medical Policy Implementation Committee approval. Coverage statement unchanged.
11/03/2016	Medical Policy Committee review
11/16/2016	Medical Policy Implementation Committee approval. No change to coverage.
01/01/2017	Coding update: Removing ICD-9 Diagnosis Codes
11/02/2017	Medical Policy Committee review
11/15/2017	Medical Policy Implementation Committee approval. No change to coverage. Added new FDA information.
01/01/2018	Coding update
11/08/2018	Medical Policy Committee review
11/21/2018	Medical Policy Implementation Committee approval. No change to coverage.
11/07/2019	Medical Policy Committee review
11/13/2019	Medical Policy Implementation Committee approval. No change to coverage.
11/05/2020	Medical Policy Committee review
11/11/2020	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
12/11/2020	Coding update
11/04/2021	Medical Policy Committee review
11/10/2021	Medical Policy Implementation Committee approval. Policy statements revised to remove outdated eligibility criteria, but intent unchanged.
11/03/2022	Medical Policy Committee review
11/09/2022	Medical Policy Implementation Committee approval. No change to coverage.
11/02/2023	Medical Policy Committee review
11/08/2023	Medical Policy Implementation Committee approval. Minor changes to coverage. Intent unchanged.

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

11/07/2024 Medical Policy Committee review

11/13/2024 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

04/03/2025 Medical Policy Committee review

04/09/2025 Medical Policy Implementation Committee approval. Coverage for percutaneous ventricular assist devices (PVADs) changed from investigational to eligible for coverage with criteria.

04/02/2026 Medical Policy Committee review

04/10/2026 Medical Policy Implementation Committee review.

04/16/2026 Medical Quality Management Committee approval. Coverage for implantable VADs as a bridge to heart transplantation broken out into adults and children separately. Also added “is not expected to survive until a donor heart can be obtained” to this criteria.

Added the following criteria to total artificial hearts (TAHs) as a bridge to heart transplantation coverage:

- Absence of any of the contraindications:
 - o Chronic irreversible hepatic, renal, or respiratory failure
 - o Active systemic infection
 - o Coagulation disorders and inability to tolerate anticoagulation therapy.

Added “left heart pVAD is FDA approved” to the percutaneous ventricular assist devices (pVADs) for the treatment of cardiogenic shock criteria. Added “in accordance with FDA approval” to the ventricular assist devices criteria statements. Added Not Medically Necessary section. Added “Based on review of available data, the Company considers the use of percutaneous VADs when criteria are not met to be not medically necessary.” Added “Based on review of available data, the Company considers other applications of implantable ventricular assist devices (VADs) or total artificial hearts (TAHs) including, but not limited to, the use of TAHs as destination therapy to be not medically necessary.” Removed “Other Indications Based on review of available data, the Company considers other applications of implantable ventricular assist devices (VADs) or total artificial hearts (TAHs) including, but not limited to, the use of TAHs as destination therapy to be investigational.” Added “Based on review of available data, the Company considers the use of right heart percutaneous VADs to be investigational.”

Next Scheduled Review Date: 04/2027

Coding

The five character codes included in the Louisiana Blue Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)†, copyright 2025 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

The responsibility for the content of Louisiana Blue Medical Policy Coverage Guidelines is with Louisiana Blue and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Louisiana Blue Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Louisiana Blue Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	33927, 33928, 33929, 33975, 33976, 33977, 33978, 33979, 33980, 33981, 33982, 33983, 33990, 33991, 33992, 33993, 33995, 33997, 93750
HCPCS	L8698, Q0477, Q0478, Q0479, Q0480, Q0481, Q0482, Q0483, Q0484, Q0485, Q0486, Q0487, Q0488, Q0489, Q0490, Q0491, Q0492, Q0493, Q0494, Q0495, Q0496, Q0497, Q0498, Q0499, Q0500, Q0501, Q0502, Q0503, Q0504, Q0506, Q0507, Q0508, Q0509
ICD-10 Diagnosis	I21.01 -I21.02, I21.11-I21.19, I21.21-I21.29, I21.3, I21.4, I21.9, I22.0-I22.9, R57.0, T81.11XA-T81.11XS

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
 1. Consultation with technology evaluation center(s);
 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
 3. Reference to federal regulations.

Total Artificial Hearts and Implantable Ventricular Assist Devices

Policy # 00246

Original Effective Date: 01/20/2010

Current Effective Date: 07/01/2026

****Medically Necessary** (or “Medical Necessity”) - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

For these purposes, “nationally accepted standards of medical practice” means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: If the Patient’s health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

NOTICE: Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage.

NOTICE: If an authorization for an ongoing course of treatment has been provided to a member and the member changes from one health plan to another health plan (e.g., a member moves from carrier A to Louisiana Blue), Louisiana Blue may honor the previous health plan’s authorization for the same service under the same type of in-network benefit for a 90-day transition period. Documentation of the authorization for the ongoing course of treatment from the previous health plan must be provided to us by the member or their provider and the services provided for the course of treatment must otherwise be a covered service under the Louisiana Blue health plan. This provision does not apply to medications covered under the plan’s pharmacy benefit.