

evinacumab-dgnb (Evkeeza™)

Policy # 00751

Original Effective Date: 08/09/2021

Current Effective Date: 08/01/2026

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the “Company”), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Investigational or experimental services are not covered. This includes any drug, device, procedure, or service provided under the investigational arm of a clinical trial or clinical study. These services are excluded from coverage under benefits.

When Services May Be Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member’s contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of available data, the Company may consider evinacumab-dgnb (Evkeeza™)† for the treatment of homozygous familial hypercholesterolemia (HoFH) to be **eligible for coverage**.**

Patient Selection Criteria

Coverage eligibility for evinacumab-dgnb (Evkeeza) will be considered when the following criteria are met:

Initial Authorization: (Patient must meet I, II, III, IV, V, **and** VI)

- I. Patient is greater than or equal to 1 year of age; AND
- II. Patient is adherent to any medications required for therapy prior to receiving authorization for evinacumab-dgnb (Evkeeza); AND
*(Note: This specific patient selection criterion is an additional company requirement for coverage eligibility and will be denied as not medically necessary** if not met.)*
- III. evinacumab-dgnb (Evkeeza) will be used along with a maximally tolerated statin [in those who are not considered statin intolerant (see below for statin intolerance)] AND a maximally dosed proprotein convertase subtilisin kexin type 9 (PCSK9) inhibitor, such as Repatha®‡ (evolocumab) or Praluent®‡ (alirocumab), unless the patient has two LDL-receptor negative alleles, AND generic ezetimibe 10 mg daily; AND
- IV. Patient meets one of the following criteria which confirms the diagnosis of HoFH (Patient must meet A, B, **or** C):
 - A. Patient has had genetic confirmation of two mutant alleles at the low-density lipoprotein receptor (LDLR), apolipoprotein B (apo B), proprotein convertase subtilisin kexin type 9 (PCSK9), or low-density lipoprotein receptor adaptor protein 1 (LDLRAP1) gene locus; OR
 - B. Patient has an untreated (i.e. prior to therapy with any anti-hyperlipidemic agent) low-density lipoprotein cholesterol (LDL-C) level greater than 400 mg/dL AND meets one of the following criteria (i **or** ii):
(Note: Untreated refers to prior to therapy with any antihyperlipidemic agent.)

Policy # 00751

Original Effective Date: 08/09/2021

Current Effective Date: 08/01/2026

- i. Patient had clinical manifestations of HoFH (i.e. cutaneous xanthomas, tendon xanthomas, arcus cornea, tuberous xanthomas, or xanthelasma) before the age of 10 years; OR
 - ii. Both parents of the patient had untreated (i.e. prior to therapy with any anti-hyperlipidemic agent) LDL-C levels or total cholesterol levels consistent with heterozygous familial hypercholesterolemia (HeFH); OR
(Note: An example of HeFH in both parents would be if both had an untreated LDL-C level greater than or equal to 190 mg/dL and/or an untreated total cholesterol level greater than 250 mg/dL.)
(Note: In digenic form, one parent may have normal LDL-C levels and the other may have LDL-C levels consistent with HoFH)
 - C. Patient has a treated (i.e. therapy with at least one anti-hyperlipidemic agent) LDL-C level greater than or equal to 300 mg/dL AND meets one of the following criteria (i or ii):
(Note: Treated refers to after therapy with at least one antihyperlipidemic agent. Some examples of antihyperlipidemic agents include statins [e.g., atorvastatin, rosuvastatin, lovastatin, simvastatin, pravastatin], ezetimibe, a PCSK9 inhibitor [e.g., Praluent (alirocumab subcutaneous injection)], or Juxtapid® [lomitapide capsules])
 - i. Patient had clinical manifestations of HoFH (i.e. cutaneous xanthomas, tendon xanthomas, arcus cornea, tuberous xanthomas, or xanthelasma) before the age of 10 years; OR
 - ii. Both parents of the patient had untreated (i.e. prior to therapy with any anti-hyperlipidemic agent) LDL-C levels or total cholesterol levels consistent with heterozygous familial hypercholesterolemia (HeFH); AND
(Note: An example of HeFH in both parents would be if both had an untreated LDL-C greater than or equal to 190 mg/dL and/or an untreated total cholesterol greater than 250 mg/dL.)
- V. Patient meets one of the following criteria (A or B):
 - A. Patient meets ALL of the following criteria (i, ii, and iii)
 - i. Patient has tried one high-intensity maximum daily dose statin therapy (i.e., generic atorvastatin greater than or equal to 80 mg daily; generic rosuvastatin greater than or equal to 40 mg daily [as a single-entity or as a combination product]) for greater than or equal to 12 continuous weeks OR patient has tried a maximally tolerated stable daily dose statin of any potency for greater than or equal to 12 continuous weeks if proof is given that a high potency maximum daily dose statin was not well tolerated [as a single-entity or as a combination product]); AND
*(Note that the 12-week timeframe is an additional company requirement and will be denied as not medically necessary** if not met)*
 - ii. Patient has taken the statin along with generic ezetimibe 10 mg daily (as a single entity or as a combination product) for greater than or equal to 12 continuous weeks, unless the patient is statin intolerant (defined below), then generic ezetimibe 10 mg daily without a statin should be taken for greater than or equal to 12 continuous weeks; AND

Policy # 00751

Original Effective Date: 08/09/2021

Current Effective Date: 08/01/2026

*(Note that the 12-week timeframe is an additional company requirement and will be denied as not medically necessary** if not met)*

- iii. Patient's LDL-C level after this treatment regimen remains greater than or equal to 70 mg/dL or greater than or equal to 55 mg/dL based on "high risk" or "very high risk" classifications of the patient; OR
- B. Patient has been determined to be statin intolerant by meeting the following criteria (i **and** ii):
 - i. Patient's LDL-C level remains greater than or equal to 70 mg/dL or greater than or equal to 55 mg/dL based on "high risk" or "very high risk" classifications of the patient due to statin intolerance; AND
 - ii. Patient meets all of the following criteria [(1), (2), **and** (3)]:
 - 1. Patient was unable to tolerate at least 2 different statins. [The inability to tolerate one statin will be accepted if the patient experienced rhabdomyolysis or clinically significant myonecrosis secondary to a statin. Rhabdomyolysis/myonecrosis is considered to be a muscle breakdown with signs and symptoms such as muscle pain, weakness, tenderness **ALONG WITH** either: a.) acute renal failure or myoglobinuria **AND** elevated creatine kinase levels (i.e., greater than or equal to 10 times the upper limit of normal) **OR** b.) elevated creatine kinase levels (i.e., greater than or equal to 10 times the upper limit of normal) alone]; AND
 - 2. Patient's intolerance was associated with confirmed, intolerable statin-related adverse effects [i.e., skeletal related muscle symptoms (myopathy [muscle weakness] or myalgia [muscle aches, soreness, stiffness, or tenderness])]; AND
 - 3. Patient's symptoms or biomarker changes resolved or showed significant improvement on dose decrease or discontinuation; AND
- VI. Patient meets **ONE** of the following criteria (A, B, **or** C):
 - A. Patient meets **BOTH** of the following criteria (i **and** ii):
 - i. Patient has tried a PCSK9 inhibitor for greater than or equal to 12 continuous weeks at the FDA approved maximum dose for HoFH; AND
(Note: Examples of PCSK9 inhibitors at their maximum dose include Repatha (evolocumab) 420 mg once every 2 weeks and Praluent (alirocumab) 150 mg once every 2 weeks)
*(Note that the 12-week timeframe is an additional company requirement and will be denied as not medically necessary** if not met)*
 - ii. Patient's LDL-C level after the max PCSK9 inhibitor HoFH dosing regimen remains greater than or equal to 70 mg/dL or greater than or equal to 55 mg/dL based on "high risk" or "very high risk" classifications of the patient; OR
 - B. Patient is known to have two LDL-receptor negative alleles; OR
 - C. Patient is 1 to 9 years of age.

evinacumab-dgnb (Evkeeza™)

Policy # 00751

Original Effective Date: 08/09/2021

Current Effective Date: 08/01/2026

Re-authorization: (Patient must meet I **and** II)

- I. Patient previously met the initial criteria and received an approval for evinacumab-dgnb (Evkeeza); AND
- II. Patient has achieved clinically significant LDL-C lowering AND is adherent to evinacumab-dgnb (Evkeeza)
*(Note that the re-authorization criteria are additional company requirements and will be denied as not medically necessary** if not met)*

When Services Are Considered Not Medically Necessary

Based on review of available data, the Company considers the use of evinacumab-dgnb (Evkeeza) when the member has not tried the required pre-requisite medications for a timeframe of at least 12 weeks to be **not medically necessary.****

Based on review of available data, the Company considers the use of evinacumab-dgnb (Evkeeza) when the re-authorization criteria are NOT met to be **not medically necessary.****

Based on review of available data, the Company considers the use of evinacumab-dgnb (Evkeeza) when the patient is not compliant with their cholesterol lowering regimen to be **not medically necessary.****

When Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers the use of evinacumab-dgnb (Evkeeza) when patient selection criteria are not met (except those listed above as **not medically necessary****) to be **investigational.***

Policy Guidelines

The National Lipid Association sets LDL-C treatment goal levels for patients at various risk stratifications. Those with clinical atherosclerotic cardiovascular disease would fall into the “very high risk” or “high risk” category and would therefore be treated to an LDL-C of less than 55 mg/dL or less than 70 mg/dL depending on patient specific factors. Patients with familial hypercholesterolemia could fall into either the “very high risk” or “high risk” categories, based on their patient characteristics and would therefore have a treatment goal of less than 55 mg/dL or 70 mg/dL (respectively). Risk stratification (per the National Lipid Association) is as follows:

Policy # 00751

Original Effective Date: 08/09/2021

Current Effective Date: 08/01/2026

Very High-Risk Individuals	
Major ASCVD Events ☐ ACS (< last 12 months) ☐ MI history (other than ACS above)	☐ Ischemic stroke history ☐ Symptomatic PAD*
High-Risk Conditions ☐ Age ≥ 65 years ☐ Diabetes mellitus ☐ Hypertension ☐ Smoker (current) ☐ LDL-C ≥ 100 mg/dL despite maximally tolerated statin & ezetimibe	
☐ CHF history ☐ Heterozygous FH (LDL-C ≥ 190 mg/dL) ☐ CKD (eGFR 15-59 mL/min/1.73m ²) ☐ Prior coronary artery bypass surgery^ ☐ Prior percutaneous coronary intervention^	
Risk Assessment 2+ Major ASCVD Events or 1 Major ASCVD Event & 2+ High-Risk Conditions	
LDL-C Objective < 55 mg/dL	
*History of claudication with ABI <0.85 or previous revascularization or amputation ^Outside of the major ASCVD event(s)	
High-Risk Individuals	
Patients with 1+ of the Following: ☐ Type 2 diabetes mellitus ☐ Familial Hypercholesterolemia, without a prior event	
☐ CAC score > 100 AU or ≥ 75th percentile of the CAC score distribution for a particular age and sex ☐ Clinical ASCVD without other high-risk features ☐ Primary prevention patient with an estimated 10-year event risk of ≥ 20% in a pooled cohort risk calculation	
LDL-C Objective < 70 mg/dL	
Patients with 1+ of the Following: ☐ No clinical ASCVD and baseline LDL-C ≥ 190 mg/dL ☐ Calculated 10-year ASCVD risk of 7.5-19.9% by AHA/ACC pooled cohort equation	
LDL-C Objective < 100 mg/dL	
Optimal LDL-C for healthy adults < 100 mg/dL	

The above chart comes from National Lipid Association 2025. Available at: [https://www.lipid.org/sites/default/files/files/LDL%20Management%20Simplified%20\(6_30_2020_5%20v2\).pdf](https://www.lipid.org/sites/default/files/files/LDL%20Management%20Simplified%20(6_30_2020_5%20v2).pdf)

Clinical ASCVD (includes one or more of the following):

- I. Myocardial infarction or other acute coronary syndrome
- II. Angina (stable or unstable)
- III. Coronary or other revascularization procedure
- IV. Transient ischemic attack
- V. Ischemic stroke
- VI. Atherosclerotic peripheral arterial disease (ABI of less than 0.90)
- VII. Other documented atherosclerotic diseases such as
 - A. Clinically significant coronary atherosclerosis diagnosed by coronary angiography, stress test, stress echocardiography, or myocardial perfusion imaging
 - B. Renal atherosclerosis
 - C. Aortic aneurysm secondary to atherosclerosis
 - D. Carotid plaque (greater than or equal to 50% stenosis)

Policy # 00751

Original Effective Date: 08/09/2021

Current Effective Date: 08/01/2026

Background/Overview

Evkeeza is an ANGPTL3 (angiopoietin-like 3) inhibitor indicated as an adjunct to other low-density lipoprotein-cholesterol (LDL-C) lowering therapies for the treatment of adult and pediatric patients, aged 1 year and older, with homozygous familial hypercholesterolemia (HoFH). ANGPTL3 is a member of the angiopoietin-like protein family that is expressed primarily in the liver and plays a role in the regulation of lipid metabolism by inhibiting lipoprotein lipase (LPL) and endothelial lipase (EL). Evkeeza's inhibition of ANGPTL3 leads to reduction in LDL-C, HDL-C, and triglycerides (TG). Evkeeza reduces LDL-C independent of the presence of LDL receptor (LDLR) by promoting very low-density lipoprotein (VLDL) processing and clearance upstream of LDL formation. Evkeeza's blockade of ANGPTL3 lowers TG and HDL-C by rescuing LPL and EL activities, respectively. Evkeeza is supplied in 345 mg and 1,200 mg single-dose vials and is dosed at 15 mg/kg administered intravenously (IV) once monthly over 60 minutes. It should be noted that the safety and efficacy of Evkeeza have not been established in patients with other causes of hypercholesterolemia, including those with heterozygous familial hypercholesterolemia (HeFH). The effects of Evkeeza on cardiovascular morbidity and mortality have not been determined.

Hypercholesterolemia/Treatment Guidelines

Approximately 30% of the United States population has elevated LDL-C (low density lipoprotein cholesterol). There is also a subset of hypercholesterolemia, known as familial hypercholesterolemia, which can affect nearly 1 in 300 individuals. Familial hypercholesterolemia can further be broken down into homozygous and heterozygous forms of familial hypercholesterolemia. The homozygous form is by far the rarest with an estimated incidence of 1 in 1,000,000 individuals. HoFH is most commonly due to impaired functionality of the LDL receptor, which leads to a low or absence of clearance of LDL-C from the circulation. Genetic testing is available to determine whether or not an individual has familial hypercholesterolemia, however clinical signs/symptoms are often a more practical method of diagnosing this condition. Known mutations causing HoFH are located in the LDL receptor, apolipoprotein B (apo B), or proprotein convertase subtilisin kexin type 9 (PCSK9) genes. Patients with HoFH may have physical findings including tendon or cutaneous xanthomas coupled with elevated LDL-C levels (> 500 mg/dL [untreated] and > 300 mg/dL [treated]). The gold standard for the treatment of elevated LDL-C levels is a statin given along with ezetimibe (Zetia) to provide the greatest amount of LDL-C lowering. Statin products also have proven cardiovascular outcomes. Next steps in therapy for HoFH patients would include the PCSK9 inhibitors, Praluent and Repatha, which also have cardiovascular outcomes.

Statin Intolerance

Statins have been associated with muscle-related adverse effects such as myalgia (e.g., muscle aches, soreness, stiffness, or tenderness), myopathy (muscle weakness), and/or myositis (muscle inflammation). Although the incidence is variable, muscle adverse effects are reported in around 5% of patients receiving statins, but may be due to other causes (e.g., excessive exercise, other medical conditions [hypothyroidism], non-statin medications). It is advisable to assess for drug interactions as well as to check vitamin D levels and thyroid function status. Rhabdomyolysis, which is uncommon with statin therapy, is a severe muscle-related adverse effect that results in muscle breakdown associated with muscle-related symptoms (e.g., muscle pain, weakness, tenderness) along with acute renal failure and elevated creatine kinase [CK] levels (myonecrosis). In patients

Policy # 00751

Original Effective Date: 08/09/2021

Current Effective Date: 08/01/2026

with statin-related muscle adverse events, symptoms may not re-occur if the patient switches to a different statin therapy. Pravastatin and fluvastatin appear to have much less intrinsic muscle toxicity than other statins and could be considered for those who had statin related intolerable muscle symptoms.

In 2014, the NLA Statin Intolerance Panel published an update. It was stated that most statin intolerance is due to myalgia. The strongest evidence at present for statin intolerance in a population is that myalgia appears but then remits with withdrawal but reoccurs with re-challenge. The incidence of statin intolerance is widely variable. The Panel states that statins are among the safest medications available. The Panel does advise that due to statin benefits, it is safe to recommend a patient continue statin therapy even when some degree of statin intolerance is present, if the patient can reasonably tolerate the statin. A pivotal trial with Praluent called ODYSSEY ALTERNATIVE defined statin intolerance as the inability to take at least two different statins due to muscle-related adverse effects, of which one statin was administered at the lowest approved starting dose. Data also suggests that many patients who are re-challenged with statin therapy after an adverse event may be able to tolerate statin therapy long-term. Of note, in the ODYSSEY ALTERNATIVE trial with Praluent, 69.8% of patients who were considered statin intolerant were treated with atorvastatin 20 mg daily and completed the double-blind 24-week portion of the trial. This suggests that re-challenge with a statin in those purported to be statin intolerant is reasonable and may lead to successful use of a statin therapy.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

Evkeeza is indicated as an adjunct to other low-density lipoprotein-cholesterol (LDL-C) lowering therapies for the treatment of adult and pediatric patients, aged 5 years and older, with homozygous familial hypercholesterolemia. It should be noted that the safety and efficacy of Evkeeza have not been established in patients with other causes of hypercholesterolemia, including those with heterozygous familial hypercholesterolemia (HeFH). The effects of Evkeeza on cardiovascular morbidity and mortality have not been determined. In September 2025, the Evkeeza label was updated to include approval for patients aged 1 year and older.

Rationale/Source

This medical policy was developed through consideration of peer-reviewed medical literature generally recognized by the relevant medical community, U.S. Food and Drug Administration approval status, nationally accepted standards of medical practice and accepted standards of medical practice in this community, technology evaluation centers, reference to regulations, other plan medical policies, and accredited national guidelines.

ELIPSE-HoFH was a multicenter, double-blind, randomized, placebo-controlled trial evaluating the efficacy and safety of Evkeeza compared to placebo in 65 patients with HoFH. During the 24-week, double-blind treatment period, 43 patients were randomized to receive Evkeeza 15 mg/kg IV every 4 weeks and 22 patients to receive placebo. After the double-blind treatment period, 64 of 65 patients entered a 24-week open-label extension period in which all patients received Evkeeza 15 mg/kg IV every 4 weeks.

evinacumab-dgnb (Evkeeza™)

Policy # 00751

Original Effective Date: 08/09/2021

Current Effective Date: 08/01/2026

Patients were on a background of other lipid-lowering therapies, including maximally tolerated statins, ezetimibe, PCSK9 inhibitor antibodies, lomitapide, and lipoprotein apheresis. The mean LDL-C at baseline was 255 mg/dL. In patients with limited LDLR function, the mean LDL-C at baseline was 307 mg/dL. At baseline, 94% of patients were on statins, 75% on ezetimibe, 77% on a PCSK9 inhibitor antibody, 22% on lomitapide, and 34% were receiving lipoprotein apheresis. The mean age at baseline was 42 years (range 12 to 75) with 12% $\geq$ 65 years old; 54% women, 3% Hispanic, 74% White, 15% Asian, 3% Black, and 8% Other or not reported.

The primary efficacy endpoint was percent change in LDL-C from baseline to Week 24. At Week 24, the least squares (LS) mean treatment difference between Evkeeza and placebo in mean percent change in LDL-C from baseline was -49% (95% confidence interval: -65% to -33%; $p < 0.0001$). After 24 weeks of open-label Evkeeza treatment (Week 24 to Week 48), the observed LDL-C reduction from baseline was similar in patients who cross over from placebo to Evkeeza and was maintained in patients who remained on Evkeeza for 48 weeks.

In an open-label trial (Trial 2), 13 pediatric patients with HoFH (aged 12 to 17 years) received 15 mg/kg of Evkeeza given intravenously every 4 weeks as an adjunct to other lipid-lowering therapies (e.g., statins, ezetimibe, PCSK9 inhibitor antibodies and lipoprotein apheresis) for a median treatment duration of 33 weeks. The mean percent change from baseline in LDL-C at Week 24 was -52% in the 9 patients who completed treatment and had a lipid assessment at Week 24. Overall, the effect of Evkeeza on lipid parameters in pediatric patients aged 12 to 17 years with HoFH was generally similar to that seen in adults with HoFH.

Trial R1500-CL-17100 (Trial 3) was a multicenter, three-part, single-arm, open-label trial in pediatric patients aged 5 to 11 years with HoFH. Part B of this trial evaluated the efficacy of Evkeeza 15 mg/kg given intravenously every 4 weeks as an adjunct to other lipid-lowering therapies (e.g., statins, ezetimibe, lomitapide, and lipoprotein apheresis) for 24 weeks in 14 patients with HoFH.

In Part B, the mean LDL-C at baseline was 264 mg/dL. The primary efficacy endpoint was percent change in calculated LDL-C from baseline to Week 24. At Week 24, the mean percent change in calculated LDL-C from baseline was -48% (95% confidence interval: -69% to -28%).

Clinical data for pediatric patients aged 1 to less than 5 years with HoFH were collected through an expanded access program. The program required each patient's treating physician to provide documentation confirming eligibility and effectiveness information, including LDL-C values. The efficacy of Evkeeza was evaluated in 6 patients aged 1 to less than 5 years with HoFH. Four females and two males were enrolled. At baseline, 83% of patients were receiving statins, 83% ezetimibe, 17% PCSK9 inhibitors, and 17% plasmapheresis. Mean baseline LDL-C was 499 mg/dL (range 238 to 872 mg/dL), mean age was 3.2 years (range 1.1 to 4.4 years), and mean weight was 14.5 kg (range 8 to 17 kg). Patients received Evkeeza 15 mg/kg every 4 weeks for up to 98 weeks. Treatment with Evkeeza resulted in a mean LDL-C reduction of 60% at Week 40 [mean (standard deviation) change of -60% (16.8)].

Policy # 00751

Original Effective Date: 08/09/2021

Current Effective Date: 08/01/2026

References

1. Evkeeza [package insert]. Regeneron Pharmaceuticals, Inc. Tarrytown, New York. Updated September 2025.
2. Stone NJ, Robinson J, Lichtenstein AH, et al. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice guidelines. *Circulation*. 2014;129(25 Suppl 2):S1-S45. Available at <http://circ.ahajournals.org/content/early/2013/11/11/01.cir.0000437738.63853.7a>.
3. Jacobson TA, Ito MK, Maki KC, et al. National Lipid Association recommendations for patient-centered management of dyslipidemia: Part 1-executive summary. *J Clin Lipidol*. 2014;8:473-488. Available at: [http://www.lipidjournal.com/article/S1933-2874\(14\)00274-8/pdf](http://www.lipidjournal.com/article/S1933-2874(14)00274-8/pdf).
4. Jacobson TA, Ito MK, Maki KC, et al. National Lipid Association recommendations for patient-centered management of dyslipidemia: Part 1-full report. *J Clin Lipidol*. 2015;9:129-169.
5. Rosenson RS, Baker SK, Jacobson TA, et al. An assessment by the statin muscle safety task force: 2014 update. *J Clin Lipidol*. 2014;8:S58-S71.
6. Guyton JR, Bays HE, Grundy SM, Jacobson TA. An assessment by the Statin Intolerance Panel: 2014 update. *J Clin Lipidol*. 2014;8:S72-S81.
7. Moriarty PM, Jacobson TA, Bruckert E, et al. Efficacy and safety of alirocumab, a monoclonal antibody to PCSK9, in statin-intolerant patients: design and rationale of ODYSSEY ALTERNATIVE, a randomized phase 3 trial. *J Clin Lipidol*. 2014;8:554-561.
8. Zhang H, Plutzky J, Skentzos S, et al. Discontinuation of statins in routine care settings. *Ann Intern Med*. 2013;158(7):526-534.
9. Mampuya WM, Frid D, Rocco M, et al. Treatment strategies in patients with statin intolerance: the Cleveland Clinic Experience. *Am Heart J*. 2013;166(3):597-603.
10. 95th Percentile Chart: Chapter 31: Cholesterol, Triglycerides, and Associated Lipoproteins; In *Clinical Methods: The History, Physical, and Laboratory Examinations*. 3rd edition. Walker HK, Hall WD, Hurst JW, et al. Available at: <http://www.ncbi.nlm.nih.gov/books/NBK351/table/A968/?report=objectonly>.
11. Alternative percentile link: ATP III Final Report Appendix III-A Distributions of Total Cholesterol, LDL Cholesterol, HDL Cholesterol, and Triglycerides in the U.S. Adult Population, NHANES III Data (1988-1994). Available at: <http://circ.ahajournals.org/content/106/25/3237/T2.expansion.html>.
12. Steg G, Schwartz GG, Szarek M, et al, on behalf of the ODYSSEY Outcomes Investigators and Committees. The ODYSSEY Outcomes trial: topline results. Alirocumab in patients after acute coronary syndrome. Presented at: the American College of Cardiology 67th Scientific Sessions; Orlando, FL; March 10-12, 2018. Presented March 10, 2018. Available at: <http://www.acc.org/educationand-meetings/image-and-slide-gallery/media-detail?id=90e30055844c402ba16d54fb05e45136>.
13. Schwartz GG, Bessac L, Berdan LG, et al. Effect of alirocumab, a monoclonal antibody to PCSK9, on long-term cardiovascular outcomes following acute coronary syndromes: rationale and design of the ODYSSEY Outcomes trial. *Am Heart J*. 2014;168:682-689.e1.
14. Sabatine MS, Giugliano RP, Keech A, et al. Evolocumab and clinical outcomes in patients with cardiovascular disease. *N Engl J Med*. 2017;376(18):1713-1722.

evinacumab-dgnb (Evkeeza™)

Policy # 00751

Original Effective Date: 08/09/2021

Current Effective Date: 08/01/2026

Policy History

Original Effective Date: 08/09/2021

Current Effective Date: 08/01/2026

07/01/2021 Medical Policy Committee review

07/14/2021 Medical Policy Implementation Committee approval. New policy.

09/30/2021 Coding update

07/07/2022 Medical Policy Committee review

07/13/2022 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

07/06/2023 Medical Policy Committee review

07/12/2023 Medical Policy Implementation Committee approval. Updated the age for use from 12 years to 5 years to align with the FDA approval update.

07/02/2024 Medical Policy Committee review

07/10/2024 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

07/03/2025 Medical Policy Committee review

07/09/2025 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

07/21/2026 Pre Pharmacy and Therapeutics Committee review

07/22/2026 Pharmacy and Therapeutics Committee approval. Updated age according to FDA label update (approved in ages 1 and up). Updated LDL-C treatment goals to align with latest guideline recommendations. Added Policy Guidelines section.

Next Scheduled Review Date: 07/2027

Coding

The five character codes included in the Louisiana Blue Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)‡, copyright 2025 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Louisiana Blue Medical Policy Coverage Guidelines is with Louisiana Blue and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Louisiana Blue Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Louisiana Blue Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Policy # 00751

Original Effective Date: 08/09/2021

Current Effective Date: 08/01/2026

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	No codes
HCPCS	J1305
ICD-10 Diagnosis	All Related Diagnoses

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
 1. Consultation with technology evaluation center(s);
 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
 3. Reference to federal regulations.

**Medically Necessary (or “Medical Necessity”) - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

For these purposes, “nationally accepted standards of medical practice” means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

evinacumab-dgnb (Evkeeza™)

Policy # 00751

Original Effective Date: 08/09/2021

Current Effective Date: 08/01/2026

NOTICE: If the Patient's health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

NOTICE: Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage.

NOTICE: If an authorization for an ongoing course of treatment has been provided to a member and the member changes from one health plan to another health plan (e.g., a member moves from carrier A to Louisiana Blue), Louisiana Blue may honor the previous health plan's authorization for the same service under the same type of in-network benefit for a 90-day transition period. Documentation of the authorization for the ongoing course of treatment from the previous health plan must be provided to us by the member or their provider and the services provided for the course of treatment must otherwise be a covered service under the Louisiana Blue health plan. This provision does not apply to medications covered under the plan's pharmacy benefit.